




3/16" X 3/16" ARRAY ALPHA NUMERIC DATA MATRIX CODE (ECC 200) PRINTED IN BLACK OR DARK BLUE ON WHITE FIELD WITH 1/8" QUIET ZONE ON ALL 4 SIDES
DATA MATRIX CODE = 083398

PRODUCTION (NOT COLOR ACCURATE)



DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



Infants' Advil
Concentrated Drops
Ibuprofen Oral Suspension
50 mg per 1.25 mL
Pain Reliever/Fever Reducer (NSAID)

White Grape-Flavored Drops
1 FL OZ (30 mL)

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Active ingredient Ibuprofen 50 mg in each 1.25 mL. Uses temporarily reduces fever, relieves minor aches and pains due to the common cold, flu, headaches and toothaches.

SHAKE WELL

Do Not Use if plastic wrap imprinted with "SAFETY SEAL" is broken or missing.

Warnings
Ask doctor before use if child is under a doctor's care for a serious condition, taking other drugs or has stomach problems.

TOP PANEL 1



Directions Does not contain directions or complete warnings for adult use. Do not give more than directed. Shake well before using. Find right dose on chart. Use weight to dose; otherwise use age. Repeat dose every 6-8 hours, if needed. Do not use more than 4 times a day. Measure with the dosing device provided.

Weight (lb)	Age (mos)	Dose (mL)
under 6 mos		ask a doctor
12-17 lb	6-11 mos	1.25 mL
18-23 lb	12-23 mos	1.875 mL

Questions or comments?
call 1-800-88-Advil
Pfizer, Madison, NJ 07940 USA
For most recent product information, visit www.Advil.com.
© 2016 Pfizer Inc. PAA083399

BASE PANEL 3

UNVARNISHED AREA

VARNISHED 1/8" x 1/8" DATA MATRIX CODE (ECC 200), IN BLACK OR DARK BLUE.
DATA MATRIX CODE = 083399



May cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If allergic reaction occurs, stop use, seek medical help right away. Do not use if child has ever had an allergic reaction to any pain reliever/fever reducer. Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if the child has had stomach ulcers or bleeding problems, takes a blood thinning (anticoagulant) or steroid drug, takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), takes more or for a longer time than directed. Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

RELEASED BACK PANEL 2

5/16" HINGE AREA UNVARNISHED

PRODUCTION

(NOT COLOR ACCURATE)



DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	6 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.25 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• WARNING BOX LINE, HAIRLINES	.5 pt




3/16" X 3/16" ARRAY ALPHA NUMERIC DATA MATRIX CODE (ECC 200) PRINTED IN BLACK OR DARK BLUE ON WHITE FIELD WITH 1/8" QUIET ZONE ON ALL 4 SIDES
DATA MATRIX CODE = 082444

PRODUCTION (NOT COLOR ACCURATE)



(b) (4)

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



Infants' Advil
Concentrated Drops
Ibuprofen Oral Suspension
50 mg per 1.25 mL
Pain Reliever/Fever Reducer (NSAID)

White Grape-Flavored Drops
3.4 FL OZ (22.5 mL)

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Active ingredient Ibuprofen 50 mg in each 1.25 mL.
Uses: temporarily reduces fever, relieves minor aches and pains due to the common cold, flu, headaches and toothaches.

SHAKE WELL

Do Not Use if plastic wrap imprinted with "SAFETY SEAL" is broken or missing.

Warnings:
Ask doctor before use if child is under a doctor's care for a serious condition, taking other drugs or has stomach problems.

LET
HERE

**TOP
PANEL 1**



Directions: Does not contain directions or complete warnings for adult use. Do not give more than directed. Shake well before using. Find right dose on chart. Use weight to dose; otherwise use age. Repeat dose every 6-8 hours, if needed. Do not use more than 4 times a day. Measure with the dosing device provided.

Weight (lb)	Age (mos)	Dose (mL)
under 6 mos		ask a doctor
12-17 lb	6-11 mos	1.25 mL
18-23 lb	12-23 mos	1.875 mL

Pfizer, Madison, NJ 07940 USA
For most recent product information, visit www.Advil.com.
© 2016 Pfizer Inc. PAA082445

Questions or comments? call 1-800-88-Advil

**BASE
PANEL 3**

UNVARNISHED AREA

VARNISHED 1/8" x 1/8"
DATA MATRIX CODE (ECC 200),
IN BLACK OR DARK BLUE.
DATA MATRIX CODE = 082445



May cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If allergic reaction occurs, stop use, seek medical help right away. Do not use if child has ever had an allergic reaction to any pain reliever/fever reducer. **Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if the child has had stomach ulcers or bleeding problems, takes a blood thinning (anticoagulant) or steroid drug, takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), takes more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**RELEASED BACK
PANEL 2**

5/16" HINGE AREA UNVARNISHED

PRODUCTION

(NOT COLOR ACCURATE)



DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	5.45 pt
• SUBHEADINGS/BODY TEXT	5.45 pt
• LEADING	5.95 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• WARNING BOX LINE, HAIRLINES	.5 pt



3/16" X 3/16" ARRAY ALPHA NUMERIC DATA MATRIX CODE (ECC 200) PRINTED IN BLACK OR DARK BLUE ON WHITE FIELD WITH 1/8" QUIET ZONE ON ALL 4 SIDES
DATA MATRIX CODE = 082315

PRODUCTION

(NOT COLOR ACCURATE)



DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces






3/16" X 3/16" ARRAY ALPHA NUMERIC DATA MATRIX CODE (ECC 200) PRINTED IN BLACK OR DARK BLUE ON WHITE FIELD WITH 1/8" QUIET ZONE ON ALL 4 SIDES
DATA MATRIX CODE = 082451



PRODUCTION (NOT COLOR ACCURATE)



DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



BASE



RELEASED BACK

PRODUCTION

(NOT COLOR ACCURATE)



DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	5.45 pt
• SUBHEADINGS/BODY TEXT	5.45 pt
• LEADING	5.8 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• WARNING BOX LINE, HAIRLINES	.5 pt







090177e18afa5399\Final\Final On: 03-Jan-2017 16:10

PRODUCTION

(NOT COLOR ACCURATE)

(b) (4)





PRODUCTION

(NOT COLOR ACCURATE)



DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	5.45 pt
• SUBHEADINGS/BODY TEXT	5.45 pt
• LEADING	5.8 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• WARNING BOX LINE, HAIRLINES	.5 pt

PEEL HERE
AREA

LIFT HERE
for easy
drug facts

TEXT FREE AREA
1/16"

COPY AREA
PANEL
1

Drug Facts

Active ingredient (in each 1.25 mL) **Purpose**
Ibuprofen 50 mg (NSAID) Fever reducer/Pain reliever
*nonsteroidal anti-inflammatory drug

Uses temporarily:

- reduces fever
- relieves minor aches and pains due to the common cold, flu, headaches and toothaches

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives ■ facial swelling ■ asthma (wheezing)
- shock ■ skin reddening ■ rash ■ blisters

 If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if the child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

3/16" x 3/16" DATA MATRIX CODE (ECC200), PRINTED IN BLACK OR DARK BLUE ON UNPRINTED WHITE BACKGROUND

DATA MATRIX CODE=078966

COPY AREA
PANEL
2

Drug Facts (continued)

Do not use

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to the child
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has a history of stomach problems, such as heartburn
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
 - feels faint
 - vomits blood
 - has bloody or black stools
 - has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

COPY AREA
BASE
PANEL

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- repeat dose every 6-8 hours, if needed
- do not use more than 4 times a day
- measure with the dosing device provided. Do not use with any other device.

Dosing Chart

Weight (lb)	Age (mos)	Dose (mL)
under 6 mos		ask a doctor
12-17 lb	6-11 mos	1.25 mL
18-23 lb	12-23 mos	1.875 mL

Other Information

- one dose lasts 6-8 hours
- store at 20-25°C (68-77°F)
- see bottom of box for lot number and expiration date

Inactive ingredients
 acetic acid, artificial flavor, butylated hydroxytoluene, carboxymethylcellulose sodium, citric acid monohydrate, edetate disodium, glycerin, microcrystalline cellulose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum

Questions or comments?
 Call toll-free 1-800-88-ADVIL or ask your pharmacist, doctor or health care professional

PAA078966.FDA01

ADHESIVE
AREAS 1/8"

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

PRODUCTION

(NOT COLOR ACCURATE)

(b) (4)



This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

VALERIE S PRATT
03/24/2017