

4.75" width

Front

(b) (4)

Cat. No. 260480 / 260480NSB NDC 054365-400-06

Chloraprep® One-Step

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact
1 ml APPLICATOR



Not made with natural rubber latex



Do Not Reuse

Professional Use Only
External Use Only

Drug Facts	Purposes
Active ingredients	
Chlorhexidine gluconate 2% w/v.....	Antiseptic
Isopropyl alcohol 70% v/v.....	Antiseptic
Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.	
Warnings	
For external use only. Flammable, keep away from fire or flame.	
<ul style="list-style-type: none"> ■ do not use with electrocautery procedures 	
Allergy Alert	
This product may cause a severe allergic reaction. Symptoms may include:	
<ul style="list-style-type: none"> ■ wheezing/difficulty breathing ■ shock ■ facial swelling ■ hives ■ rash 	
If an allergic reaction occurs, stop use and seek medical help right away.	
Do not use	
<ul style="list-style-type: none"> ■ on patients allergic to chlorhexidine gluconate or any other ingredient in this product ■ for lumbar puncture or in contact with the meninges ■ on open skin wounds or as a general skin cleanser 	

3.8" length

Chloraprep® One-Step

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

Drug Facts (continued)
When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.
Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.
Directions
<ul style="list-style-type: none"> ■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. ■ maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm²) ■ remove applicator from package; do not touch sponge ■ hold the applicator with the sponge down. Pinch wings <u>only once</u> to activate the ampule and release the antiseptic. ■ wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin ■ completely wet the treatment area with antiseptic ■ dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for approximately 30 seconds. Allow the area to air dry for approximately 30 seconds. ■ moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes for approximately 2 minutes. Allow the area to air dry for approximately 1 minute. ■ do not blot or wipe away ■ discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.
Other information
<ul style="list-style-type: none"> ■ store between 15–30 °C (59–86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F)
Inactive ingredient USP purified water
Questions?
<ul style="list-style-type: none"> ■ www.chloraprep.com ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

CareFusion, El Paso, TX 79912 • Artwork 6-775480 • EXXXXXX • Made in the USA of US and imported parts

Back

Artwork No. 6-001480, ChloroPrep One-Step 1 mL

(b) (4)


(b) (4)

(b) (4)




Exp. Date and Lot No.: placement - for manufacturing purpose only.

Reference ID: 4151410



Chloraprep® One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact



Drug Facts	Purposes
Active ingredients: Chlorhexidine gluconate 2% w/v Isopropyl alcohol 70% v/v	Antiseptic Antiseptic
Uses: For the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.	
Warnings: For external use only. Flammable, keep away from fire or flame. Do not use with electrocautery procedures.	
Allergy Alert: This product may cause a severe allergic reaction. Symptoms may include: • wheezing/difficulty breathing • facial swelling • hives • rash If an allergic reaction occurs, stop use and seek medical help right away.	
Do not use: • on patients allergic to chlorhexidine gluconate or any other ingredient in this product • on similar particles or in contact with the meninges • on open skin wounds or as a general skin cleanser	
When using this product: Keep out of eyes, ears, and mouth. May cause redness or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor. Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
Directions: • Use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. • Maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (62 cm²) • Remove applicator from package; do not touch sponge • Hold the applicator with the sponge down. Pinch wings only, close to activate the ampule and release the antiseptic. • Wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin. • Completely wet the treatment area with antiseptic. • Dry applied area (e.g., abdomen or arm); use gentle repeated back-and-forth strokes of the sponge for 30 seconds. Allow the area to air dry for approximately 30 seconds. • moist surgical sites (e.g., abdominal or arm); use gentle repeated back-and-forth strokes of the sponge for 2 minutes. Allow the area to air dry for approximately 1 minute. • Do not use applicator on face. • Discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.	
Other information: • Store between 15-30 °C (59-86 °F) • Avoid freezing and excessive heat above 40 °C (104 °F)	
Inactive ingredient: USP purified water	
Questions? • www.chloraprep.com • toll 1-800-523-0502 (M-F 8 a.m.-5 p.m. CST)	

Chloraprep® One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact



1 ml
APPLICATORS
Clear

1 ml
APPLICATORS
Clear

60 applicators

CareFusion
El Paso, TX 79912
1-800-523-0502
www.chloraprep.com
Made in the USA of US and imported parts


1 ml
APPLICATORS
Clear

Do Not Reuse
Not made with natural rubber latex
STERILE

Professional Use Only
Store between 15-30 °C (59-86 °F)
Avoid freezing and excessive heat above 40 °C (104 °F)

60 applicators
0.03 fl. oz (1 ml) each

Chloraprep® One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact



Chloraprep® One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact



1 ml
APPLICATORS
Clear

LOT XXXXXX (printed in-line) EXP DATE XXXXX
No Yarnish

BARCODE PLACEHOLDER

©2014 CareFusion. All rights reserved. Chloraprep® is a registered trademark of CareFusion Corporation or one of its subsidiaries.

No Yarnish / No printing area

1 15/32

3 1/4

1 7/4

6 1/2

3 1 1/16

2 5 1/6 4

1 2 7/8

5 5/8

1 2 7/8

5 1 9/32

3 7 9/32

(b) (4)

Dimensions: 7.5" x 2.5"

← 2.5" →

ChloroPrep® One-Step
 2% w/v chlorhexidine gluconate (CHG)
 and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation

Non-sterile Solution
 Applicator is sterile if package is intact
 1.5 ml FREPP® APPLICATOR

Drug Facts						
<table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%;">Active ingredients</td> <td style="width: 50%;">Purposes</td> </tr> <tr> <td>Chlorhexidine gluconate 2% w/v</td> <td>Antiseptic</td> </tr> <tr> <td>Isopropyl alcohol 70% v/v</td> <td>Antiseptic</td> </tr> </table>	Active ingredients	Purposes	Chlorhexidine gluconate 2% w/v	Antiseptic	Isopropyl alcohol 70% v/v	Antiseptic
Active ingredients	Purposes					
Chlorhexidine gluconate 2% w/v	Antiseptic					
Isopropyl alcohol 70% v/v	Antiseptic					
Use For the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.						
Warnings For external use only Flammable, keep away from fire or flame. <ul style="list-style-type: none"> ■ do not use with electrocautery procedures 						
Allergy Alert This product may cause a severe allergic reaction. Symptoms may include: <ul style="list-style-type: none"> ■ wheezing/difficulty breathing ■ shock ■ facial swelling ■ hives ■ rash If an allergic reaction occurs, stop use and seek medical help right away.						
Do not use <ul style="list-style-type: none"> ■ on patients allergic to chlorhexidine gluconate or any other ingredient in this product ■ for lumbar puncture or in contact with the meninges ■ on open skin wounds or as a general cleanser 						
When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.						
Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.						
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.						
Directions <ul style="list-style-type: none"> ■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. ■ maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm²) ■ remove applicator from package; do not touch sponge ■ hold the applicator horizontally with the sponge down. Pinch wings only once to activate the ampule and release the antiseptic. ■ wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin ■ completely wet the treatment area with antiseptic ■ dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for approximately 30 seconds. Allow the area to air dry for approximately 30 seconds. ■ moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes for approximately 2 minutes. Allow the area to air dry for approximately 1 minute. ■ do not blot or wipe away ■ discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available. 						
Other information ■ store between 15–30 °C (59–86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F)						
Inactive ingredient USP purified water						
Questions? ■ www.chloroprep.com ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)						

See other side for additional information

Front

NDC 054365-400-02
 Cat. No. 260299NSB / 260599NSB

ChloroPrep® One-Step
 2% w/v chlorhexidine gluconate (CHG)
 and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation

Non-sterile Solution
 Applicator is sterile if package is intact
 1.5 ml FREPP® APPLICATOR

See Drug Facts for Full Labeling Information

 Do Not Reuse

 Not made with natural rubber latex

External Use Only
Professional Use Only

CareFusion, El Paso, TX 79912

Artwork 6-775403F · PXXXXXX
 Made in the USA of US and imported parts

Back

(b) (4)

Cat. No. 260299 NDC 054365-400-02

ChloroPrep® One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation

Non-sterile Solution
Applicator is sterile if package is intact

1.5 ml FREPP® APPLICATOR

WARNING. FLAMMABLE.
KEEP AWAY FROM FIRE OR FLAME.

CareFusion, El Paso, TX 79912
1-800-523-0502 (M-F 8 a.m. - 5 p.m. CST)

Artwork No. 6-000299 · M090314
Made in the USA of US and imported parts

Exp. Date xx/xx
Lot No. xxxxx

Do Not Reuse

Bend and Peel

72 mm

40 mm

Trim Area

(b) (4)

(b) (4)

66 mm

40 mm

Placeholder for lot and expiry data

Cat. No. 260299 NDC 054365-400-02
ChloroPrep® One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact
1.5 ml FREPP® APPLICATOR



WARNING. FLAMMABLE.
KEEP AWAY FROM FIRE OR FLAME.

CareFusion, El Paso, TX 79912
1-800-523-0502 (M-F 8 a.m. - 5 p.m. CST)
Artwork No. 6-001299 · F090314
Made in the USA of US and imported parts


Do Not Reuse



Bend and Peel

Exp. Date and Lot No.: placement - for
manufacturing purpose only.

Artwork No. 6-824299, ChloroPrep One-Step Frepp Intermediate Carton



(b) (4)

4.75" width

Front


(b) (4)

<div style="background-color: red; color: white; text-align: center; padding: 2px;">WARNING</div> <div style="background-color: red; color: white; text-align: center; padding: 2px;">FLAMMABLE</div> <p>Keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:</p> <ul style="list-style-type: none"> • solution contains alcohol and gives off flammable vapors • avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable. • do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) • do not allow solution to pool • remove wet materials from prep area 	<p style="font-size: small;">Cat. No. 260415 / 260415NS / 260415NSB NDC 054365-400-11 Cat. No. 260400 / 260400NS / 260500NS NDC 054365-400-01</p> <p style="text-align: center;">Chloraprep® One-Step Chloraprep® With Tint</p> <p style="text-align: center; font-size: small;">2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)</p> <p style="text-align: center;">Patient Preoperative Skin Preparation</p> <p style="text-align: center; font-size: small;">Non-sterile Solution Applicator is sterile if package is intact</p> <p style="text-align: center; font-size: large;">3 ml APPLICATOR</p> <p style="font-size: x-small;">Professional Use Only External Use Only Do Not Reuse Not made with natural rubber latex</p>																								
<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <th colspan="2" style="text-align: left; padding: 2px;">Drug Facts</th> </tr> <tr> <td style="width: 60%; padding: 2px;">Active ingredients</td> <td style="width: 40%; padding: 2px;">Purposes</td> </tr> <tr> <td style="padding: 2px;">Chlorhexidine gluconate 2% w/v.....</td> <td style="padding: 2px;">.....Antiseptic</td> </tr> <tr> <td style="padding: 2px;">Isopropyl alcohol 70% v/v.....</td> <td style="padding: 2px;">.....Antiseptic</td> </tr> <tr> <td colspan="2" style="padding: 2px;">Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.</td> </tr> <tr> <td colspan="2" style="padding: 2px;">Warnings</td> </tr> <tr> <td colspan="2" style="padding: 2px;">For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:</td> </tr> <tr> <td colspan="2" style="padding: 2px;"> <ul style="list-style-type: none"> ■ solution contains alcohol and gives off flammable vapors ■ avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable. ■ do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) ■ do not allow solution to pool ■ remove wet materials from prep area </td> </tr> <tr> <td colspan="2" style="padding: 2px;">Allergy Alert</td> </tr> <tr> <td colspan="2" style="padding: 2px;">This product may cause a severe allergic reaction. Symptoms may include:</td> </tr> <tr> <td colspan="2" style="padding: 2px;"> <ul style="list-style-type: none"> ■ wheezing/difficulty breathing ■ shock ■ facial swelling ■ hives ■ rash </td> </tr> <tr> <td colspan="2" style="padding: 2px;">If an allergic reaction occurs, stop use and seek medical help right away. ▶</td> </tr> </table>		Drug Facts		Active ingredients	Purposes	Chlorhexidine gluconate 2% w/v.....Antiseptic	Isopropyl alcohol 70% v/v.....Antiseptic	Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.		Warnings		For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:		<ul style="list-style-type: none"> ■ solution contains alcohol and gives off flammable vapors ■ avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable. ■ do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) ■ do not allow solution to pool ■ remove wet materials from prep area 		Allergy Alert		This product may cause a severe allergic reaction. Symptoms may include:		<ul style="list-style-type: none"> ■ wheezing/difficulty breathing ■ shock ■ facial swelling ■ hives ■ rash 		If an allergic reaction occurs, stop use and seek medical help right away. ▶	
Drug Facts																									
Active ingredients	Purposes																								
Chlorhexidine gluconate 2% w/v.....Antiseptic																								
Isopropyl alcohol 70% v/v.....Antiseptic																								
Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.																									
Warnings																									
For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:																									
<ul style="list-style-type: none"> ■ solution contains alcohol and gives off flammable vapors ■ avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable. ■ do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) ■ do not allow solution to pool ■ remove wet materials from prep area 																									
Allergy Alert																									
This product may cause a severe allergic reaction. Symptoms may include:																									
<ul style="list-style-type: none"> ■ wheezing/difficulty breathing ■ shock ■ facial swelling ■ hives ■ rash 																									
If an allergic reaction occurs, stop use and seek medical help right away. ▶																									


3.8" length

<p>Chloraprep® One-Step / Chloraprep® With Tint</p> <p>2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)</p>
<p>Drug Facts (continued)</p> <p>Do not use</p> <ul style="list-style-type: none"> ■ on patients allergic to chlorhexidine gluconate or any other ingredient in this product ■ for lumbar puncture or in contact with the meninges ■ on open skin wounds or as a general skin cleanser <p>When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.</p> <p>Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p> <p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> ■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. ■ use in a well ventilated area ■ maximal treatment area for one applicator is approximately 4 in. x 5 in. (130 cm²) ■ remove applicator from package; do not touch sponge ■ hold the applicator with the sponge down. Pinch wings only once to activate the ampule and release the antiseptic. ■ wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin ■ completely wet the treatment area with antiseptic ■ dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes of the sponge for approximately 30 seconds. Allow solution to air dry for approximately 30 seconds. Do not blot or wipe away. ■ moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes of the sponge for approximately 2 minutes. Allow solution to air dry for approximately 1 minute. Do not blot or wipe away. ■ if using an ignition source, allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away. ■ discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available. <p>Other information ■ store between 15–30 °C (59–86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F) ■ the tint will slowly fade from the skin. Soap and water, or alcohol may be used to remove the tint if desired.</p> <p>Inactive ingredients FD&C yellow #6 dye (Hi-Lite Orange applicators only), USP purified water</p> <p>Questions? ■ www.chloraprep.com ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)</p>
<p>CareFusion, El Paso, TX 79912 · Artwork 6-775400 · IXXXXXX · Made in the USA of US and imported parts</p>

Back




Lot/Exp Placeholder - for reference only.



(b) (4)

Lot/Exp Placeholder -
for reference only.

1¹⁵/₃₂



ChloroPrep® One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact

WARNING, FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts	Purposes
Active ingredients Chlorhexidine gluconate 2% w/v Isopropyl alcohol 70% v/v	Antiseptic Anesthetic
Use For the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.	
Warnings For external use only. Flammable, keep away from fire or flame. To reduce the risk of fire, PREP CAREFULLY: • solution contains alcohol and gives off flammable vapors • avoid getting solution into hair areas that may take up to 1 hour to dry. Wet hair is flammable. • do not drag or use ignition sources (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) • do not allow solution to pool • remove wet materials from prep area	
Allergy Alert: This product may cause a severe allergic reaction. Symptoms may include: • wheezing/difficulty breathing • shock • hives • itching • rash If an allergic reaction occurs, stop use and seek medical help right away.	
Do not use • on patients allergic to chlorhexidine gluconate or any other ingredient in this product • on umbilical puncture or in contact with the meninges • on open skin wounds or as a general skin cleanser	
When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor. Stop use and seek a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
Directions • use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. • use in a well-ventilated area • use on treatment area for one application at approximately 4" x 6" in (130 cm ²) • remove applicator from package. Do not touch sponge. • hold the applicator with the sponge side. Firmly single stroke to activate the sponge and release the antiseptic. • wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin. • use on the treatment area with antiseptic. • dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes of the sponge for approximately 30 seconds. Allow solution to air dry for approximately 30 seconds. Do not blot or wipe away. • moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes of the sponge for approximately 2 minutes. Allow solution to air dry for approximately 1 minute. Do not blot or wipe away. • flaming all surgical sites: allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away. • use the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.	
Other information • store between 15-30 °C (59-86 °F) • avoid freezing and excessive heat above 40 °C (104 °F)	
Inactive ingredient USP purified water	
Questions? www.chloroprep.com 1-800-523-0592 M-F 9 a.m.-5 p.m. CST	

ChloroPrep® One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact

3 ml
APPLICATORS
Clear

25 applicators

ChloroPrep® One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact

3 ml
APPLICATORS
Clear

CareFusion
El Paso, TX 79912
1-800-523-0592
www.chloroprep.com
Made in the USA of US and imported parts

3 ml
APPLICATORS
Clear

Do Not Reuse
Not made with natural rubber latex
STERILE EO

25 applicators
0.10 fl. oz. (3 ml) each

ChloroPrep® One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact

WARNING
FLAMMABLE
Keep away from fire or flame.
To reduce risk of fire, PREP CAREFULLY:
• solution contains alcohol and gives off flammable vapors
• avoid getting solution into hair areas
• hair may take up to 1 hour to dry. **Wet hair is flammable.**
• do not drag or use ignition sources (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
• do not allow solution to pool
• remove wet materials from prep area

ChloroPrep® One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact

3 ml
APPLICATORS
Clear

LOT XXXXXX (Printed in-line)
No Yarnish
EXP DATE XXXXX
BARCODE PLACEHOLDER

No Yarnish/No printing area

3¹¹/₁₆

1¹/₁₆

1 2⁷/₈

5⁵/₈

1 2⁷/₈

5¹⁹/₃₂

3 7⁹/₃₂

3³/₄

6¹/₂

2⁵/₁₆

1 7¹/₄

1¹⁵/₃₂

3¹¹/₁₆

1¹¹/₁₆

1 2⁷/₈

5⁵/₈

1 2⁷/₈

5¹⁹/₃₂


3 7⁹/₃₂

2⁵/₁₆

6¹/₂

1 7¹/₄

3³/₄



ChloroPrep® With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact

WARNING, FLAMMABLE, KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use For the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

- For external use only. Flammable, keep away from fire or flame. To reduce the risk of fire, PREP CAREFULLY!
- Solution contains alcohol and gives off flammable vapors.
- Avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable.
- Do not scrape or use lotion source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair).
- Do not allow solution to pool.
- Remove wet materials from prep area.

Caution: This product may cause a severe allergic reaction. Symptoms may include:

- Intermittently breathing
- Hives
- Facial swelling
- Itching
- Redness

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- On patients allergic to chlorhexidine gluconate or any other ingredient in this product.
- For lacerations or in contact with the meninges.
- On open skin wounds or as a general skin cleanser.

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- Use in a well-ventilated area.
- Maximal treatment area for one applicator is approximately 4 in. x 5 in. (100 cm²).
- Remove applicator from package; do not touch sponge.
- Hold the applicator with the sponge down. From angle 095°/205° to activate the sponge and release the antiseptic.
- Wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin.
- Continuously use the treatment area with strokes.
- Apply surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes of the sponge for approximately 30 seconds. Allow solution to air dry for approximately 30 seconds. Do not blot or wipe away.
- Moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes of the sponge for approximately 2 minutes. Allow solution to air dry for approximately 1 minute. Do not blot or wipe away.
- When using an applicator, allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- Discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- Store between 15-30 °C (59-86 °F).
- Avoid freezing and excessive heat above 40 °C (104 °F).

Inactive ingredients FDAC yellow #6 dye, USP purified water.

Questions?

- www.chloroprep.com
- 1-800-523-0592 M-F 8 a.m.-6 p.m. CST.

ChloroPrep® With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact

3 ml
APPLICATORS
Hi-Lite Orange®

25 applicators

ChloroPrep® With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact

3 ml
APPLICATORS
Hi-Lite Orange®

CareFusion
El Paso, TX 79912
1-800-523-0592
www.chloroprep.com
Made in the USA of US and imported parts

3 ml
APPLICATORS
Hi-Lite Orange®

Do Not Reuse
Not made with natural rubber latex
STERILE EO

Professional Use Only
Store between 15-30 °C (59-86 °F)
Avoid freezing and excessive heat above 40 °C (104 °F)

25 applicators
0.10 fl. oz. (3 ml) each

ChloroPrep® With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact

**WARNING
FLAMMABLE**
Keep away from fire or flame.
To reduce risk of fire, PREP CAREFULLY:
- solution contains alcohol and gives off flammable vapors
- avoid getting solution into hairy areas
- hair may take up to 1 hour to dry. Wet hair is flammable.
- do not scrape or use lotion source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

ChloroPrep® With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact

3 ml
APPLICATORS
Hi-Lite Orange®

LOT XXXXXX (Printed in Red) EXP DATE XXXX
No Variants / No printing area

©2009 CareFusion. All rights reserved. ChloroPrep® is a registered trademark of CareFusion Corporation or one of its subsidiaries.

Artwork No. 6-824415 OXXXXXX

NDC 054365-400-11
Cat. No. 260415

4.75" width

Front

3.8" length


⚠ WARNING FLAMMABLE	Cat. No. 260700 / 260700NS / 260700NSB NDC 054365-400-04 Cat. No. 260715 / 260715NS / 260715NSB NDC 054365-400-12 Cat. No. 260725 / 260725NS / 260725NSB NDC 054365-400-09						
<p>Keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:</p> <ul style="list-style-type: none"> • solution contains alcohol and gives off flammable vapors • avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable. • do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) • do not allow solution to pool • remove wet materials from prep area 	<p>ChloroPrep® One-Step ChloroPrep® With Tint</p> <p>2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA) Patient Preoperative Skin Preparation</p> <p>Non-sterile Solution Applicator is sterile if package is intact</p> <p>10.5 ml APPLICATOR </p> <p>Professional Use Only Do Not Reuse Not made with natural rubber latex</p>						
<p>Drug Facts</p> <table border="0" style="width: 100%; border-collapse: collapse;"> <tr> <td style="border-bottom: 1px solid black; padding: 2px;">Active ingredients</td> <td style="border-bottom: 1px solid black; padding: 2px; text-align: right;">Purposes</td> </tr> <tr> <td style="padding: 2px;">Chlorhexidine gluconate 2% w/v.....</td> <td style="padding: 2px; text-align: right;">Antiseptic</td> </tr> <tr> <td style="padding: 2px;">Isopropyl alcohol 70% v/v.....</td> <td style="padding: 2px; text-align: right;">Antiseptic</td> </tr> </table> <p>Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.</p> <p>Warnings</p> <p>For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY</p> <ul style="list-style-type: none"> ■ solution contains alcohol and gives off flammable vapors ■ avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable. ■ do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) ■ do not allow solution to pool ■ remove wet materials from prep area <p>Allergy Alert</p> <p>This product may cause a severe allergic reaction. Symptoms may include:</p> <ul style="list-style-type: none"> ■ wheezing/difficulty breathing ■ shock ■ facial swelling ■ hives ■ rash <p>If an allergic reaction occurs, stop use and seek medical help right away. ▶</p>		Active ingredients	Purposes	Chlorhexidine gluconate 2% w/v.....	Antiseptic	Isopropyl alcohol 70% v/v.....	Antiseptic
Active ingredients	Purposes						
Chlorhexidine gluconate 2% w/v.....	Antiseptic						
Isopropyl alcohol 70% v/v.....	Antiseptic						

<p>ChloroPrep® One-Step / ChloroPrep® With Tint</p> <p>2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)</p>
<p>Drug Facts (continued)</p> <p>Do not use</p> <ul style="list-style-type: none"> ■ on patients allergic to chlorhexidine gluconate or any other ingredient in this product ■ for lumbar puncture or in contact with the meninges ■ on open skin wounds or as a general skin cleanser <p>When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.</p> <p>Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p> <p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> ■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. ■ use in a well ventilated area ■ maximal treatment area for one applicator is approximately 8.4 in. x 8.4 in. (457 cm²) ■ remove applicator from package; do not touch sponge ■ hold the applicator with the sponge down. Pinch wings only once to activate the ampule and release the antiseptic. ■ wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin ■ completely wet the treatment area with antiseptic ■ dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds ■ moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes ■ do not allow solution to pool; tuck prep towels to absorb solution, and then remove ■ allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away. ■ discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available. <p>Other information ■ store between 15–30 °C (59–86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F)</p> <p>■ the tint will slowly fade from the skin. Soap and water, or alcohol may be used to remove the tint if desired.</p> <p>Inactive ingredients FD&C green #3 dye (Scrub-Teal applicators only), FD&C yellow #6 dye (Hi-Lite Orange applicators only), USP purified water</p> <p>Questions? ■ www.chloroprep.com ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)</p>
<p>CareFusion, El Paso, TX 79912 · Artwork No. 6-640248 · MXXXXXX · Made in the USA of US and imported parts</p>

Back




Lot/Exp Placeholder - for reference only




(b) (4)

(b) (4)

Lot/Exp Placeholder - for reference only.






Lot/Exp Placeholder - for reference only.

(b) (4)


Part No. 6-773700
10.5-ml ChloraPrep One-Step
Line Mold Labels



(b) (4)

(b) (4)


Part No. 6-773715
10.5-ml ChloroPrep With Tint
In-Line Mold Labels



(b) (4)

(b) (4)

Part No. 6-773725
10.5-ml ChloroPrep With Tint
In-Line Mold Labels




(b) (4)

(b) (4)

Artwork No. # 6-824248, Intermediate Carton 10.5-mL Applicator ChloroPrep 2% One-Step

1¹⁵/₃₂

ChloroPrep® One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact



WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts	Purposes
<p>Active ingredients Chlorhexidine gluconate 2% w/v Isopropyl alcohol 70% v/v</p> <p>Use Use on the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.</p> <p>Warnings For external use only. Flammable, keep away from fire or flame. To reduce the risk of fire, PREP CAREFULLY: <ul style="list-style-type: none"> solution contains alcohol and gives off flammable vapors avoid getting solution into hair areas. Hair may take up to 1 hour to dry. Wet hair is flammable. do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) do not allow solution to pool remove wet materials from prep area </p> <p>Other Warnings This product may cause a severe allergic reaction. Symptoms may include: <ul style="list-style-type: none"> itching/difficulty breathing facial swelling hives itch </p> <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Do not use <ul style="list-style-type: none"> if you are allergic to chlorhexidine gluconate or any other ingredients on this product for longer duration or in contact with the mucous membranes on open skin wounds or as a general skin cleanser </p> <p>When using this product, keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.</p> <p>Stop use and seek medical help if irritation, sensitization, or allergic reaction occurs. There may be signs of a serious condition. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p> <p>Directions <ul style="list-style-type: none"> use with care in geriatric patients or infants under 2 months of age. These products may cause irritation or chemical burns. use in a well-ventilated area prepare treatment area for one applicator is approximately 8.4 in. x 4.4 in. (457 cm) remove applicator from package; do not touch sponge hold the applicator with the sponge down. Press away against the ampule to activate the ampule and release the antiseptic. wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin completely wet the treatment area with antiseptic any surgical sites (i.e., abdominal or other) are gently prepared back-and-forth strokes for 30 seconds moist surgical sites (e.g., inguinal fold) use gentle repeated back-and-forth strokes for 2 minutes do not allow solution to pool; keep applicator in alcohol solution and then remove allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away. discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available. </p> <p>Other information <ul style="list-style-type: none"> store between 15-30 °C (59-86 °F) avoid freezing and exposure heat above 40 °C (104 °F) </p> <p>Inactive ingredient USP purified water</p> <p>Questions? <ul style="list-style-type: none"> www.chloroprep.com call 1-800-523-0502 (M-F 8 a.m.-4 p.m. CST) </p>	<p>2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA) Patient Preoperative Skin Preparation Non-sterile Solution Applicator is sterile if package is intact</p>

25 applicators
10.5 ml APPLICATORS
Clear


ChloroPrep® One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
10.5 ml APPLICATORS
Clear

ChloroPrep® One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact
10.5 ml APPLICATORS
Clear




10.5 ml APPLICATORS
Clear
Not made with natural rubber latex
STERILE EO
Professional Use Only
Store between 15-30 °C (59-86 °F)
Avoid freezing and excessive heat above 40 °C (104 °F)
25 applicators
0.36 fl. oz. (10.5 ml) each

ChloroPrep® One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact



ChloroPrep® One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact
10.5 ml APPLICATORS
Clear



LOT XXXXXX (PRINTED IN) EXP DATE XXXXX
No Variish
(01) 20354365400048
©2000 CareFusion. All rights reserved. ChloroPrep® is a registered trademark of CareFusion Corporation or one of its subsidiaries.

No Variish / No printing area

3¹/₄

6¹/₂

2⁵¹/₆⁴

1 7¹/₄

11¹/₁₆

1 2⁷/₈

5⁵/₈

1 2⁷/₈

5¹⁹/₃₂

3 7¹⁹/₃₂

1 15/32

3 1 1/16

1 1/16

1 2 7/8

5 5/8

1 2 7/8

5 1 9/32


3 7 9/32

2 5 1/64

6 1/2

1 7 1/4

3 1/4



ChloraPrep® With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Non-sterile Solution
Patient Preoperative Skin Preparation
Applicator is sterile if package is intact

WARNING: FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.	
Drug Facts	Purpose(s)
Active ingredients Chlorhexidine gluconate 2% w/v Isopropyl alcohol 70% v/v	Indication(s) Antiseptic
Use Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.	
Warnings For external use only. Flammable, keep away from fire or flame. To reduce the risk of fire, PREP CAREFULLY: <ul style="list-style-type: none"> solution contains alcohol and gives off flammable vapors avoid getting solution into hair areas. Hair may take up to 1 hour to dry. Wet hair is flammable. do not drink or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) do not allow solution to pool remove any residual prep from area Allergy Alert: This product may cause a severe allergic reaction. Symptoms may include: <ul style="list-style-type: none"> redness/itching/swelling hives swelling itching redness stinging difficulty breathing If an allergic reaction occurs, stop use and seek medical help right away.	
Do not use <ul style="list-style-type: none"> on patients allergic to chlorhexidine gluconate or any other ingredient in this product for longer periods or in contact with the mucosae on open skin wounds or as a general skin cleanser When using this product, keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.	
Stop use and ask a doctor if: irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
Directions <ul style="list-style-type: none"> use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. use in a well-ventilated area shaded treatment area for one applicator is approximately 8.4 in. x 4.4 in. (457 cm²) remove applicator from package; do not touch sponge hold the applicator with the sponge down. Press sponge gently against the treatment area until liquid is visible on the skin press the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin repeat until the hairless area is completely dry any residual prep (e.g., inguinal fold) use gentle repeated back-and-forth strokes for 30 seconds do not allow solution to pool; keep prep areas to avoid pooling, and then remove allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away. discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available. 	
Other information <ul style="list-style-type: none"> store between 15-30 °C (59-86 °F) avoid freezing and exposure to heat above 40 °C (104 °F) 	
Inactive ingredients FD-C120 80 mg, USP purified water	
Questions? <ul style="list-style-type: none"> www.chloraprep.com call 1-800-523-0502, 9 A.M.-5 P.M., 5 days a week (CST) 	

ChloraPrep® With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)

10.5 ml
APPLICATORS
Hi-Lite Orange®


ChloraPrep® With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact

10.5 ml
APPLICATORS
Hi-Lite Orange®

CareFusion
El Paso, TX 79912
1-800-523-0502
www.chloraprep.com
Made in the USA of US and imported parts

10.5 ml
APPLICATORS
Hi-Lite Orange®

Not made with natural rubber latex
STERILE/EO
Professional Use Only
Store between 15-30 °C (59-86 °F)
Avoid freezing and excessive heat above 40 °C (104 °F)
25 applicators
0.36 fl. oz. (10.5 ml) each



ChloraPrep® With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact

WARNING
FLAMMABLE
Keep away from fire or flame.
To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off flammable vapors
- avoid getting solution into hair areas
- hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drink or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

NDC 054365-400-12
Cat. No. 260715

ChloraPrep® With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact

10.5 ml
APPLICATORS
Hi-Lite Orange®

LOT XXXXXX (printed on line)
EXP DATE XXXX
No Varnish
BARCODE PLACEHOLDER

©2008 CareFusion. All rights reserved. ChloraPrep® is a registered trademark of CareFusion Corporation or one of its subsidiaries.


No Varnish / No printing area

Artwork No. 6-824715 NXXXXXX

Artwork No. 6-824725, 10.5-mL ChloroPrep with Tint Intermediate Carton

1¹⁵/₃₂

ChloroPrep® With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact



25 applicators
10.5 ml
APPLICATORS
Scrub Teal®

WARNING, FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.	
Drug Facts	Purposes
Active Ingredients Chlorhexidine gluconate 2% w/v Isopropyl alcohol 70% v/v	Antiseptic Preoperative Skin Preparation
Warnings For external use only. Flammable. Keep away from fire or flame. To reduce the risk of fire, PREP CAREFULLY! • solution contains alcohol and gives off flammable vapors • avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wash hair in hairless areas. • do not shake or use ignition source (e.g., candle, heater) until solution is completely dry (minimum of 2 minutes on hairless skin up to 1 hour in hair) • do not allow solution to pool • remove wet materials from prep area Caution: This product may cause a severe allergic reaction. Symptoms may include: • wheezing/difficultly breathing • hives • facial swelling • dizziness • rash • eye allergic reaction occurs, stop use and seek medical help right away. Do not use • do not use on patients allergic to chlorhexidine gluconate or any other ingredient in this product • for laceration or in contact with the mucous membranes • for open skin wounds or as a general skin cleanser When using the product: keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cool water right away and consult a doctor. Stop use and ask a doctor if: irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
Directions • use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. • use in a well-ventilated area. • maximal treatment area for one applicator is approximately 6.4 in. x 8.4 in. (167 cm²) • remove applicator from package, do not touch applicator. • wet the applicator with the solution. Then, with the scrub side to activate the applicator and release the antiseptic. • wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin completely wet the treatment area with solution. • dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds • dry surgical sites (e.g., inguinal folds): use gentle repeated back-and-forth strokes for 2 minutes • do not allow solution to pool; back prep towels to absorb solution, and then remove • when the solution is completely dry (minimum of 2 minutes on hairless skin, up to 1 hour in hair). Do not blot or wipe away. • discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.	
Other Information • store between 15-30 °C (59-86 °F) • avoid freezing and excessive heat above 40 °C (104 °F)	
Inactive Ingredients FDAC green #3 dye, USP purified water	
Questions? • visit www.chloroprep.com • call 1-800-832-0982 M-F 9 a.m.-6 p.m. CST	

ChloroPrep® With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact
10.5 ml
APPLICATORS
Scrub Teal®

ChloroPrep® With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact
10.5 ml
APPLICATORS
Scrub Teal®



10.5 ml
APPLICATORS
Scrub Teal®
Not made with natural rubber latex
STERILE EO
Professional Use Only
Store between 15-30 °C (59-86 °F)
Avoid freezing and excessive heat above 40 °C (104 °F)
25 applicators
0.36 fl. oz. (10.5 ml) each

NDC 054365-400-09
Cat. No. 260725
ChloroPrep® With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact

**WARNING
FLAMMABLE**
Keep away from fire or flame.
To reduce risk of fire, PREP CAREFULLY!
• solution contains alcohol and gives off flammable vapors
• avoid getting solution into hairy areas.
• hair may take up to 1 hour to dry. Wash hair in hairless areas.
• do not shake or use ignition source (e.g., candle, heater) until solution is completely dry (minimum of 2 minutes on hairless skin, up to 1 hour in hair)
• do not allow solution to pool
• remove wet materials from prep area

ChloroPrep® With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact
10.5 ml
APPLICATORS
Scrub Teal®



LOT XXXXXX
EXP DATE XXXX
No Version
BARCODE PLACEHOLDER

No Varnish/No printing area

3¹/₁₆

1¹/₁₆

1 2⁷/₈

5⁵/₈

1 2⁷/₈

5¹/₃₂

3 7¹/₃₂

2⁵/₆₄

6¹/₂

1 7¹/₄

3¹/₄

Artwork No. 6-824725 LXXXXXX

5" width

Front

(b) (4)

WARNING FLAMMABLE	Cat. No. 260800 / 260800NS / 260800NSB NDC 054365-400-14 Cat. No. 260815 / 260815NS / 260815NSB NDC 054365-400-13 Cat. No. 260825 / 260825NS / 260825NSB NDC 054365-400-05						
<p>Keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:</p> <ul style="list-style-type: none"> • do not use 26-ml applicator for head and neck surgery or on an area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead. • solution contains alcohol and gives off flammable vapors • avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable. • do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) • do not allow solution to pool • remove wet materials from prep area 	<p>ChloroPrep® One-Step ChloroPrep® With Tint</p> <p>2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)</p> <p>Patient Preoperative Skin Preparation</p> <p>Non-sterile Solution</p> <p>Applicator is sterile if package is intact</p> <p>26 ml APPLICATOR</p> <p style="font-size: 8px;">Professional Use Only Do Not Reuse Not made with natural rubber latex</p>						
<p>Drug Facts</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="font-size: 8px;">Active ingredients</td> <td style="font-size: 8px;">Purposes</td> </tr> <tr> <td style="font-size: 8px;">Chlorhexidine gluconate 2% w/v.....</td> <td style="font-size: 8px;">Antiseptic</td> </tr> <tr> <td style="font-size: 8px;">Isopropyl alcohol 70% v/v.....</td> <td style="font-size: 8px;">Antiseptic</td> </tr> </table> <p>Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.</p> <p>Warnings</p> <p>For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:</p> <ul style="list-style-type: none"> ■ do not use 26-ml applicator for head and neck surgery ■ do not use on an area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead ■ solution contains alcohol and gives off flammable vapors ■ avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable. ■ do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) ■ do not allow solution to pool ■ remove wet materials from prep area 		Active ingredients	Purposes	Chlorhexidine gluconate 2% w/v.....	Antiseptic	Isopropyl alcohol 70% v/v.....	Antiseptic
Active ingredients	Purposes						
Chlorhexidine gluconate 2% w/v.....	Antiseptic						
Isopropyl alcohol 70% v/v.....	Antiseptic						

4" length


<p>ChloroPrep® One-Step / ChloroPrep® With Tint</p> <p>2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)</p>
<p>Drug Facts (continued)</p> <p>Allergy Alert This product may cause a severe allergic reaction. Symptoms may include: ■ wheezing/difficulty breathing ■ shock ■ facial swelling ■ hives ■ rash If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Do not use</p> <ul style="list-style-type: none"> ■ on patients allergic to chlorhexidine gluconate or any other ingredient in this product ■ for lumbar puncture or in contact with the meninges ■ on open skin wounds or as a general skin cleanser <p>When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.</p> <p>Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p> <p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> ■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. ■ use in a well ventilated area ■ maximal treatment area for one applicator is approximately 13.2 in. x 13.2 in. (1126 cm²) ■ remove applicator from package; do not touch sponge ■ hold the applicator with the sponge down. Pinch wings only once to activate the ampules and release the antiseptic. ■ wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin ■ completely wet the treatment area with antiseptic ■ dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds ■ moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes ■ do not allow solution to pool; tuck prep towels to absorb solution, and then remove ■ allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away. ■ discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available. <p>Other information ■ store between 15–30 °C (59–86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F) ■ the tint will slowly fade from the skin. Soap and water, or alcohol may be used to remove the tint if desired.</p> <p>Inactive ingredients FD&C green #3 dye (Scrub-Teal applicators only), FD&C yellow #6 dye (Hi-Lite Orange applicators only), USP purified water</p> <p>Questions? ■ www.chloroprep.com ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)</p>
CareFusion, El Paso, TX 79912 · Artwork No. 6-775815 · LXXXXXX · Made in the USA of US and imported parts

Back

(b) (4)

Lot/Exp Placeholder -
for reference only

Lot # and Exp. Date are displayed
at this end of the package.





(b) (4)

Lot/Exp Placeholder - for reference only

Lot # and Exp. Date are displayed at this end of the package.

Barcode Placeholder

NDC 054365-400-13
Cat. No. 260815

Single Use
ChloroPrep® With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact
26 ml APPLICATOR
Hi-Lite Orange®

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings
For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery
- do not use on an area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead.
- solution contains alcohol and gives off flammable vapors
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Allergy Alert: This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- on patients allergic to chlorhexidine gluconate or any other ingredient in this product
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 13.2 in. x 13.2 in. (1126 cm²). Do not use 26-ml applicator for area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead.
- do not use 26-ml applicator for head and neck surgery
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wing **only once** to activate the ampules and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- do not allow solution to pool; tuck prep towels to absorb solution, and then remove
- **allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15–30 °C (59–86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients
FD&C yellow #6 dye, USP purified water

Questions?

- www.chloroprep.com
- call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

CareFusion, El Paso, TX 79912 • Artwork No. 6-786815 • OXXXXXX
Made in the USA of US and imported parts

8.25

243


305

60

80.5

(b) (4)

Lot # and Exp. Date are displayed at this end of the package.




Lot/Exp Placeholder - for reference only

(b) (4)

Part No. 6-773800-L
26-ml ChloraPrep One-Step
In-Line Mold Labels

(b) (4)




(b) (4)

(b) (4)


Part No. 6-773800-R
26-ml ChloroPrep One-Step
In-Line Mold Labels

(b) (4)




(b) (4)

(b) (4)



(b) (4)

Part No. 6-773815-R
26-ml ChloraPrep with tint
In-Line Mold Labels



(b) (4)


(b) (4)

(b) (4)

(b) (4)

Part No. 6-773825-L
26-ml ChloroPrep with tint
In-Line Mold Labels

(b) (4)




(b) (4)

(b) (4)

Part No. 6-773825-R
26-ml ChloroPrep with tint
In-Line Mold Labels

(b) (4)



(b) (4)

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

VALERIE S PRATT
09/12/2017