


14-Jan-2017 02:22

PRODUCTION

(NOT COLOR ACCURATE)

(b) (4

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces




PRODUCTION


(NOT COLOR ACCURATE)

(b)	(4)
· · · /	· /

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.5 pt
• BULLETS	N/A
SPACE BEFORE BULLET	N/A
WARNING BOX LINE	.5 pt

Reference ID: 4075656








(NOT COLOR ACCURATE)

(b) (4)

090177e18b3676e3\Final\Final On: 07-Feb-2017 07:08 (GMT)

Reference ID: 4075656



3.175mm x 3.175mm DATA MATRIX CODE (ECC200), PRINTED IN BLACK OR DARK BLUE ON UNPRINTED WHITE BACKGROUND **DATA MATRIX = 083985**

PANEL 1

PANEL 2

	of use
	he child has ever had an allergic reaction to any other pain reliever/fever reducer ht before or after heart surgery
	doctor hefore use if
 chi chi chi or chi chi 	mach bleeding warning applies to the child Id has problems or serious side effects from taking pain relievers or fever reducers Id has a history of stomach problems, such as heartburn Id has high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, had a stroke Id has not been drinking fluids Id has lost a lot of fluid due to vomiting or diarrhea Id is taking a diuretic
∎ un ■ tak	doctor or pharmacist before use if the child is der a doctor's care for any serious condition ing any other drug
Whe	using this product at the with food or milk if stomach upset occurs
chi chi chi chi fev rec	use and ask a doctor if Id experiences any of the following signs of stomach bleeding: feels faint — worits blood — has bloody or black stools has stomach pain that does not get better Id has symptoms of heart problems or stroke: chest pain — trouble breathing — weakness in one part or side of body surred speech — Ieg swelling Id does not get any relief within first day (24 hours) of treatment er or pain gets worse or lasts more than 3 days ness or swelling is present in the painful area / new symptoms appear
	out of reach of children. In case of overdose, get medical help or contact a on Control Center right away.

do not find rig repeat o	duct does not contain (give more than directed	i sible, use weight to o needed	ete warnings for adult i dose; otherwise use age	
		Dosing Chart		
	Weight (lb)	Age (yr)	Dose (tablets)]
	under 24 lb	under 2 yr	ask a doctor]
	24-35 lb	2-3 yr	2 tablets]
	36-47 lb	4-5 yr	3 tablets]
	48-59 lb	6-8 yr	4 tablets]
	60-71 lb	9-10 yr	5 tablets	1
	72-95 lb	11 yr	6 tablets	1
Phenyl one dos	nformation ketonurics: contains ph se lasts 6-8 hours a dry place at 20-25°C		per tablet	-

BASE PANEL

	glycolate, sorbitol
	Cuestions or comments? call toll-free 1-800-88-ADVIL or ask your pharmacist, doctor or health care professional
l	PAA083985.FDA01

(b) (4)

PRODUCTION (NOT COLOR ACCURATE)

DRUG FACTS TEXT DEFINED TYPE SIZE • DRUG FACTS TITLE 9 pt • DRUG FACTS CONTINUED 8 pt 8 pt HEADINGS • SUBHEADINGS/BODY TEXT 6 pt • LEADING 6.5 pt • # OF CHARACTERS PER INCH <39 • BULLETS 5 pt • SPACE BEFORE BULLET 2 ems • BARLINES, HAIRLINES 1.5 pt, .5 pt SPACE BETWEEN HAIRLINES AND BOX END 2 spaces