

Drug Facts (continued)

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- redness or swelling is present in the painful area
- any new symptoms appear
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing or the caplet feels stuck in your throat
- you get nervous, dizzy, or sleepless
- nasal congestion lasts more than 7 days

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- swallow whole; do not crush or chew
- drink a full glass of water with each dose
- adults and children 12 years and older: 1 caplet every 12 hours; do not take more than 2 caplets in 24 hours
- children under 12 years: do not use

Other information

- each caplet contains: sodium 22 mg
- store at 20-25°C (68-77°F)
- store in a dry place

Inactive ingredients colloidal silicon dioxide, hypromellose, lactose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments? 1-800-986-0369 (Mon-Fri 9AM - 5PM EST)

Bayer

Bayer HealthCare LLC
Whippany, NJ 07981
Bayer, the Bayer Cross, Aleve-D and All Day Strong are registered trademarks of Bayer.

XXXXXX

3 258666 56952 3

Do not use if the blister seal is cut or broken.

10 CAPLETS
CAPSULE-SHAPED TABLETS

1 CAPLET - 12 HOURS

12 HOUR MULTI-SYMPTOM RELIEF OF
SINUS PRESSURE
HEADACHE • SINUS CONGESTION
NASAL CONGESTION • BODY ACHES

aproxen sodium 220 mg (NSAID) / Pseudoephedrine HCl 120 mg extended-release tablet
Pain reliever/fever reducer/nasal decongestant

ALEVE-D
ALL DAY STRONG

SINUS & COLD

NON-DROWSY

ALL DAY STRONG
ALEVE-D
SINUS & COLD

Drug Facts

Active ingredients (in each caplet)
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID) Pain reliever/fever reducer
Pseudoephedrine HCl 120 mg, extended-release Nasal decongestant
*nonsteroidal anti-inflammatory drug

Uses temporarily relieves these cold, sinus, and flu symptoms:

- sinus pressure
- minor body aches and pains
- headache
- nasal and sinus congestion (promotes sinus drainage and restores freer breathing through the nose)
- fever

Warnings
Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- stomach
- skin redness
- rash
- blisters

 If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (an anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- in children under 12 years of age

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, have trouble urinating due to an enlarged prostate gland, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Drug Facts (continued)

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- redness or swelling is present in the painful area
- any new symptoms appear
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing or the caplet feels stuck in your throat
- you get nervous, dizzy, or sleepless
- nasal congestion lasts more than 7 days

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- swallow whole: do not crush or chew
- drink a full glass of water with each dose
- adults and children 12 years and older: 1 caplet every 12 hours; do not take more than 2 caplets in 24 hours
- children under 12 years: do not use

Other information

- each caplet contains: sodium 22 mg
- store at 20-25°C (68-77°F)
- store in a dry place

Inactive ingredients colloidal silicon dioxide, hypromellose, lactose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments? 1-800-986-0369 (Mon-Fri 9AM - 5PM EST)

Bayer

Do not use if the blister seal is cut or broken.

Dist. by: Bayer HealthCare LLC
Whippany, NJ 07981
Bayer, the Bayer Cross, Alvee-D and All Day Strong are registered trademarks of Bayer.

XXXXXX



20 CAPLETS
CAPSULE-SHAPED TABLETS

1 CAPLET - 12 HOURS

ALL DAY STRONG
ALEVE-D
SINUS & COLD

NON-DROWSY

12 HOUR MULTI-SYMPTOM RELIEF OF
SINUS PRESSURE
HEADACHE • SINUS CONGESTION
NASAL CONGESTION • BODY ACHES

naproxen sodium 220 mg (5.1%)
Pseudoephedrine HCl 120 mg extended-release tablet
Pain reliever-fever reducer/nasal decongestant

BAYER

ALL DAY STRONG
ALEVE-D
SINUS & COLD

Drug Facts

Active ingredients (in each caplet)
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)* Pain reliever/fever reducer
Pseudoephedrine HCl 120 mg, extended-release Nasal decongestant
*nonsteroidal anti-inflammatory drug

Uses temporarily relieves these cold, sinus, and flu symptoms:
• sinus pressure • minor body aches and pains
• headache • nasal and sinus congestion (promotes sinus drainage and restores freer breathing through the nose)
• fever

Warnings
Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Purposes

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- in children under 12 years of age

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heart burn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, have trouble urinating due to an enlarged prostate gland, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

LIFT HERE

LIFT HERE

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

VALERIE S PRATT
09/12/2017