


3 0045-0212-122



Questions or comments?
call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Inactive ingredients
acesulfame potassium, croscarmellose sodium, dibasic calcium phosphate, flavor, microcrystalline cellulose, stearic acid

Do not use if carton is open or if blister unit is open or torn



TEAR OFF A SECTION



FOLD CORNER INWARD
AND PEEL BACK FOIL

Drug Facts (continued)

Directions
In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Purposes
Anti-diarrheal
Anti-gas

Active ingredients (in each caplet)
Loperamide HCl 2 mg
Simethicone 125 mg

Uses
relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas

Warnings
Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl
Heart alert: Taking more than directed can cause serious heart problems or death

Do not use
if you have bloody or black stool
if you have difficulty swallowing

Ask a doctor before use if you have
ever a history of liver disease
mucus in the stool

Ask a doctor or pharmacist before use if you are taking antibiotics
When using this product, tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if
diarrhea lasts for more than 2 days
if you get abdominal swelling or bloating. These may be signs of a serious condition.
if you have difficulty swallowing the caplet


If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.

Other information
each caplet contains calcium 165 mg, sodium 5 mg
store between 20-25°C (68-77°F). Protect from light.

children 9-11 years	1 caplet after the first loose stool, but no more than 3 caplets in 24 hours
children 6-8 years	1 caplet after the first loose stool, but no more than 2 caplets in 24 hours
children 2-5 years	ask a doctor
children under 2 years	do not use

Care To Recycle™

Manufactured by
JOHNSON & JOHNSON CONSUMER INC.
Medicine Division
Fort Washington, PA 19084 USA
© 18/CI 2016
www.imodium.com
U.S. Patent No. 6,103,260



LOT
EXP

Imodium®

How can we help?
1-877-895-3665

See New Warning and Directions NDC 50580-338-12

Imodium®

Loperamide HCl, 2 mg/Simethicone, 125 mg Antidiarrheal/Anti-gas

Multi-Symptom Relief

Caplet


Relieves symptoms of diarrhea **plus**

- Cramps & Pressure
- Bloating
- Gas

12 caplets*



*capsule-shaped tablets Actual Size



00000000



Drug Facts (continued)
Inactive ingredients
 ace sulfamethoxazole, croscarmellose sodium, dibasic calcium phosphate, fava, microcrystalline cellulose, stearic acid

Questions or comments?
 Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)



TEAR OFF A SECTION



FOLD CORNER INWARD AND PEEL BACK FOIL

Drug Facts (continued)

Directions
 Drink plenty of clear fluids to help prevent dehydration caused by diarrhea.
 Take only on an empty stomach (1 hour before or 2 hours after a meal).
 Take with a full (8 oz.) glass of water.
 Find right dose on chart below. If possible, use weight to dose, otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool, but no more than 4 caplets in 24 hours
children 9-11 years	1 caplet after the first loose stool, but no more than 3 caplets in 24 hours
children 6-8 years	1/2 caplet after each subsequent loose stool, but no more than 3 caplets in 24 hours
children 2-5 years	1/4 caplet after each subsequent loose stool, but no more than 2 caplets in 24 hours
children under 2 years (up to 33 lbs)	do not use

ask a doctor

Other information
 Each caplet contains calcium 165 mg, sodium 3 mg

store between 20-25°C (68-77°F). Protect from light.
 do not use if carton is open or if blister unit is open or torn

Drug Facts

Active ingredients (in each caplet)
 Loperamide HCl 2 mg
 Simethicone 125 mg

Purposes
 Anti-diarrheal
 Anti-gas

Uses
 Relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas

Warnings
 Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Heart alert
 Taking more than directed can cause serious heart problems or death

Do not use
 if you have bloody or black stool
 if you have difficulty swallowing

Ask a doctor before use if you have
 ever a history of liver disease

Ask a doctor or pharmacist before use if you are taking antibiotics
 or a history of liver disease

When using this product
 drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if
 symptoms get worse
 diarrhea lasts for more than 2 days
 you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children.



Distributed by:
JOHNSON & JOHNSON
 JOHNSON OTC Inc.
 Medical Devices Division
 First Floor, 1900
 E. Washington Ave., Allentown, PA 19034 USA
 © J&J, 2016
 www.Imodium.com
 U.S. Patent No. 6,103,260



Imodium®
 How can we help?
 1-877-895-3665

00000000

See New Warning and Directions NDC 50580-338-18

Imodium®

Loperamide HCl, 2 mg/Simethicone, 125 mg Antidiarrheal/Anti-gas

Multi-Symptom Relief
 Caplet




Relieves symptoms of diarrhea **plus**

- Cramps & Pressure
- Bloating
- Gas


18 Caplets*



*capsule-shaped tablets Actual Size




00000000



Drug Facts (continued)
Inactive ingredients
 ace sulfanilamide potassium, croscarmellose sodium, dibasic calcium phosphate, fava, microcrystalline cellulose, stearic acid

Questions or comments?
 Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)



Drug Facts (continued)

Directions
 ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea
 ■ take only on an empty stomach (1 hour before or 2 hours after a meal)
 ■ take in a full (8 oz.) glass of water
 ■ and right dose on chart below; if possible, use weight to dose; otherwise, use age

adults and children 12 years and over	2 caplets after the first loose stool, but no more than 4 caplets in 24 hours
children 9-11 years	1 caplet after the first loose stool, but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool, but no more than 2 caplets in 24 hours
children 2-5 years (24 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information
 ■ each caplet contains calcium 165 mg, sodium 3 mg
 ■ store between 20-25°C (68-77°F). Protect from light.
 ■ do not use if carton is open or if blister unit is open or torn

Drug Facts

Active ingredients (in each caplet)
 Loperamide HCl 2 mg; Simethicone 125 mg
 Anti-diarrheal
 Anti-gas

Purposes
 Uses: relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas

Warnings
 Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl
 Heart alert: Taking more than directed can cause serious heart problems or death

Do not use
 ■ if you have bloody or black stool
 ■ if you have difficulty swallowing

Ask a doctor before use if you have
 ■ ever ■ mucus in the stool ■ a history of liver disease

Ask a doctor or pharmacist before use if you are taking antibiotics
 When using this product, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if
 ■ symptoms get worse ■ diarrhea lasts for more than 2 days
 ■ you get abdominal swelling or bulging. These may be signs of a serious condition.

■ you have difficulty swallowing the caplet

If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children.



Distributed by:
JOHNSON & JOHNSON
 JANSSEN-ORIGEN
 Pharmaceutical Research Division
 P.O. Box 2009
 Kenilworth, NJ 07033 USA
 © J&J, 2018
 www.imodium.com
 U.S. Patent No. 6,103,260



Imodium®
 How can we help?
 1-877-895-3665

00000000

See New Warning and Directions NDC 50580-338-18

Imodium®

Loperamide HCl, 2 mg/Simethicone, 125 mg Antidiarrheal/Anti-gas

Multi-Symptom Relief
 Caplet




Relieves symptoms of diarrhea **plus**

- Cramps & Pressure
- Bloating
- Gas

18 Caplets*



*capsule-shaped tablets Actual Size



00000000

See New Warning and Directions

NDC 50580-338-30

Imodium®

Loperamide HCl, 2 mg/Simethicone, 125 mg Antidiarrheal/Anti-gas

Multi-Symptom Relief Caplet

Relieves symptoms
of diarrhea **plus**

- Cramps & Pressure
- Bloating
- Gas

30 Caplets*

*capsule-shaped tablets





Actual Size

Multi-Symptom Relief 30 Caplets

Loperamide HCl, 2 mg/Simethicone, 125 mg Antidiarrheal/Anti-gas

Imodium®



3 0045-02 12-30 6

LOT
EXP

Imodium®
How can we help?
1-877-895-3665

www.imodium.com U.S. Patent No. 6,103,260
©J&JCI 2016
Fort Washington, PA 19034 USA
McNeil Consumer Healthcare Division
JOHNSON & JOHNSON CONSUMER INC.
Distributed by

Care To Recycle™
215-273-8755 (collect)
call 1-877-895-3665 (toll-free) or
recycle.steak.com

Questions or comments?
do not use
children under
(up to 55 lbs)
2 years

Other information
■ each caplet contains: calcium 165 mg,
sodium 3 mg
■ store between 20-25°C (68-77°F).
Protect from light.
■ do not use if carton is open or if printed
foil seal under bottle cap is open or torn

Drug Facts (continued)
ask a doctor
children 2-5
years
(34 to 47 lbs)
children under
do not use
(up to 55 lbs)
2 years

Drug Facts (continued)
ask a doctor
children 2-5
years
(34 to 47 lbs)
children under
do not use
(up to 55 lbs)
2 years

Drug Facts (continued)
ask a doctor
children 2-5
years
(34 to 47 lbs)
children under
do not use
(up to 55 lbs)
2 years

Drug Facts (continued)
ask a doctor
children 2-5
years
(34 to 47 lbs)
children under
do not use
(up to 55 lbs)
2 years

Drug Facts (continued)
ask a doctor
children 2-5
years
(34 to 47 lbs)
children under
do not use
(up to 55 lbs)
2 years

Drug Facts (continued)
ask a doctor
children 2-5
years
(34 to 47 lbs)
children under
do not use
(up to 55 lbs)
2 years

Drug Facts (continued)
ask a doctor
children 2-5
years
(34 to 47 lbs)
children under
do not use
(up to 55 lbs)
2 years

Drug Facts (continued)
ask a doctor
children 2-5
years
(34 to 47 lbs)
children under
do not use
(up to 55 lbs)
2 years

Drug Facts (continued)
ask a doctor
children 2-5
years
(34 to 47 lbs)
children under
do not use
(up to 55 lbs)
2 years

Drug Facts (continued)
ask a doctor
children 2-5
years
(34 to 47 lbs)
children under
do not use
(up to 55 lbs)
2 years

Drug Facts (continued)
ask a doctor
children 2-5
years
(34 to 47 lbs)
children under
do not use
(up to 55 lbs)
2 years

Drug Facts (continued)
ask a doctor
children 2-5
years
(34 to 47 lbs)
children under
do not use
(up to 55 lbs)
2 years

Drug Facts (continued)
ask a doctor
children 2-5
years
(34 to 47 lbs)
children under
do not use
(up to 55 lbs)
2 years

Drug Facts (continued)
ask a doctor
children 2-5
years
(34 to 47 lbs)
children under
do not use
(up to 55 lbs)
2 years

Drug Facts (continued)
ask a doctor
children 2-5
years
(34 to 47 lbs)
children under
do not use
(up to 55 lbs)
2 years

<p>Active ingredients (in each caplet) Purposes Loperamide HCl 2 mg Anti-diarrheal Simethicone 125 mg Anti-gas Uses relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas Warnings: Allergy alert: Do not use if you have</p>	<p>NDC 50580 338 30 Imodium Loperamide HCl, 2 mg/Simethicone, 125 mg Antidiarrheal/Anti-gas</p>	<p>Directions or pharmacist help: Use only if you are taking antibiotics. When using this product, drowsiness or dizziness may occur. Be careful when driving or operating machinery. Stop use and ask a doctor if symptoms get worse, diarrhea lasts for more than 2 days, you get abdominal swelling or bulging. These may be signs of a serious condition, or you have difficulty swallowing the caplet. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Directions: Drink plenty of clear fluids to help prevent dehydration caused by diarrhea. Take only on an empty stomach (1 hour before or 2 hours after a meal). Take with a full (8 oz.) glass of water. If possible, use weight to dose. Otherwise, use age. Adults and children 12 years and over: 2 caplets after the first loose stool; 1 caplet after each subsequent loose stool, but no more than 4 caplets in 24 hours. Children 9-11 years (60-95 lbs): 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool, but no more than 3 caplets in 24 hours. Children 6-8 years (40-59 lbs): 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool, but no more than 2 caplets in 24 hours. Children 2-5 years (14 to 47 lbs): ask a doctor. Children under 2 years (up to 33 lbs): do not use. Other information: Each caplet contains calcium 165 mg, sodium 3 mg. Store between 20-25°C (68-77°F). Protect from light. Do not use if printed foil seal on bottle is broken or missing.</p>
<p>LOT: EXP: 00000000</p>	<p>Multi-Symptom Relief 30 Caplets* *capsule-shaped tablets</p> <p>Heart alert: Taking more than directed can cause dizziness, fainting, or slow heart rate. Do not use if you have a heart condition or are taking heart medicine. Do not use if you have a history of liver disease. Do not use if you have bloody or black stool, if you have difficulty swallowing. Ask a doctor before use if you have a history of liver disease.</p>	<p>Questions or comments? call 1-877-895-3665 (toll-free) or 215-273-8755 (collect) Distributed by JOHNSON & JOHNSON CONSUMER INC. McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA © J&JCI 2016 www.imodium.com U.S. Patent No. 6,103,260</p>

See New Warning and Directions NDC 50580-338-30

Imodium®

Loperamide HCl, 2 mg/Simethicone, 125 mg Antidiarrheal/Anti-gas

Multi-Symptom Relief Caplet

Relieves symptoms
of diarrhea **plus**

- Cramps & Pressure
- Bloating
- Gas

30 Caplets*




*capsule-shaped tablets

Actual Size



000000



000000

Multi-Symptom Relief 30 Caplets

Loperamide HCl, 2 mg/Simethicone, 125 mg Antidiarrheal/Anti-gas

Imodium®

Imodium
How can we help?
1-877-895-3665

U.S. Patent No. 6,103,260
www.imodium.com
© J&J 2016
Fort Washington, PA 19034 USA
Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
Care To Recycle™

Drug Facts (continued)
Inactive ingredients
ascorbic acid, croscarmellose sodium, dibasic calcium phosphate, flavor, microcrystalline cellulose, stearic acid
Questions or comments?
Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Drug Facts (continued)
ask a doctor
children 2-5 years
(34 to 47 lbs)
children under 2 years
(up to 33 lbs)
do not use
Other information
each caplet contains: calcium 165 mg, sodium 3 mg
Store between 20°-25°C (68°-77°F).
Protect from light.
Do not use if carton is open or if printed foil seal under bottle cap is open or torn.



3 0045-02 12-30 6

Drug Facts (continued)
Directions
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)
Warnings
Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.
Heart alert: Taking more than directed can cause serious heart problems or death.
Do not use if you have blood or black stool or if you have difficulty swallowing.
Ask a doctor before use if you have ever taken antibiotics, are taking antiacids, or dizziness may occur. Be careful when driving or operating machinery.
Stop use and ask a doctor if symptoms get worse, diarrhea lasts for more than 2 days, you get abdominal swelling or bloating, or you have difficulty swallowing the caplet.
When using this product (drowsiness, dizziness, or drowsiness may occur). Be careful when driving or operating machinery.
Ask a doctor or pharmacist before use if you are taking antibiotics, are taking antiacids, or dizziness may occur. Be careful when driving or operating machinery.
Do not use if you have blood or black stool or if you have difficulty swallowing.
Ask a doctor before use if you have ever taken antibiotics, are taking antiacids, or dizziness may occur. Be careful when driving or operating machinery.
Stop use and ask a doctor if symptoms get worse, diarrhea lasts for more than 2 days, you get abdominal swelling or bloating, or you have difficulty swallowing the caplet.
When using this product (drowsiness, dizziness, or drowsiness may occur). Be careful when driving or operating machinery.
Ask a doctor or pharmacist before use if you are taking antibiotics, are taking antiacids, or dizziness may occur. Be careful when driving or operating machinery.

Drug Facts (continued)
Directions
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)
Warnings
Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.
Heart alert: Taking more than directed can cause serious heart problems or death.
Do not use if you have blood or black stool or if you have difficulty swallowing.
Ask a doctor before use if you have ever taken antibiotics, are taking antiacids, or dizziness may occur. Be careful when driving or operating machinery.
Stop use and ask a doctor if symptoms get worse, diarrhea lasts for more than 2 days, you get abdominal swelling or bloating, or you have difficulty swallowing the caplet.
When using this product (drowsiness, dizziness, or drowsiness may occur). Be careful when driving or operating machinery.
Ask a doctor or pharmacist before use if you are taking antibiotics, are taking antiacids, or dizziness may occur. Be careful when driving or operating machinery.



<p>Active ingredients (in each caplet)</p> <p>Loperamide HCl 2 mg Anti-diarrheal Simethicone 125 mg Anti-gas</p> <p>Uses - relieves symptoms of diarrhea plus bloating, pressure and gas, commonly referred to as gas.</p> <p>Warnings: Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl. Heart alert: Taking more than directed can cause serious heart problems or death. Do not use - if you have bloody or black stool - if you have difficulty swallowing. Ask a doctor before use if you have a fever - flu up in the stool - a history of liver disease. Ask a doctor or pharmacist before use if you are taking antibiotics. When using this product, dizziness, lightheadedness or dizziness may occur. Be careful when driving or operating machinery. Stop use and ask a doctor if symptoms get worse - diarrhea lasts for more than 2 days - you experience abdominal swelling or bloating. These may be signs of a serious condition - if you have difficulty swallowing the caplet. If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-2222)</p>	<p>Purposes</p> <p>Anti-diarrheal Anti-gas</p>	<p>NDC 50580-338-30</p> <p>Imodium® Loperamide HCl, 2mg/Simethicone, 125mg Antidiarrheal/Anti-gas</p> <p>Multi-Symptom Relief</p> <p>30 Caplets* *capsule-shaped tablets</p> <p>Directions - drink plenty of clear fluids to help prevent dehydration caused by diarrhea - take only on an empty stomach (1 hour before or 2 hours after a meal) - take with a full (8 oz.) glass of water - find right dose on chart. If possible, use weight-based dose; otherwise, use age.</p>	<table border="1"> <tr> <td>adults and children 2 years and over</td> <td>2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours</td> </tr> <tr> <td>children 9-11 years (60-95 lbs)</td> <td>1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours</td> </tr> <tr> <td>children 6-8 years (48-59 lbs)</td> <td>1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours</td> </tr> <tr> <td>children 2-5 years (34 to 47 lbs)</td> <td>ask a doctor</td> </tr> <tr> <td>children under 2 years (up to 33 lbs)</td> <td>do not use</td> </tr> </table> <p>Other information - each caplet contains: calcium 165 mg, sodium 3 mg - store between 20-25°C (68-77°F). Protect from light. - do not use if printed foil seal on bottle is broken or missing</p> <p>Questions or comments? call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)</p>	adults and children 2 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours	children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours	children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours	children 2-5 years (34 to 47 lbs)	ask a doctor	children under 2 years (up to 33 lbs)	do not use	<p>Johnson & Johnson Consumer Inc. Molecular HealthCare Division Fort Washington, PA 19004 USA © J&J 2016 USA 03-2930</p> <p>EXP. LOT.</p>
adults and children 2 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours													
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours													
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours													
children 2-5 years (34 to 47 lbs)	ask a doctor													
children under 2 years (up to 33 lbs)	do not use													

See New Warning and Directions

NDC 50580-338-42

Imodium®

Loperamide HCl, 2 mg/Simethicone, 125 mg Antidiarrheal/Anti-gas

Multi-Symptom Relief Caplet

Relieves symptoms
of diarrhea **plus**

- Cramps & Pressure
- Bloating
- Gas

42 Caplets*

*capsule-shaped tablets



Actual Size

Multi-Symptom Relief 42 Caplets

Loperamide HCl, 2 mg/Simethicone, 125 mg Antidiarrheal/Anti-gas

Imodium®



3 0045-02 12-42 9

Imodium
How can we help?
1-877-895-3665

LOT:
EXP:

Care To Recycle™
Distributed by
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
©J&JCI 2016
www.imodium.com U.S. Patent No. 6,103,260

Drug Facts (continued)
Inactive ingredients
acesulfame potassium, croscarmellose sodium, dibasic calcium phosphate, flavor, microcrystalline cellulose, stearic acid

Questions or comments?
Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Drug Facts (continued)
children 2-5 ask a doctor
years (34 to 47 lbs) do not use
children under 2 years (up to 33 lbs)
Other information
■ each caplet contains: calcium 165 mg, sodium 3 mg
■ store between 20-25°C (68-77°F). Protect from light.
■ do not use if carton is open or if printed foil seal under bottle cap is open or torn.

Drug Facts (continued)
adults and children 12 years and over
2 caplets after the first loose stool; 1 caplet after each subsequent loose stool, but no more than 4 caplets in 24 hours

children 9-11 years
1/2 caplet after the first loose stool; 1 caplet after each subsequent loose stool, but no more than 3 caplets in 24 hours

children 6-8 years
1/2 caplet after each subsequent loose stool, but no more than 2 caplets in 24 hours

(48-59 lbs)

Drug Facts (continued)
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions
■ think plenty of clear fluids to help prevent dehydration caused by diarrhea
■ take only on an empty stomach (1 hour before or 2 hours after a meal)
■ take with a full (8 oz.) glass of water
■ find right dose on chart, if possible, use weight to dose; otherwise, use age

Drug Facts (continued)
Do not use ■ if you have bloody or black stool
■ if you have difficulty swallowing
■ ever ■ mucus in the stool ■ a history of liver disease

Ask a doctor or pharmacist before use if you
are taking antibiotics
When using this product (tiredness, drowsiness or dizziness may occur; Be careful when driving or operating machinery;
Stop use and ask a doctor if
■ symptoms get worse
■ symptoms last for more than 2 days
■ diarrhea lasts for more than 2 days
■ you get abdominal swelling or bloating
■ these may be signs of a serious condition.
■ you have difficulty swallowing the caplet

Drug Facts (continued)
Warnings
Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.
Heart alert: Taking more than directed can cause serious heart problems or death

Purposes
Antidiarrheal Loperamide HCl 2 mg, Ant-gas Simethicone 125 mg.

<p>Active ingredients (in each caplet)</p> <p>Loperamide HCl 2 mg, Simethicone 125 mg</p> <p>Purposes</p> <p>Anti-diarrheal, Anti-gas</p> <p>Uses</p> <p>relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas</p> <p>Warnings: Allergy alert: Do not use if you have</p>	<p>NDC 50580 338 42</p> <p>Imodium</p> <p>Loperamide HCl, 2 mg/Simethicone, 125 mg</p> <p>Antidiarrheal/Anti-gas</p> <p>Multi-Symptom Relief</p> <p>42 Caplets*</p> <p>*capsule-shaped tablets</p>	<p>Ask a doctor or pharmacist before use if you are taking any blood thinners. When using this product, drowsiness or dizziness may occur. Be careful when driving or operating machinery. Stop use and ask a doctor if symptoms get worse or diarrhea lasts for more than 2 days or you get abdominal swelling or bulging. These may be signs of a serious condition. If you have difficulty swallowing the caplet, if pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-322-2222) Directions: Drink plenty of clear fluids to help prevent dehydration caused by diarrhea. Take only on an empty stomach (1 hour before or 2 hours after a meal). Take with a full (8 oz.) glass of water. If possible, use weight to dose; otherwise, use age. Adults and children 12 years and older: 2 caplets after the first loose stool; 1 caplet after each subsequent loose stool, but no more than 4 caplets in 24 hours. Children 9-11 years (60-95 lbs): 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool, but no more than 3 caplets in 24 hours. Children 6-8 years (40-59 lbs): 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool, but no more than 2 caplets in 24 hours. Children 2-5 years (34 to 47 lbs): ask a doctor. Children under 2 years (up to 33 lbs) do not use. Other information: Each caplet contains calcium 165 mg, sodium 3 mg. Store between 20-25°C (68-77°F). Protect from light. Do not use if printed foil seal on bottle is broken or missing. Questions or comments? Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect). Distributed by JOHNSON & JOHNSON CONSUMER INC., McKeesport Consumer Healthcare Division, Fort Washington, PA 19034 USA ©JAJCI 2016 www.imodium.com U.S. Patent No. 6,103,260</p>
<p>LOT: 00000000</p> <p>EXP:</p>	<p>never had a rash or other allergic reaction to loperamide HCl. Heart alert: Taking more than directed can cause serious heart problems or death. Do not use if you have bloody or black stool or if you have difficulty swallowing. Ask a doctor before use if you have a fever, muscle aches in the stool, or a history of liver disease.</p>	

See New Warning and Directions NDC 50580-338-42

Imodium®

Loperamide HCl, 2 mg/Simethicone, 125 mg Antidiarrheal/Anti-gas

Multi-Symptom Relief Caplet

Relieves symptoms
of diarrhea **plus**

- Cramps & Pressure
- Bloating
- Gas


42 Caplets*

*capsule-shaped tablets

Actual Size



000000



000000

Multi-Symptom Relief 30 Caplets

Loperamide HCl, 2 mg/Simethicone, 125 mg Antidiarrheal/Anti-gas

Imodium®

Imodium
How can we help?
1-877-895-3665

www.imodium.com U.S. Patent No. 6,103,260

Dispersed by:
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
JOHNSON & JOHNSON CONSUMER INC.

Care To Recycle™

Questions or comments?
Call 1-877-895-3665 (toll-free) or
215-273-8755 (collect)

Inactive ingredients
(continued)

acesulfame potassium, croscarmellose sodium,
dibasic calcium phosphate, flavor, microcrystalline
cellulose, stearic acid

Drug Facts (continued)

children 2-5 ask a doctor
years (34 to 47 bs)
children under do not use
(up to 33 bs)

Other information
each caplet contains: calcium 165 mg,
sodium 3 mg
Store between 20-25°C (68-77°F).
Protect from light.
Do not use if carton is open or if printed
foil seal under bottle cap is open or torn.



3 0045-02 12-42 9

Drug Facts (continued)

adults and children 12 years and over
1 caplet after each subsequent loose stool, but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)
1/2 caplet after each subsequent loose stool, but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)
1/2 caplet after the first loose stool, but no more than 2 caplets in 24 hours

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Drug Facts (continued)

Directions
dehydration caused by diarrhea
2 hours after a meal)
Take with a full (8 oz.) glass of water
Find right dose on chart. If possible, use weight to dose; otherwise, use age.

Drug Facts (continued)

Do not use if you have bloody or black stool or if you have difficulty swallowing.
Ask a doctor before use if you have ever had a history of liver disease or mucus in the stool.
Loperamide HCl 2 mg Antidiarrheal
Simethicone 125 mg Anti-gas
Uses relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas.
Warnings
Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.
Heart alert: Taking more than directed can cause serious heart problems or death.

Drug Facts (continued)

Active ingredients
Loperamide HCl 2 mg Antidiarrheal
Simethicone 125 mg Anti-gas
Purposes
relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas.
Warnings
Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.
Heart alert: Taking more than directed can cause serious heart problems or death.

<p>Active ingredients (in each caplet)</p> <p>Loperamide HCl 2 mg Anti-diarrheal Simethicone 125 mg Anti-gas</p> <p>Uses - relieves symptoms of diarrhea plus bloating, pressure and gas, commonly referred to as gas.</p> <p>Warnings: Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl. Heart alert: Taking more than directed can cause serious heart problems or death. Do not use - if you have bloody or black stool - if you have difficulty swallowing. Ask a doctor before use if you have: - fever - flu or stomach infection - a history of liver disease. Ask a doctor or pharmacist before use if you are taking anti-biotics. When using this product, drowsiness, dizziness or lightheadedness may occur. Be careful when driving or operating machinery. Stop use and ask a doctor if symptoms get worse - diarrhea lasts for more than 2 days - you experience abdominal swelling or bloating. These may be signs of a serious condition. - you have difficulty swallowing the caplet. If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-2222)</p>	<p>Purposes</p> <p>Anti-diarrheal Anti-gas</p>	<p>NDC 50580-338-42</p> <p>Imodium® Loperamide HCl, 2mg/Simethicone, 125 mg Antidiarrheal/Anti-gas</p> <p>Multi-Symptom Relief</p> <p>42 Caplets* *capsule-shaped tablets</p> <p>Directions - drink plenty of clear fluids to help prevent dehydration caused by diarrhea - take only on an empty stomach (1 hour before or 2 hours after a meal) - take with a full (8 oz.) glass of water - find right dose on chart. If possible, use weight-based dose; otherwise, use age.</p>	<table border="1"> <tr> <td>adults and children 2 years and over</td> <td>2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours</td> </tr> <tr> <td>children 9-11 years (60-95 lbs)</td> <td>1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours</td> </tr> <tr> <td>children 6-8 years (48-59 lbs)</td> <td>1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours</td> </tr> <tr> <td>children 2-5 years (34 to 47 lbs)</td> <td>ask a doctor</td> </tr> <tr> <td>children under 2 years (up to 33 lbs)</td> <td>do not use</td> </tr> </table> <p>Other information - each caplet contains: calcium 165 mg, sodium 3 mg - store between 20-25°C (68-77°F). Protect from light. - do not use if printed foil seal on bottle is broken or missing</p> <p>Questions or comments? call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)</p>	adults and children 2 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours	children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours	children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours	children 2-5 years (34 to 47 lbs)	ask a doctor	children under 2 years (up to 33 lbs)	do not use	<p>Johnson & Johnson Consumer Inc. Medical Center New Brunswick, NJ 08901 Fort Washington, PA 19004 USA © J&J 2016 EXP. DATE LOT</p>
adults and children 2 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours													
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours													
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours													
children 2-5 years (34 to 47 lbs)	ask a doctor													
children under 2 years (up to 33 lbs)	do not use													

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

VALERIE S PRATT
04/12/2017