



# NOT OFFICIAL SIZE/DIELINE



Label Size : 1.5" x 4.5"  
 Track : 19/02/2017, 22/02/2017



Art Production		DRUG FACTS INFO.	
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	-
Drug Facts	N/A	8 pt	-
(continued):	N/A	8 pt	-
Headings:	N/A	8 pt	6 pt
Subheadings:	N/A	6 pt	5 pt
Text:	N/A	6 pt	5 pt
Bullets:	N/A	5 pt (no larger)	4 pt
Leading:	N/A	+0.5 pt	5.4 pt
Characters Per Inch:	N/A	No more than 39	39



Carton Size : 3.0628" x 2.1253" x 1.5934"  
 Track : 19/02/2017, 22/02/2017

	Font	Size
Drug Facts	helvetica bold Oblique	10 pt.
Headings	helvetica bold Oblique	8 pt.
Subheadings	helvetica bold condensed	8 pt.
Body Text	helvetica medium condensed	6 pt.
Leading	N/A	0.5 pt
Bullets	Zapf Dingbats	5 pt.
Barline	N/A	2.5 pt.
Hairlines	N/A	0.5 pt.



NDC 69452-178-26

**Ibuprofen Capsules,  
200 mg**  
Pain Reliever/Fever Reducer (NSAID)

200 Capsules

**Drug Facts**

**Active ingredient (in each capsule)**  
Ibuprofen 200 mg (NSAID)\* .....Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches
  - the common cold
  - temporarily reduces fever
- minor pain of arthritis
- menstrual cramps
- toothache

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- skin redness
- asthma (wheezing)
- hives
- facial swelling
- rash
- shock
- skin redness
- blisters

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

- take with food or milk, if stomach upset occurs

Manufactured for: **Bionpharma Inc.**  
600 Alexander Road,  
Princeton, NJ 08540

Rev # 02/17  
Lot No.:  
Exp. Date:

**SEE HERE**

**Drug Facts (continued)**

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over:
  - take 1 capsule every 4 to 6 hours while symptoms persist
  - if pain or fever does not respond to 1 capsule, 2 capsules may be used
  - do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

**Other information**

- store at 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F)
- do not use if inner-child seal under bottle cap imprinted with "Sealed for Your Protection" is broken or missing
- read all warnings and directions before use. Keep carton.

**Inactive ingredients**  
FD&C Blue #1, gelatin, hypromellose, mannitol, polyethylene glycol, propylene glycol, povidone, purified water, sorbitan, sorbitol, titanium dioxide, vitamin E, polyethylene glycol succinate

**Questions or comments?** call toll free 1-888-235-2465

Label Size : 3" x 7"  
Track : 19/02/2017, 22/02/2017



	Font	Size
Drug Facts	helvetica bold Oblique	10 pt.
Headings	helvetica bold Oblique	8 pt.
Subheadings	helvetica bold condensed	8 pt.
Body Text	helvetica medium condensed	6 pt.
Leading	N/A	0.5 pt
Bullets	Zapf Dingbats	5 pt.
Barline	N/A	2.5 pt.
Hairlines	N/A	0.5 pt.

(b) (4)

9.4375 x 8.59375 Flat  
2.03125 x 2.03125 x 3.875

NDC 69452-178-26

## Ibuprofen Capsules, 200 mg Pain Reliever/Fever Reducer (NSAID)

### 200 Capsules

Softgels\*  
\*Liquid filled capsules

KEEP OUTER CARTON FOR COMPLETE  
WARNINGS AND PRODUCT INFORMATION

Raw # 02/17  
Lot No.:  
Exp. Date:

3  
6 9 4 5 2 1 7 8 2 6  
3

Manufactured for: **Bionpharma Inc.**  
600 Alexander Road, Princeton, NJ 08540  
Manufactured by: **Patheon Softgels Inc.**  
Highpoint, NC, 27285 USA.

**Drug Facts (continued)**

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ do not take more than directed ■ the smallest effective dose should be used  
 ■ adults and children 12 years and over: ■ take 1 capsule every 4 to 6 hours while symptoms persist  
 ■ if pain or fever does not respond to 1 capsule, 2 capsules may be used  
 ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor  
 ■ children under 12 years: ask a doctor

**Other information** ■ store at 20-25°C (68-77°F) ■ avoid high humidity and excessive heat above 40°C (104°F)  
 ■ do not use if Tamper-Evident seal under bottle cap imprinted with "Sealed for Your Protection" is broken or missing  
 ■ read all warnings and directions before use. Keep carton.

**Inactive ingredients** FD&C Blue #1, gelatin, hypromellose, mannitol, polyethylene glycol, propylene glycol, povidone, purified water, sorbitan, sorbitol, titanium dioxide, vitamin E polyethylene glycol succinate

**Questions or comments?** call toll free 1-888-235-2466

KEEP OUTER CARTON FOR COMPLETE  
WARNINGS AND PRODUCT INFORMATION

(b) (4)

**Drug Facts**

**Active ingredient (in each capsule)** .....Pain reliever/fever reducer  
 Ibuprofen 200 mg (NSAID)\* .....nonsteroidal anti-inflammatory drug

**Purposes**

**Uses**  
 ■ temporarily relieves minor aches and pains due to: ■ headache ■ muscular aches ■ minor pain of arthritis  
 ■ toothache ■ backache ■ the common cold ■ menstrual cramps ■ temporarily reduces fever

**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters  
 If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older  
 ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug  
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
 ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

KEEP OUTER CARTON FOR COMPLETE  
WARNINGS AND PRODUCT INFORMATION

(b) (4)

**Drug Facts (continued)**

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer  
 ■ right before or after heart surgery

**Ask a doctor before use if**  
 ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn  
 ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke  
 ■ you are taking a diuretic ■ you have problems or serious side effects from taking pain relievers or fever reducers

**Ask a doctor or pharmacist before use if you are**  
 ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug

**When using this product** ■ take with food or milk if stomach upset occurs

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding:  
 ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better  
 ■ you have symptoms of heart problems or stroke:  
 ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling  
 ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days  
 ■ redness or swelling is present in the painful area ■ any new symptoms appear

KEEP OUTER CARTON FOR COMPLETE  
WARNINGS AND PRODUCT INFORMATION

Colors



Font	Size
Drug Facts	10 pt.
Headings	8 pt.
Subheadings	8 pt.
Body Text	6 pt.
Leading	0.5 pt
Bullets	5 pt.
Barline	2.5 pt.
Hairlines	0.5 pt.

Carton Size : 2.031" x 2.031" x 3.875"  
Track : 19/02/2017, 22/02/2017