


# NOT OFFICIAL SIZE/DIELINE



Label Size : 4.625" x 1"  
Track : 14/09/2016, 23/02/2017-II

## Colors

(b) (4)



DRUG FACTS INFO.			
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	-
Drug Facts	N/A	8 pt	-
(continued):	N/A	8 pt	-
Headings:	N/A	8 pt	4.5 pt
Subheadings:	N/A	6 pt	4.5 pt
Text:	N/A	6 pt	4.5 pt
Bullets:	N/A	5 pt (no larger)	4.5 pt
Leading:	N/A	+0.5 pt	4.9 pt
Characters Per Inch:	N/A	No more than 39	45



Carton Size : 1.65" x 1.6" x 3"  
 Track : 14/09/2016, 08/02/2017,  
 23/02/2017-II

**MODIFIED DFL**

DRUG FACTS INFO.		
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED
Drug Facts Title:	N/A	>7.0 pt
Drug Facts	N/A	7 pt
(continued):	N/A	7 pt
Headings:	N/A	7 pt
Subheadings:	N/A	6 pt
Text:	N/A	6 pt
Bullets:	N/A	5 pt (no larger)
Leading:	N/A	+0.5 pt
Characters Per Inch:	N/A	No more than 39

(b) (4)

Colors



# NOT OFFICIAL SIZE/DIELINE



Label Size : 4.625" x 1"  
Track : 14/09/2016, 23/02/2017-II

## Colors




DRUG FACTS INFO.			
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	-
Drug Facts	N/A	8 pt	-
(continued):	N/A	8 pt	-
Headings:	N/A	8 pt	4.5 pt
Subheadings:	N/A	6 pt	4.5 pt
Text:	N/A	6 pt	4.5 pt
Bullets:	N/A	5 pt (no larger)	4.5 pt
Leading:	N/A	+0.5 pt	4.9 pt
Characters Per Inch:	N/A	No more than 39	45



Carton Size : 1.875" x 1.812" x 4.18"  
 Track : 14/09/2016, 08/02/2017,  
 23/02/2017-II

Colors



DRUG FACTS INFO.		PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A			10 pt
Drug Facts	N/A			9 pt
(continued):	N/A			9 pt
Headings:	N/A			8 pt
Subheadings:	N/A			6 pt
Text:	N/A			6 pt
Bullets:	N/A			5 pt (no larger)
Leadings:	N/A			+0.5 pt
Characters Per Inch:	N/A			6.5 pt
				No more than 39
				39

# NOT OFFICIAL SIZE/DIELINE

NDC 69452-179-15
Compare to the active ingredient  
of ALEVE® Liquid Gels\*\*\*

## Naproxen Sodium Capsules, 220 mg (NSAID)

**Pain Reliever/Fever Reducer**

**40 CAPSULES**

*Strength to last 12 hours*

Do not use if printed safety seal under cap is torn or missing.

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

Warnings: Heavy alcohol consumption may cause stomach bleeding, ulcers, and other complications. Symptoms include: heartburn, indigestion, stomach pain, nausea, vomiting, loss of appetite, weight loss, dizziness, headache, blurred vision, ringing in ears, nosebleeds, bleeding gums, skin rashes, itching, or hives. If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using the product • take more or for a longer time than directed. Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. American Heart Association (AHA) and American Stroke Association (ASA) advise that you should not take NSAIDs if you are at high risk for heart attack or stroke. Stop use and seek medical help right away if you experience any of the following signs or symptoms: chest pain, shortness of breath, sudden weakness or numbness in the face, arm, or leg, slurred speech, or changes in vision. If you experience any of the following signs or symptoms, stop use and seek medical help right away: black stools, have stomach pain (that does not go better), or if pain or fever gets worse. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. *Directions*: Adults and children 12 years and older: Take 1 capsule with water every 8 to 12 hours while symptoms last. For capsules in 24-hour period, do not take more than 12 capsules in 24 hours. *Warnings*: Do not use if you are allergic to naproxen sodium or any of the ingredients. *Other information*: • Contains 40 capsules. • USP-N 657 • This product is not for sale in the state of Florida. • **Boehr Ingelheim LLC, distributor of ALEVE® Liquid Gels** • Manufactured for: **Bayer Pharma Inc.** • 800 Aleander Road, Princeton, NJ 08540 • Lot/EXP

Label Size : 5.25" x 1.875"  
Track : 14/09/2016, 23/02/2017-II

### Colors



(b) (4)

DRUG FACTS INFO.			
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	-
Drug Facts	N/A	8 pt	-
(continued):	N/A	8 pt	-
Headings:	N/A	8 pt	5 pt
Subheadings:	N/A	6 pt	5 pt
Text:	N/A	6 pt	5 pt
Bullets:	N/A	5 pt (no larger)	4 pt
Leading:	N/A	+0.5 pt	5.4 pt
Characters Per Inch:	N/A	No more than 39	43



Carton Size : 1.875" x 1.812" x 4.18"  
 Track : 15/09/2016, 08/02/2017,  
 23/02/2017-II

Colors

(b) (4)

DRUG FACTS INFO.		PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A			10 pt
Drug Facts	N/A			9 pt
(continued):	N/A			9 pt
Headings:	N/A			8 pt
Subheadings:	N/A			6 pt
Text:	N/A			6 pt
Bullets:	N/A			5 pt (no larger)
Leadings:	N/A			+0.5 pt
Characters Per Inch:	N/A			6.5 pt
				No more than 39
				39

# NOT OFFICIAL SIZE/DIELINE

NDC 69452-179-16 *Compare to the active ingredient of ALEVE® Liquid Gels\*\*\**

**Naproxen Sodium Capsules, 220 mg (NSAID)**  
Pain Reliever/Fever Reducer

**50 CAPSULES**

*Strength to last 12 hours*

Do not use if printed safety seal under cap is torn or missing.

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

Warnings: Heavy alcohol consumption may cause stomach bleeding, ulcers, and other complications. Symptoms include: heartburn, indigestion, stomach pain, nausea, vomiting, diarrhea, constipation, bloating, gas, or belching. If any of these symptoms occur, stop use and seek medical help right away. Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using the product • take more or for a longer time than directed. Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. American Heart Association (AHA) and American Stroke Association (ASA) advise that you should not take NSAIDs if you are taking aspirin for your heart or stroke prevention. Stop use and seek medical help if you see black or tarry stools, blood in your stool, or blood in your urine. Stop use and seek medical help if you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not go better, or if pain or fever gets worse. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. *Directions:* Adults and children 12 years and older: Take 1 capsule with water every 8 to 12 hours while symptoms last. For other uses, ask your health care provider. *Caution:* Do not exceed 2 capsules in a 24-hour period. Do not take more than 12 capsules a day. *Other information:* Contains 50 capsules. Each capsule contains 220 mg of naproxen sodium (as the sodium salt). *©2017 Bayer Healthcare LLC, distributor of ALEVE® Liquid Gels. Manufactured for: Bayer Pharma Inc., 800 Alexander Road, Princeton, NJ 08540. Lot#EXP*

Label Size : 5.25" x 1.875"  
Track : 14/09/2016, 23/02/2017-II

## Colors



DRUG FACTS INFO.			
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	-
Drug Facts	N/A	8 pt	-
(continued):	N/A	8 pt	-
Headings:	N/A	8 pt	5 pt
Subheadings:	N/A	6 pt	5 pt
Text:	N/A	6 pt	5 pt
Bullets:	N/A	5 pt (no larger)	4 pt
Leading:	N/A	+0.5 pt	5.4 pt
Characters Per Inch:	N/A	No more than 39	43



Carton Size : 1.875" x 1.812" x 4.18"  
 Track : 15/09/2016, 08/02/2017,  
 23/02/2017-II

Colors

(b) (4)

DRUG FACTS INFO.		PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A			10 pt
Drug Facts	N/A			9 pt
(continued):	N/A			9 pt
Headings:	N/A			8 pt
Subheadings:	N/A			6 pt
Text:	N/A			6 pt
Bullets:	N/A			5 pt (no larger)
Leadings:	N/A			+0.5 pt
Characters Per Inch:	N/A			6.5 pt
				No more than 39
				39



# NOT OFFICIAL SIZE/DIELINE

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

NDC 69452-179-78 Compare to the active ingredient of ALEVE® Liquid Gels\*\*\*

## Naproxen Sodium Capsules, 220 mg (NSAID)

### Pain Reliever/Fever Reducer

EASY OPEN **ARTHRITIS CAP** **80 CAPSULES**

*Strength to last 12 hours*

**Do not use if printed safety seal under cap is torn or missing.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

**Warnings** Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin redness, rash, or blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains a non-steroidal anti-inflammatory drug (NSAID), which may cause a severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer or right before or after heart surgery. Ask a doctor before use if the stomach bleeding warning applies to you, you have a history of stomach problems (such as heartburn), you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke, you are taking a diuretic, you have problems or serious side effects from taking pain relievers or fever reducers, Ask a doctor or pharmacist before use if you are under a doctor's care for any serious condition or taking any other drug. **When using this product take with food or milk if stomach upset occurs.** Stop use and ask a doctor if you experience any of the following signs of stomach bleeding (flet/fat, vomit blood, have bloody or black stools, have stomach pain that does not get better), you have symptoms of heart problems or stroke (face pain, trouble breathing, weakness in one part or side of body, slurred speech, big swelling), pain gets worse or lasts more than 10 days, fever gets worse or lasts more than 3 days, redness or swelling is present in the genital area, any new symptoms appear, you have difficulty swallowing, or it feels like the capsule is stuck in your throat. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact Poison Control Center right away. **Directions** Do not take more than directed. The smallest effective dose should be used. Drink a full glass of water with each dose. If taken with food, this product may take longer to work. Adults and children 12 years and older: take 1 capsule every 8 to 12 hours while symptoms last. For the first dose you may take 2 capsules within the first hour. Do not exceed 2 capsules in any 8- to 12-hour period. Do not exceed 3 capsules in a 24-hour period. Children under 12 years: ask a doctor. Store at 20-25°C (68-77°F); avoid high humidity and excessive heat above 40°C (104°F). **Questions or comments?** 1-888-235-2466 (Mon-Fri 9AM - 5PM EST)

\*\*\*This product is not manufactured or distributed by Bayer Healthcare LLC, distributor of ALEVE® Liquid Gels. Manufactured for: Ampharma Inc., 600 Alexander Road, Princeton, NJ 08540

03/17  
LOVEXP

Label Size : 7.5" x 2.5"  
Track : 14/09/2016, 12/02/2017,  
23/02/2017-II, 01/03/2017

### Colors



DRUG FACTS INFO.			
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	-
Drug Facts	N/A	8 pt	-
(continued):	N/A	8 pt	-
Headings:	N/A	8 pt	6 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6.5 pt
Characters Per Inch:	N/A	No more than 39	39

NO COATING

**Naproxen Sodium Capsules, 220 mg (NSAID)**  
Pain Reliever/Fever Reducer

**80 CAPSULES**

Strength to last 12 hours

EASY OPEN ARTHRITIS CAP

Compare to the active ingredient of ALEVE® Liquid Gels \*\*\*

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

NDC 69452-179-78

NO COATING

**Drug Facts**

**Active ingredient (in each capsule)**  
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)\*.....Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Purposes**  
Pain reliever/fever reducer

**Uses**  
temporarily relieves minor aches and pains due to:  

- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

**Warnings**  
**Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  

- hives
- facial swelling
- asthma (wheezing)
- blisters
- shock
- skin reddening
- rash

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**  

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**  

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

**Ask a doctor or pharmacist before use if you are**  

- under a doctor's care for any serious condition
- taking any other drug

**When using this product**  

- take with food or milk if stomach upset occurs

**Drug Facts (continued)**

**Stop use and ask a doctor if**  

- you experience any of the following signs of stomach bleeding:  
  - feel faint
  - vomit blood
  - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:  
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

**adults and children 12 years and older:**  

- take 1 capsule every 8 to 12 hours while symptoms last
- for the first dose you may take 2 capsules within the first hour
- do not exceed 2 capsules in any 8- to 12-hour period
- do not exceed 3 capsules in a 24-hour period

**children under 12 years:**  

- ask a doctor

**Other information**  

- each capsule contains: sodium 20 mg
- store at 20-25°C (68-77°F) avoid high humidity and excessive heat above 40°C (104°F)
- read all directions and warnings before use. Keep carton.

**Inactive ingredients** FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

**Questions or comments?** 1-888-235-2466 (Mon-Fri 9AM - 5PM EST)

\*\*\*This product is not manufactured or distributed by Bayer Healthcare LLC, distributor of ALEVE® Liquid Gels.

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

**Naproxen Sodium Capsules, 220 mg (NSAID)**  
Pain Reliever/Fever Reducer

**80 CAPSULES**

Strength to last 12 hours

EASY OPEN ARTHRITIS CAP

Manufactured for:  
**Bionpharma Inc.**  
 600 Alexander Road,  
 Princeton, NJ 08540

3 69452 17978 9

Do not use if printed safety seal under cap is torn or missing.

Rev # 02/17  
 Lot No.:  
 Exp. Date:

LOT & EXP

**Colors**

(b) (4)

Carton Size : 2.625" x 2.625" x 4.875"  
 Track : 14/09/2016, 08/02/2017,  
 23/02/2017-II

DRUG FACTS INFO.			
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	10 pt
Drug Facts (continued):	N/A	8 pt	9 pt
Headings:	N/A	8 pt	8 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6.5 pt
Characters Per Inch:	N/A	No more than 39	39

# NOT OFFICIAL SIZE/DIELINE

NDC 69452-179-78
Compare to the active ingredient  
of ALEVE® Liquid Gels\*\*\*

## Naproxen Sodium Capsules, 220 mg (NSAID)

### Pain Reliever/Fever Reducer

#### 80 CAPSULES

*Strength to last 12 hours*

Do not use if printed safety seal under cap is torn or missing.

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

Warnings: Allergy reactions: Naproxen sodium may cause a severe allergic reaction, especially in people with asthma. Symptoms include: hives; facial swelling; itching; wheezing; shock; skin redness; rash; or blisters. If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding/warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using the product • take more or for a longer time than directed. Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. American Heart Association: If you are more than 65 years old, your doctor may suggest you stop taking NSAIDs. Do not use if you experience any of the following symptoms: chest pain, dizziness, fainting, or lightheadedness. Stop use and seek a doctor if you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not go better, or if pain or fever gets worse. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. *Directions:* Adults and children 12 years and older: Take 1 capsule with water every 8 to 12 hours while symptoms last. For other uses, ask a health professional. Do not exceed 2 capsules in a 24-hour period. Children under 12 years: See label directions. *Warnings:* Do not use if you are allergic to naproxen sodium, aspirin, or NSAIDs. \*\*\*This product is not comparable to Aleve® Liquid Gels. Manufactured for: Bion Pharma Inc., 800 Alexander Road, Princeton, NJ 08540. Lot#EXP 02/17

Label Size : 5.25" x 1.875"  
Track : 14/09/2016, 23/02/2017-II


### Colors



DRUG FACTS INFO.			
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	-
Drug Facts	N/A	8 pt	-
(continued):	N/A	8 pt	-
Headings:	N/A	8 pt	5 pt
Subheadings:	N/A	6 pt	5 pt
Text:	N/A	6 pt	5 pt
Bullets:	N/A	5 pt (no larger)	4 pt
Leading:	N/A	+0.5 pt	5.4 pt
Characters Per Inch:	N/A	No more than 39	43



Carton Size : 1.875" x 1.812" x 4.18"  
 Track : 15/09/2016, 08/02/2017,  
 23/02/2017-II  
**Colors**



DRUG FACTS INFO.		ACTUAL SIZE IN FILE
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED
Drug Facts Title:	N/A	10 pt
Drug Facts	N/A	8 pt
(continued):	N/A	8 pt
Headings:	N/A	8 pt
Subheadings:	N/A	6 pt
Text:	N/A	6 pt
Bullets:	N/A	5 pt (no larger)
Leading:	N/A	+0.5 pt
Characters Per Inch:	N/A	6.5 pt
		No more than 39
		39

# NOT OFFICIAL SIZE/DIELINE

NDC 69452-179-20      Compare to the active ingredient  
of ALEVE® Liquid Gels\*\*\*

**Naproxen Sodium  
Capsules, 220 mg (NSAID)**  
Pain Reliever/Fever Reducer

**100 CAPSULES**

*Strength to last 12 hours*

Do not use if printed safety seal under cap is torn or missing.

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

Warnings: Heavy alcohol consumption may cause stomach irritation, bleeding, or ulcers. Symptoms include: heartburn, indigestion, stomach pain, bloating, gas, or belching. If you experience any of these symptoms, stop using naproxen sodium capsules and call your doctor. If you experience any of these symptoms, stop using naproxen sodium capsules and call your doctor. If you experience any of these symptoms, stop using naproxen sodium capsules and call your doctor.

Precautions: Tell your doctor if you are taking any other medicines, especially blood thinners, aspirin, or other NSAIDs. Tell your doctor if you have ever had stomach ulcers, bleeding, or ulcers. Tell your doctor if you have ever had kidney disease, liver disease, or heart disease. Tell your doctor if you are pregnant or breastfeeding. Tell your doctor if you are taking any other medicines, especially blood thinners, aspirin, or other NSAIDs. Tell your doctor if you have ever had stomach ulcers, bleeding, or ulcers. Tell your doctor if you have ever had kidney disease, liver disease, or heart disease. Tell your doctor if you are pregnant or breastfeeding. Tell your doctor if you are taking any other medicines, especially blood thinners, aspirin, or other NSAIDs.

Side Effects: The most common side effects of naproxen sodium capsules are: heartburn, indigestion, stomach pain, bloating, gas, or belching. Other side effects include: dizziness, headache, drowsiness, and dry mouth. Tell your doctor if you experience any of these side effects. Tell your doctor if you experience any of these side effects. Tell your doctor if you experience any of these side effects.

Directions: Take 1 capsule with water every 8 to 12 hours while symptoms last. Do not exceed 2 capsules in a 24-hour period. Do not take more than 12 capsules in a 30-day period. Do not take more than 12 capsules in a 30-day period. Do not take more than 12 capsules in a 30-day period.

Warnings for Children: Do not give naproxen sodium capsules to children under 12 years of age. Do not give naproxen sodium capsules to children under 12 years of age. Do not give naproxen sodium capsules to children under 12 years of age.

Manufactured for: Bayer HealthCare LLC, distributor of ALEVE® Liquid Gels  
Parsippany, NJ 07054  
Lot#EXP

02/17

Label Size : 5.25" x 1.875"  
Track : 14/09/2016, 23/02/2017-II

### Colors



(b) (4)

DRUG FACTS INFO.			
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	-
Drug Facts	N/A	8 pt	-
(continued):	N/A	8 pt	-
Headings:	N/A	8 pt	5 pt
Subheadings:	N/A	6 pt	5 pt
Text:	N/A	6 pt	5 pt
Bullets:	N/A	5 pt (no larger)	4 pt
Leading:	N/A	+0.5 pt	5.4 pt
Characters Per Inch:	N/A	No more than 39	43



Carton Size : 1.875" x 1.812" x 4.18"  
 Track : 15/09/2016, 08/02/2017,  
 23/02/2017-II

Colors



DRUG FACTS INFO.		ACTUAL SIZE IN FILE
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED
Drug Facts Title:	N/A	10 pt
Drug Facts	N/A	8 pt
(continued):	N/A	8 pt
Headings:	N/A	8 pt
Subheadings:	N/A	6 pt
Text:	N/A	6 pt
Bullets:	N/A	5 pt (no larger)
Leading:	N/A	+0.5 pt
Characters Per Inch:	N/A	No more than 39
		39

# NOT OFFICIAL SIZE/DIELINE

NDC 69452-179-79 *Compare to the active ingredient of ALEVE® Liquid Gels\*\*\**

**Naproxen Sodium Capsules, 220 mg (NSAID)**  
Pain Reliever/Fever Reducer

**160 CAPSULES**

*Strength to last 12 hours*

Do not use if printed safety seal under cap is torn or missing.

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

Warnings: Heavy alcohol consumption may cause stomach irritation, bleeding, or ulcers. Symptoms include: heartburn, indigestion, stomach pain, stomach bleeding, black or bloody stools, or blood in urine. Stop use and seek medical help right away. Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using the product • take more or for a longer time than directed. Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. American Heart Association (AHA) and American Stroke Association (ASA) advise that you should not take NSAIDs if you are at high risk for heart attack or stroke. Stop use and seek medical help right away if you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not go better, or if pain or fever gets worse. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. *Directions:* Adults and children 12 years and older: Take 1 capsule with water every 8 to 12 hours while symptoms last. For 2 capsules in a 24-hour period. Do not take more than 12 capsules in 10 days. *Warnings:* See Warnings and Precautions on page 2. *Directions:* See Directions on page 2. *Other information:* See Other information on page 2. *© 2017 Bayer Healthcare LLC, distributor of ALEVE® Liquid Gels. Manufactured for: Bayer Pharma Inc., 800 Aleander Road, Princeton, NJ 08540. Lot/EXP*

Label Size : 5.25" x 1.875"  
Track : 14/09/2016, 23/02/2017-II

## Colors



DRUG FACTS INFO.			
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	-
Drug Facts	N/A	8 pt	-
(continued):	N/A	8 pt	-
Headings:	N/A	8 pt	5 pt
Subheadings:	N/A	6 pt	5 pt
Text:	N/A	6 pt	5 pt
Bullets:	N/A	5 pt (no larger)	4 pt
Leading:	N/A	+0.5 pt	5.4 pt
Characters Per Inch:	N/A	No more than 39	43



Carton Size : 1.875" x 1.812" x 4.18"  
 Track : 15/09/2016, 08/02/2017,  
 23/02/2017-II

Colors

(b) (4)

DRUG FACTS INFO.		PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A			10 pt
Drug Facts	N/A			8 pt
(continued):	N/A			8 pt
Headings:	N/A			8 pt
Subheadings:	N/A			6 pt
Text:	N/A			6 pt
Bullets:	N/A			5 pt (no larger)
Leadings:	N/A			+0.5 pt
Characters Per Inch:	N/A			6.5 pt
				No more than 39
				39




# NOT OFFICIAL SIZE/DIELINE

**Do not use if printed safety seal under cap is torn or missing.**  
 \*\*\*This product is not manufactured or distributed by Bayer Healthcare LLC, distributor of ALEVE® Liquid Gels.

Manufactured for:  
**Bionpharma Inc.**  
 800 Alexander Road,  
 Princeton, NJ 08540

02/17



3 69452 17979 6

Lot/EXP

NDC 69452-179-79 Compare to the active ingredient  
of ALEVE® Liquid Gels\*\*\*

## Naproxen Sodium Capsules, 220 mg (NSAID)

### Pain Reliever/Fever Reducer

160 CAPSULES

Strength to last 12 hours

PEEL HERE

**Drug Facts**

**Active ingredient (in each capsule)**  
 Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)\* .....Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Purposes**

**Uses**

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis    ● muscular aches    ● backache
- menstrual cramps        ● headache        ● toothache    ● the common cold
- temporarily reduces fever

**Warnings**

**Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives                      ● facial swelling        ● asthma (wheezing)    ● shock
- skin reddening        ● rash                    ● blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**TOP**




**BEHIND TOP**

<p><b>Drug Facts (continued)</b></p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>● if you have ever had an allergic reaction to any other pain reliever/fever reducer</li> <li>● right before or after heart surgery</li> </ul> <p><b>Ask a doctor before use if</b></p> <ul style="list-style-type: none"> <li>● stomach bleeding warning applies to you</li> <li>● you have a history of stomach problems, such as heartburn</li> <li>● you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke</li> <li>● you are taking a diuretic</li> <li>● you have problems or serious side effects from taking pain relievers or fever reducers</li> </ul> <p><b>Ask a doctor or pharmacist before use if you are</b></p> <ul style="list-style-type: none"> <li>● under a doctor's care for any serious condition</li> <li>● taking any other drug</li> </ul> <p><b>When using this product</b></p> <ul style="list-style-type: none"> <li>● take with food or milk if stomach upset occurs</li> </ul> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>● you experience any of the following signs of stomach bleeding:                     <ul style="list-style-type: none"> <li>● feel faint</li> <li>● vomit blood</li> <li>● have bloody or black stools</li> <li>● have stomach pain that does not get better</li> </ul> </li> <li>● you have symptoms of heart problems or stroke:                     <ul style="list-style-type: none"> <li>● chest pain</li> <li>● trouble breathing</li> <li>● slurred speech</li> <li>● weakness in one part or side of body</li> <li>● leg swelling</li> </ul> </li> </ul>	<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>● pain gets worse or lasts more than 10 days</li> <li>● fever gets worse or lasts more than 3 days</li> <li>● redness or swelling is present in the painful area</li> <li>● any new symptoms appear</li> <li>● you have difficulty swallowing</li> <li>● it feels like the capsule is stuck in your throat</li> </ul> <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. <b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>● do not take more than directed</li> <li>● the smallest effective dose should be used</li> <li>● drink a full glass of water with each dose</li> <li>● if taken with food, this product may take longer to work</li> </ul> <table style="width: 100%; border: none;"> <tr> <td style="width: 50%; border: none;">adults and children 12 years and older:</td> <td style="border: none;"> <ul style="list-style-type: none"> <li>● take 1 capsule every 8 to 12 hours while symptoms last</li> <li>● for the first dose you may take 2 capsules within the first hour</li> <li>● do not exceed 2 capsules in any 8- to 12-hour period</li> <li>● do not exceed 3 capsules in a 24-hour period</li> </ul> </td> </tr> <tr> <td style="border: none;">children under 12 years:</td> <td style="border: none;"> <ul style="list-style-type: none"> <li>● ask a doctor</li> </ul> </td> </tr> </table>	adults and children 12 years and older:	<ul style="list-style-type: none"> <li>● take 1 capsule every 8 to 12 hours while symptoms last</li> <li>● for the first dose you may take 2 capsules within the first hour</li> <li>● do not exceed 2 capsules in any 8- to 12-hour period</li> <li>● do not exceed 3 capsules in a 24-hour period</li> </ul>	children under 12 years:	<ul style="list-style-type: none"> <li>● ask a doctor</li> </ul>	<p><b>Drug Facts (continued)</b></p> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>● each capsule contains: sodium 20 mg</li> <li>● store at 20-25°C (68-77°F) avoid high humidity and excessive heat above 40°C (104°F)</li> <li>● read all directions and warnings before use.</li> </ul> <p><b>Inactive ingredients</b> FD&amp;C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol</p> <p><b>Questions or comments?</b> 1-888-235-2466 (Mon-Fri 9AM - 5PM EST)</p>
adults and children 12 years and older:	<ul style="list-style-type: none"> <li>● take 1 capsule every 8 to 12 hours while symptoms last</li> <li>● for the first dose you may take 2 capsules within the first hour</li> <li>● do not exceed 2 capsules in any 8- to 12-hour period</li> <li>● do not exceed 3 capsules in a 24-hour period</li> </ul>					
children under 12 years:	<ul style="list-style-type: none"> <li>● ask a doctor</li> </ul>					

**BASE**

Label Size : 8.625" x 3.25"  
 Track : 15/09/2016, 08/02/2017,  
 23/02/2017-II

**Colors**



DRUG FACTS INFO.			
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	10 pt
Drug Facts	N/A	8 pt	9 pt
(continued):	N/A	8 pt	9 pt
Headings:	N/A	8 pt	8 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6.5 pt
Characters Per Inch:	N/A	No more than 39	36