

## HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use PREZISTA safely and effectively. See Full Prescribing Information for PREZISTA.

PREZISTA (darunavir) oral suspension

PREZISTA (darunavir) tablet, for oral use

Initial U.S. Approval: 2006

## RECENT MAJOR CHANGES

Contraindications (4)

06/2017

## INDICATIONS AND USAGE

PREZISTA is a human immunodeficiency virus (HIV-1) protease inhibitor indicated for the treatment of HIV-1 infection in adult and pediatric patients 3 years of age and older. PREZISTA must be co-administered with ritonavir (PREZISTA/ritonavir) and with other antiretroviral agents. (1)

## DOSAGE AND ADMINISTRATION

- Testing:
  - In treatment-experienced patients, treatment history genotypic and/or phenotypic testing is recommended prior to initiation of therapy with PREZISTA/ritonavir to assess drug susceptibility of the HIV-1 virus (2.1, 12.4)
  - Monitor serum liver chemistry tests before and during therapy with PREZISTA/ritonavir. (2.1, 2.2, 5.2)
- Treatment-naïve adult patients and treatment-experienced adult patients with no darunavir resistance associated substitutions: 800 mg (one 800 mg tablet) taken with ritonavir 100 mg once daily and with food. (2.3)
- Treatment-experienced adult patients with at least one darunavir resistance associated substitution: 600 mg (one 600 mg tablet) taken with ritonavir 100 mg twice daily and with food. (2.3)
- Pregnant patients: 600 mg (one 600 mg tablet) taken with ritonavir 100 mg twice daily and with food. (2.4)
- Pediatric patients (3 to less than 18 years of age and weighing at least 10 kg): dosage of PREZISTA and ritonavir is based on body weight and should not exceed the adult dose. PREZISTA should be taken with ritonavir and with food. (2.5)
- PREZISTA/ritonavir is not recommended for use in patients with severe hepatic impairment. (2.6)

## DOSAGE FORMS AND STRENGTHS

- Oral suspension: 100 mg per mL (3)
- Tablets: 75 mg, 150 mg, 600 mg, and 800 mg (3)

## CONTRAINDICATIONS

- Co-administration of PREZISTA/ritonavir is contraindicated with drugs that are highly dependent on CYP3A for clearance and for which elevated plasma concentrations are associated with serious and/or life-threatening events (narrow therapeutic index). (4)

## WARNINGS AND PRECAUTIONS

- Drug-induced hepatitis (e.g., acute hepatitis, cytolytic hepatitis) has been reported with PREZISTA/ritonavir. Monitor liver function before and during therapy, especially in patients with underlying chronic hepatitis, cirrhosis, or in patients who have pre-treatment elevations of transaminases. Post-marketing cases of liver injury, including some fatalities, have been reported. (5.2)
- Skin reactions ranging from mild to severe, including Stevens-Johnson Syndrome, toxic epidermal necrolysis, drug rash with eosinophilia and systemic symptoms and acute generalized exanthematous pustulosis, have been reported. Discontinue treatment if severe reaction develops. (5.3)
- Use with caution in patients with a known sulfonamide allergy. (5.4)
- Patients may develop new onset diabetes mellitus or hyperglycemia. Initiation or dose adjustments of insulin or oral hypoglycemic agents may be required. (5.6)
- Patients may develop redistribution/accumulation of body fat or immune reconstitution syndrome. (5.7, 5.8)
- Patients with hemophilia may develop increased bleeding events. (5.9)
- PREZISTA/ritonavir is not recommended in pediatric patients below 3 years of age in view of toxicity and mortality observed in juvenile rats dosed with darunavir up to days 23 to 26 of age. (5.10)

## ADVERSE REACTIONS

- The most common clinical adverse drug reactions to PREZISTA/ritonavir (incidence greater than or equal to 5%) of at least moderate intensity (greater than or equal to Grade 2) were diarrhea, nausea, rash, headache, abdominal pain and vomiting. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Janssen Products, LP at 1-800-JANSSEN (1-800-526-7736) or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

## DRUG INTERACTIONS

- Co-administration of PREZISTA/ritonavir with other drugs can alter the concentrations of other drugs and other drugs may alter the concentrations of darunavir. The potential drug-drug interactions must be considered prior to and during therapy. (4, 5.5, 7, 12.3)

## USE IN SPECIFIC POPULATIONS

- Pregnancy: Total darunavir exposures were generally lower during pregnancy compared to postpartum period. The reduction in darunavir exposures during pregnancy were greater for once daily dosing compared to the twice daily dosing regimen. (8.1, 12.3)
- Lactation: Women infected with HIV should be instructed not to breastfeed due to the potential for HIV transmission. (8.2)
- Pediatrics: Not recommended for patients less than 3 years of age. (8.4)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 06/2017

## FULL PRESCRIBING INFORMATION: CONTENTS\*

### 1 INDICATIONS AND USAGE

### 2 DOSAGE AND ADMINISTRATION

- 2.1 Testing Prior to Initiation of PREZISTA/ritonavir
- 2.2 Monitoring During Treatment with PREZISTA/ritonavir
- 2.3 Recommended Dosage in Adult Patients
- 2.4 Recommended Dosage During Pregnancy
- 2.5 Recommended Dosage in Pediatric Patients (age 3 to less than 18 years)
- 2.6 Not Recommended in Patients with Severe Hepatic Impairment

### 3 DOSAGE FORMS AND STRENGTHS

### 4 CONTRAINDICATIONS

### 5 WARNINGS AND PRECAUTIONS

- 5.1 Importance of Co-administration with Ritonavir
- 5.2 Hepatotoxicity
- 5.3 Severe Skin Reactions
- 5.4 Sulfa Allergy
- 5.5 Risk of Serious Adverse Reactions due to Drug Interactions
- 5.6 Diabetes Mellitus/Hyperglycemia
- 5.7 Fat Redistribution
- 5.8 Immune Reconstitution Syndrome

- 5.9 Hemophilia

- 5.10 Not Recommended in Pediatric Patients Below 3 Years of Age

### 6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

### 7 DRUG INTERACTIONS

- 7.1 Potential for PREZISTA/ritonavir to Affect Other Drugs
- 7.2 Potential for Other Drugs to Affect Darunavir
- 7.3 Established and Other Potentially Significant Drug Interactions
- 7.4 Drugs without Clinically Significant Interactions with PREZISTA

### 8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.3 Females and Males of Reproductive Potential
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Hepatic Impairment
- 8.7 Renal Impairment

### 10 OVERDOSAGE

### 11 DESCRIPTION

**12 CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics
- 12.4 Microbiology

**13 NONCLINICAL TOXICOLOGY**

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

\*Sections or subsections omitted from the full prescribing information are not listed.

---

**14 CLINICAL STUDIES**

- 14.1 Description of Adult Clinical Trials
- 14.2 Treatment-Naïve Adult Subjects
- 14.3 Treatment-Experienced Adult Subjects
- 14.4 Pediatric Patients

**16 HOW SUPPLIED/STORAGE AND HANDLING**

**17 PATIENT COUNSELING INFORMATION**

## FULL PRESCRIBING INFORMATION

### 1 INDICATIONS AND USAGE

PREZISTA<sup>®</sup>, co-administered with ritonavir (PREZISTA/ritonavir), in combination with other antiretroviral agents, is indicated for the treatment of human immunodeficiency virus (HIV-1) infection in adult and pediatric patients 3 years of age and older [*see Use in Specific Populations (8.4) and Clinical Studies (14)*].

### 2 DOSAGE AND ADMINISTRATION

#### 2.1 Testing Prior to Initiation of PREZISTA/ritonavir

In treatment-experienced patients, treatment history, genotypic and/or phenotypic testing is recommended to assess drug susceptibility of the HIV-1 virus [*see Microbiology (12.4)*]. Refer to *Dosage and Administration (2.3), (2.4) and (2.5)* for dosing recommendations.

Appropriate laboratory testing such as serum liver biochemistries should be conducted prior to initiating therapy with PREZISTA/ritonavir [*see Warnings and Precautions (5.2)*].

#### 2.2 Monitoring During Treatment with PREZISTA/ritonavir

Patients with underlying chronic hepatitis, cirrhosis, or in patients who have pre-treatment elevations of transaminases should be monitored for elevation in serum liver biochemistries, especially during the first several months of PREZISTA/ritonavir treatment [*see Warnings and Precautions (5.2)*].

#### 2.3 Recommended Dosage in Adult Patients

PREZISTA must be co-administered with ritonavir to exert its therapeutic effect. Failure to correctly co-administer PREZISTA with ritonavir will result in plasma levels of darunavir that will be insufficient to achieve the desired antiviral effect and will alter some drug interactions.

Patients who have difficulty swallowing PREZISTA tablets can use the 100 mg per mL PREZISTA oral suspension.

##### Treatment-Naïve Adult Patients

The recommended oral dose of PREZISTA is 800 mg (one 800 mg tablet or 8 mL of the oral suspension) taken with ritonavir 100 mg (one 100 mg tablet or capsule or 1.25 mL of a 80 mg per mL ritonavir oral solution) once daily and with food. An 8 mL PREZISTA dose should be taken as two 4 mL administrations with the included oral dosing syringe.

##### Treatment-Experienced Adult Patients

The recommended oral dosage for treatment-experienced adult patients is summarized in Table 1.

Baseline genotypic testing is recommended for dose selection. However, when genotypic testing is not feasible, PREZISTA 600 mg taken with ritonavir 100 mg twice daily is recommended.

**Table 1: Recommended PREZISTA/ritonavir Dosage in Treatment-Experienced Adult Patients**

Baseline Resistance	Formulation and Recommended Dosing	
	PREZISTA tablets with ritonavir tablets or capsule	PREZISTA oral suspension (100 mg/mL) with ritonavir oral solution (80 mg/mL)
With no darunavir resistance associated substitutions*	One 800 mg PREZISTA tablet with one 100 mg ritonavir tablet/capsule, taken once daily with food	8 mL <sup>†</sup> PREZISTA oral suspension with 1.25 mL ritonavir oral solution, taken once daily with food
With at least one darunavir resistance associated substitutions*, or with no baseline resistance information	One 600 mg PREZISTA tablet with one 100 mg ritonavir tablet/capsule, taken twice daily with food	6 mL PREZISTA oral suspension with 1.25 mL ritonavir oral solution, taken twice daily with food

\* V11I, V32I, L33F, I47V, I50V, I54L, I54M, T74P, L76V, I84V and L89V

† An 8 mL darunavir dose should be taken as two 4 mL administrations with the included oral dosing syringe

## 2.4 Recommended Dosage During Pregnancy

The recommended dosage in pregnant patients is PREZISTA 600 mg taken with ritonavir 100 mg twice daily with food.

PREZISTA 800 mg taken with ritonavir 100 mg once daily should only be considered in certain pregnant patients who are already on a stable PREZISTA 800 mg with ritonavir 100 mg once daily regimen prior to pregnancy, are virologically suppressed (HIV-1 RNA less than 50 copies per mL), and in whom a change to twice daily PREZISTA 600 mg with ritonavir 100 mg may compromise tolerability or compliance.

## 2.5 Recommended Dosage in Pediatric Patients (age 3 to less than 18 years)

Healthcare professionals should pay special attention to accurate dose selection of PREZISTA, transcription of the medication order, dispensing information and dosing instruction to minimize risk for medication errors, overdose, and underdose.

Prescribers should select the appropriate dose of PREZISTA/ritonavir for each individual child based on body weight (kg) and should not exceed the recommended dose for adults.

Before prescribing PREZISTA, children weighing greater than or equal to 15 kg should be assessed for the ability to swallow tablets. If a child is unable to reliably swallow a tablet, the use of PREZISTA oral suspension should be considered.

The recommended dose of PREZISTA/ritonavir for pediatric patients (3 to less than 18 years of age and weighing at least 10 kg is based on body weight (see Tables 2, 3, 4, and 5) and should not exceed the recommended adult dose. PREZISTA should be taken with ritonavir and with food.

The recommendations for the PREZISTA/ritonavir dosage regimens were based on pediatric clinical trial data and population pharmacokinetic modeling and simulation [see *Use in Specific Populations* (8.4) and *Clinical Pharmacology* (12.3)].

## Dosing Recommendations for Treatment-Naïve Pediatric Patients or Antiretroviral Treatment-Experienced Pediatric Patients with No Darunavir Resistance Associated Substitutions

### *Pediatric patients weighing at least 10 kg but less than 15 kg*

The weight-based dose in antiretroviral treatment-naïve pediatric patients or antiretroviral treatment-experienced pediatric patients with no darunavir resistance associated substitutions is PREZISTA 35 mg/kg once daily with ritonavir 7 mg/kg once daily using the following table:

**Table 2: Recommended Dose for Pediatric Patients Weighing 10 kg to Less Than 15 kg Who are Treatment-Naïve or Treatment-Experienced with No Darunavir Resistance Associated Substitutions\***

Body weight (kg)	Formulation: PREZISTA oral suspension (100 mg/mL) and ritonavir oral solution (80 mg/mL)
	Dose: once daily with food
Greater than or equal to 10 kg to less than 11 kg	PREZISTA 3.6 mL <sup>†</sup> (350 mg) with ritonavir 0.8 mL (64 mg)
Greater than or equal to 11 kg to less than 12 kg	PREZISTA 4 mL <sup>†</sup> (385 mg) with ritonavir 0.8 mL (64 mg)
Greater than or equal to 12 kg to less than 13 kg	PREZISTA 4.2 mL (420 mg) with ritonavir 1 mL (80 mg)
Greater than or equal to 13 kg to less than 14 kg	PREZISTA 4.6 mL <sup>†</sup> (455 mg) with ritonavir 1 mL (80 mg)
Greater than or equal to 14 kg to less than 15 kg	PREZISTA 5 mL <sup>†</sup> (490 mg) with ritonavir 1.2 mL (96 mg)

\* darunavir resistance associated substitutions: V11I, V32I, L33F, I47V, I50V, I54M, I54L, T74P, L76V, I84V and L89V

† The 350 mg, 385 mg, 455 mg and 490 mg darunavir dose for the specified weight groups were rounded up for suspension dosing convenience to 3.6 mL, 4 mL, 4.6 mL and 5 mL, respectively.

### *Pediatric patients weighing at least 15 kg*

Pediatric patients weighing at least 15 kg can be dosed with PREZISTA oral tablet(s) or suspension using the following table:

**Table 3: Recommended Dose for Pediatric Patients Weighing At Least 15 kg Who are Treatment-Naïve or Treatment-Experienced with No Darunavir Resistance Associated Substitutions\***

Body weight (kg)	Formulation: PREZISTA tablet(s) and ritonavir capsules or tablets (100 mg)	Formulation: PREZISTA oral suspension (100 mg/mL) and ritonavir oral solution (80 mg/mL)
	Dose: once daily with food	Dose: once daily with food
Greater than or equal to 15 kg to less than 30 kg	PREZISTA 600 mg with ritonavir 100 mg	PREZISTA 6 mL (600 mg) with ritonavir 1.25 mL (100 mg)
Greater than or equal to 30 kg to less than 40 kg	PREZISTA 675 mg with ritonavir 100 mg	PREZISTA 6.8 mL <sup>†‡</sup> (675 mg) with ritonavir 1.25 mL (100 mg)
Greater than or equal to 40 kg	PREZISTA 800 mg with ritonavir 100 mg	PREZISTA 8 mL <sup>†</sup> (800 mg) with ritonavir 1.25 mL (100 mg)

\* darunavir resistance associated substitutions: V11I, V32I, L33F, I47V, I50V, I54M, I54L, T74P, L76V, I84V and L89V

† The 675 mg dose using darunavir tablets for this weight group is rounded up to 6.8 mL for suspension dosing convenience.

‡ The 6.8 mL and 8 mL darunavir dose should be taken as two (3.4 mL or 4 mL respectively) administrations with the included oral dosing syringe

## Dosing Recommendations for Treatment-Experienced Pediatric Patients with At Least One Darunavir Resistance Associated Substitutions

### *Pediatric patients weighing at least 10 kg but less than 15 kg*

The weight-based dose in antiretroviral treatment-experienced pediatric patients with at least one darunavir resistance associated substitution is PREZISTA 20 mg/kg twice daily with ritonavir 3 mg/kg twice daily using the following table:

**Table 4: Recommended Dose for Pediatric Patients Weighing 10 kg to Less Than 15 kg Who are Treatment-Experienced with At Least One Darunavir Resistance Associated Substitution\***

Body weight (kg)	Formulation: PREZISTA oral suspension (100 mg/mL) and ritonavir oral solution (80 mg/mL)
	Dose: twice daily with food
Greater than or equal to 10 kg to less than 11 kg	PREZISTA 2 mL (200 mg) with ritonavir 0.4 mL (32 mg)
Greater than or equal to 11 kg to less than 12 kg	PREZISTA 2.2 mL (220 mg) with ritonavir 0.4 mL (32 mg)
Greater than or equal to 12 kg to less than 13 kg	PREZISTA 2.4 mL (240 mg) with ritonavir 0.5 mL (40 mg)
Greater than or equal to 13 kg to less than 14 kg	PREZISTA 2.6 mL (260 mg) with ritonavir 0.5 mL (40 mg)
Greater than or equal to 14 kg to less than 15 kg	PREZISTA 2.8 mL (280 mg) with ritonavir 0.6 mL (48 mg)

\* darunavir resistance associated substitutions: V11I, V32I, L33F, I47V, I50V, I54M, I54L, T74P, L76V, I84V and L89V

### *Pediatric patients weighing at least 15 kg*

Pediatric patients weighing at least 15 kg can be dosed with PREZISTA oral tablet(s) or suspension using the following table:

**Table 5: Recommended Dose for Pediatric Patients Weighing At Least 15 kg Who are Treatment-Experienced with At Least One Darunavir Resistance Associated Substitution\***

Body weight (kg)	Formulation: PREZISTA tablet(s) and ritonavir tablets, capsules (100 mg) or oral solution (80 mg/mL)	Formulation: PREZISTA oral suspension (100 mg/mL) and ritonavir oral solution (80 mg/mL)
	Dose: twice daily with food	Dose: twice daily with food
Greater than or equal to 15 kg to less than 30 kg	PREZISTA 375 mg with ritonavir 0.6 mL (48 mg)	PREZISTA 3.8 mL (375 mg) <sup>†</sup> with ritonavir 0.6 mL (48 mg)
Greater than or equal to 30 kg to less than 40 kg	PREZISTA 450 mg with ritonavir 0.75 mL (60 mg)	PREZISTA 4.6 mL (450 mg) <sup>†</sup> with ritonavir 0.75 mL (60 mg)
Greater than or equal to 40 kg	PREZISTA 600 mg with ritonavir 100 mg	PREZISTA 6 mL (600 mg) with ritonavir 1.25 mL (100 mg)

\* darunavir resistance associated substitutions: V11I, V32I, L33F, I47V, I50V, I54M, I54L, T74P, L76V, I84V and L89V

<sup>†</sup> The 375 mg and 450 mg dose using darunavir tablets for this weight group is rounded up to 3.8 mL and 4.6 mL for suspension dosing convenience.

The use of PREZISTA/ritonavir in pediatric patients below 3 years of age is not recommended [see *Warnings and Precautions (5.10) and Use in Specific Populations (8.4)*].

## **2.6 Not Recommended in Patients with Severe Hepatic Impairment**

No dosage adjustment is required in patients with mild or moderate hepatic impairment. No data are available regarding the use of PREZISTA/ritonavir when co-administered to subjects with severe hepatic impairment; therefore, PREZISTA/ritonavir is not recommended for use in patients with severe hepatic impairment [see *Use in Specific Populations (8.6) and Clinical Pharmacology (12.3)*].

## **3 DOSAGE FORMS AND STRENGTHS**

### PREZISTA Oral Suspension

PREZISTA 100 mg per mL is supplied as a white to off-white opaque suspension for oral use, containing darunavir ethanolate equivalent to 100 mg of darunavir per mL of suspension.

## PREZISTA Tablets

- 75 mg: white, caplet-shaped, film-coated tablets containing darunavir ethanolate equivalent to 75 mg of darunavir. Each tablet is debossed with “75” on one side and “TMC” on the other side.
- 150 mg: white, oval-shaped, film-coated tablets containing darunavir ethanolate equivalent to 150 mg of darunavir. Each tablet is debossed with “150” on one side and “TMC” on the other side.
- 600 mg: orange, oval-shaped, film-coated tablets containing darunavir ethanolate equivalent to 600 mg of darunavir. Each tablet is debossed with “600MG” on one side and “TMC” on the other side.
- 800 mg: dark red, oval-shaped, film-coated tablets containing darunavir ethanolate equivalent to 800 mg of darunavir. Each tablet is debossed with “800” on one side and “T” on the other side.

## **4 CONTRAINDICATIONS**

Co-administration of PREZISTA/ritonavir is contraindicated with drugs that are highly dependent on CYP3A for clearance and for which elevated plasma concentrations are associated with serious and/or life-threatening events (narrow therapeutic index). These drugs and other contraindicated drugs (which may lead to reduced efficacy of darunavir) are listed in Table 6 [see *Drug Interactions* (7.3)]. Due to the need for co-administration of PREZISTA with ritonavir, please refer to ritonavir prescribing information for a description of ritonavir contraindications.

**Table 6: Drugs That Are Contraindicated With PREZISTA/ritonavir**

<b>Drug class</b>	<b>Drugs within class that are contraindicated with PREZISTA/ritonavir</b>	<b>Clinical comment</b>
Alpha 1-adrenoreceptor antagonist	alfuzosin	Potential for serious and/or life-threatening reactions such as hypotension.
Antianginal	ranolazine	Potential for serious and/or life-threatening reactions.
Antiarrhythmic	dronedarone	Potential for serious and/or life-threatening reactions such as cardiac arrhythmias.
Anti-gout	colchicine	Potential for serious and/or life-threatening reactions in patients with renal and/or hepatic impairment.
Antimycobacterial	rifampin	Rifampin is a potent inducer of CYP450 metabolism. PREZISTA/ritonavir should not be used in combination with rifampin, as this may cause significant decreases in darunavir plasma concentrations. This may result in loss of therapeutic effect to PREZISTA.
Antipsychotics	lurasidone	Potential for serious and/or life-threatening reactions.
	pimozide	Potential for serious and/or life-threatening reactions such as cardiac arrhythmias.
Ergot derivatives	dihydroergotamine, ergotamine, methylergonovine	Potential for serious and/or life-threatening events such as acute ergot toxicity characterized by

		peripheral vasospasm and ischemia of the extremities and other tissues.
GI motility agent	cisapride	Potential for serious and/or life-threatening reactions such as cardiac arrhythmias.
Sedative/hypnotics	Orally administered midazolam, triazolam	Triazolam and orally administered midazolam are extensively metabolized by CYP3A. Co-administration of triazolam or orally administered midazolam with PREZISTA/ritonavir may cause large increases in the concentrations of these benzodiazepines. Potential for serious and/or life-threatening events such as prolonged or increased sedation or respiratory depression.
Herbal product	St. John's wort ( <i>Hypericum perforatum</i> )	Patients taking PREZISTA/ritonavir should not use products containing St. John's wort because co-administration may result in reduced plasma concentrations of darunavir. This may result in loss of therapeutic effect and development of resistance.
Hepatitis C direct-acting antiviral	elbasvir/grazoprevir	Potential for the increased risk of alanine transaminase (ALT) elevations.
HMG-CoA reductase inhibitors	lovastatin, simvastatin	Potential for serious reactions such as myopathy including rhabdomyolysis. For dosing recommendation regarding atorvastatin, pravastatin and rosuvastatin, see Table 11: Established and Other Potentially Significant Drug Interactions: Alterations in Dose or Regimen May Be Recommended Based on Drug Interaction Studies or Predicted Interaction.
PDE-5 inhibitor	sildenafil for treatment of pulmonary arterial hypertension	A safe and effective dose for the treatment of pulmonary arterial hypertension has not been established with PREZISTA/ritonavir. There is an increased potential for sildenafil-associated adverse events (which include visual disturbances, hypotension, prolonged erection, and syncope).

## 5 WARNINGS AND PRECAUTIONS

### 5.1 Importance of Co-administration with Ritonavir

PREZISTA must be co-administered with ritonavir and food to achieve the desired antiviral effect. Failure to administer PREZISTA with ritonavir and food may result in a loss of efficacy of darunavir.

Please refer to ritonavir prescribing information for additional information on precautionary measures.

### 5.2 Hepatotoxicity

Drug-induced hepatitis (e.g., acute hepatitis, cytolytic hepatitis) has been reported with PREZISTA/ritonavir. During the clinical development program (N=3063), hepatitis was reported in 0.5% of patients receiving combination therapy with PREZISTA/ritonavir. Patients with pre-existing liver dysfunction, including chronic active hepatitis B or C, have an increased risk for liver function abnormalities including severe hepatic adverse events.



Post-marketing cases of liver injury, including some fatalities, have been reported. These have generally occurred in patients with advanced HIV-1 disease taking multiple concomitant medications, having co-morbidities including hepatitis B or C co-infection, and/or developing immune reconstitution syndrome. A causal relationship with PREZISTA/ritonavir therapy has not been established.

Appropriate laboratory testing should be conducted prior to initiating therapy with PREZISTA/ritonavir and patients should be monitored during treatment. Increased AST/ALT monitoring should be considered in patients with underlying chronic hepatitis, cirrhosis, or in patients who have pre-treatment elevations of transaminases, especially during the first several months of PREZISTA/ritonavir treatment.

Evidence of new or worsening liver dysfunction (including clinically significant elevation of liver enzymes and/or symptoms such as fatigue, anorexia, nausea, jaundice, dark urine, liver tenderness, hepatomegaly) in patients on PREZISTA/ritonavir should prompt consideration of interruption or discontinuation of treatment.

### **5.3 Severe Skin Reactions**

During the clinical development program (n=3063), severe skin reactions, accompanied by fever and/or elevations of transaminases in some cases, have been reported in 0.4% of subjects. Stevens-Johnson Syndrome was rarely (less than 0.1%) reported during the clinical development program. During post-marketing experience toxic epidermal necrolysis, drug rash with eosinophilia and systemic symptoms, and acute generalized exanthematous pustulosis have been reported. Discontinue PREZISTA/ritonavir immediately if signs or symptoms of severe skin reactions develop. These can include but are not limited to severe rash or rash accompanied with fever, general malaise, fatigue, muscle or joint aches, blisters, oral lesions, conjunctivitis, hepatitis and/or eosinophilia.

Rash (all grades, regardless of causality) occurred in 10.3% of subjects treated with PREZISTA/ritonavir [also *see Adverse Reactions (6)*]. Rash was mostly mild-to-moderate, often occurring within the first four weeks of treatment and resolving with continued dosing. The discontinuation rate due to rash in subjects using PREZISTA/ritonavir was 0.5%.

Rash occurred more commonly in treatment-experienced subjects receiving regimens containing PREZISTA/ritonavir + raltegravir compared to subjects receiving PREZISTA/ritonavir without raltegravir or raltegravir without PREZISTA/ritonavir. However, rash that was considered drug related occurred at similar rates for all three groups. These rashes were mild to moderate in severity and did not limit therapy; there were no discontinuations due to rash.

### **5.4 Sulfa Allergy**

Darunavir contains a sulfonamide moiety. PREZISTA should be used with caution in patients with a known sulfonamide allergy. In clinical studies with PREZISTA/ritonavir, the incidence and severity of rash were similar in subjects with or without a history of sulfonamide allergy.

## 5.5 Risk of Serious Adverse Reactions due to Drug Interactions

Initiation of PREZISTA/ritonavir, a CYP3A inhibitor, in patients receiving medications metabolized by CYP3A or initiation of medications metabolized by CYP3A in patients already receiving PREZISTA/ritonavir, may increase plasma concentrations of medications metabolized by CYP3A. Initiation of medications that inhibit or induce CYP3A may increase or decrease concentrations of PREZISTA/ritonavir, respectively. These interactions may lead to:

- Clinically significant adverse reactions, potentially leading to severe, life threatening, or fatal events from greater exposures of concomitant medications.
- Clinically significant adverse reactions from greater exposures of PREZISTA/ritonavir.
- Loss of therapeutic effect of PREZISTA/ritonavir and possible development of resistance.

See Table 11 for steps to prevent or manage these possible and known significant drug interactions, including dosing recommendations [*see Drug Interactions (7)*]. Consider the potential for drug interactions prior to and during PREZISTA/ritonavir therapy; review concomitant medications during PREZISTA/ritonavir therapy; and monitor for the adverse reactions associated with the concomitant drugs [*see Contraindications (4) and Drug Interactions (7)*].

## 5.6 Diabetes Mellitus/Hyperglycemia

New onset diabetes mellitus, exacerbation of pre-existing diabetes mellitus, and hyperglycemia have been reported during postmarketing surveillance in HIV-infected patients receiving protease inhibitor (PI) therapy. Some patients required either initiation or dose adjustments of insulin or oral hypoglycemic agents for treatment of these events. In some cases, diabetic ketoacidosis has occurred. In those patients who discontinued PI therapy, hyperglycemia persisted in some cases. Because these events have been reported voluntarily during clinical practice, estimates of frequency cannot be made and causal relationships between PI therapy and these events have not been established.

## 5.7 Fat Redistribution

Redistribution/accumulation of body fat, including central obesity, dorsocervical fat enlargement (buffalo hump), peripheral wasting, facial wasting, breast enlargement, and “cushingoid appearance” have been observed in patients receiving antiretroviral therapy. The mechanism and long-term consequences of these events are currently unknown. A causal relationship has not been established.

## 5.8 Immune Reconstitution Syndrome

Immune reconstitution syndrome has been reported in patients treated with combination antiretroviral therapy, including PREZISTA. During the initial phase of combination antiretroviral treatment, patients whose immune systems respond may develop an inflammatory response to indolent or residual opportunistic infections (such as *Mycobacterium avium*

infection, cytomegalovirus, *Pneumocystis jirovecii* pneumonia [PCP], or tuberculosis), which may necessitate further evaluation and treatment.

Autoimmune disorders (such as Graves' disease, polymyositis, and Guillain-Barré syndrome) have also been reported to occur in the setting of immune reconstitution; however, the time to onset is more variable, and can occur many months after initiation of antiretroviral treatment.

## 5.9 Hemophilia

There have been reports of increased bleeding, including spontaneous skin hematomas and hemarthrosis in patients with hemophilia type A and B treated with PIs. In some patients, additional factor VIII was given. In more than half of the reported cases, treatment with PIs was continued or reintroduced if treatment had been discontinued. A causal relationship between PI therapy and these episodes has not been established.

## 5.10 Not Recommended in Pediatric Patients Below 3 Years of Age

PREZISTA/ritonavir in pediatric patients below 3 years of age is not recommended in view of toxicity and mortality observed in juvenile rats dosed with darunavir (from 20 mg/kg to 1000 mg/kg) up to days 23 to 26 of age [see *Use in Specific Populations* (8.1 and 8.4) and *Clinical Pharmacology* (12.3)].

## 6 ADVERSE REACTIONS

The following adverse reactions are discussed in other sections of labeling:

- Hepatotoxicity [see *Warnings and Precautions* (5.2)]
- Severe Skin Reactions [see *Warnings and Precautions* (5.3)]
- Diabetes Mellitus/Hyperglycemia [see *Warnings and Precautions* (5.6)]
- Fat Redistribution [see *Warnings and Precautions* (5.7)]
- Immune Reconstitution Syndrome [see *Warnings and Precautions* (5.8)]
- Hemophilia [see *Warnings and Precautions* (5.9)]

Due to the need for co-administration of PREZISTA with ritonavir, please refer to ritonavir prescribing information for ritonavir-associated adverse reactions.

### 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

#### Treatment Naïve-Adults: TMC114-C211

The safety assessment is based on all safety data from the Phase 3 trial TMC114-C211 comparing PREZISTA/ritonavir 800/100 mg once daily versus lopinavir/ritonavir 800/200 mg

per day in 689 antiretroviral treatment-naïve HIV-1-infected adult subjects. The total mean exposure for subjects in the PREZISTA/ritonavir 800/100 mg once daily arm and in the lopinavir/ritonavir 800/200 mg per day arm was 162.5 and 153.5 weeks, respectively.

The majority of the adverse drug reactions (ADRs) reported during treatment with PREZISTA/ritonavir 800/100 mg once daily were mild in severity. The most common clinical ADRs to PREZISTA/ritonavir 800/100 mg once daily (greater than or equal to 5%) of at least moderate intensity (greater than or equal to Grade 2) were diarrhea, headache, abdominal pain and rash. 2.3% of subjects in the PREZISTA/ritonavir arm discontinued treatment due to ADRs.

ADRs to PREZISTA/ritonavir 800/100 mg once daily of at least moderate intensity (greater than or equal to Grade 2) in antiretroviral treatment-naïve HIV-1-infected adult subjects are presented in Table 7 and subsequent text below the table.

**Table 7: Selected Clinical Adverse Drug Reactions to PREZISTA/ritonavir 800/100 mg Once Daily\* of at Least Moderate Intensity (≥Grade 2) Occurring in ≥2% of Antiretroviral Treatment-Naïve HIV-1-Infected Adult Subjects (Trial TMC114-C211)**

System organ class, preferred term, %	PREZISTA/ritonavir 800/100 mg once daily + TDF/FTC N=343	lopinavir/ritonavir 800/200 mg per day + TDF/FTC N=346
<b>Gastrointestinal Disorders</b>		
Abdominal pain	6%	6%
Diarrhea	9%	16%
Nausea	4%	4%
Vomiting	2%	4%
<b>General Disorders and Administration Site Conditions</b>		
Fatigue	<1%	3%
<b>Metabolism and Nutrition Disorders</b>		
Anorexia	2%	<1%
<b>Nervous System Disorders</b>		
Headache	7%	6%
<b>Skin and Subcutaneous Tissue Disorders</b>		
Rash	6%	7%

N=total number of subjects per treatment group; FTC=emtricitabine; TDF=tenofovir disoproxil fumarate

\* Excluding laboratory abnormalities reported as ADRs.

### **Less Common Adverse Reactions**

Treatment-emergent ADRs of at least moderate intensity (greater than or equal to Grade 2) occurring in less than 2% of antiretroviral treatment-naïve subjects receiving PREZISTA/ritonavir 800/100 mg once daily are listed below by body system:

*Gastrointestinal Disorders:* acute pancreatitis, dyspepsia, flatulence

*General Disorders and Administration Site Conditions:* asthenia

*Hepatobiliary Disorders:* acute hepatitis (e.g., acute hepatitis, cytolytic hepatitis, hepatotoxicity)

*Immune System Disorders:* (drug) hypersensitivity, immune reconstitution syndrome

*Metabolism and Nutrition Disorders:* diabetes mellitus

*Musculoskeletal and Connective Tissue Disorders:* myalgia, osteonecrosis

*Psychiatric Disorders:* abnormal dreams

*Skin and Subcutaneous Tissue Disorders:* angioedema, pruritus, Stevens-Johnson Syndrome, urticaria

### **Laboratory Abnormalities**

Selected Grade 2 to 4 laboratory abnormalities that represent a worsening from baseline observed in antiretroviral treatment-naïve adult subjects treated with PREZISTA/ritonavir 800/100 mg once daily are presented in Table 8.

**Table 8: Grade 2 to 4 Laboratory Abnormalities Observed in Antiretroviral Treatment-Naïve HIV-1-Infected Adult Subjects\* (Trial TMC114-C211)**

Laboratory parameter %	Limit	PREZISTA/ritonavir 800/100 mg once daily + TDF/FTC	lopinavir/ritonavir 800/200 mg per day + TDF/FTC
<b>Biochemistry</b>			
Alanine Aminotransferase			
Grade 2	>2.5 to ≤5.0 X ULN	9%	9%
Grade 3	>5.0 to ≤10.0 X ULN	3%	3%
Grade 4	>10.0 X ULN	<1%	3%
Aspartate Aminotransferase			
Grade 2	>2.5 to ≤5.0 X ULN	7%	10%
Grade 3	>5.0 to ≤10.0 X ULN	4%	2%
Grade 4	>10.0 X ULN	1%	3%
Alkaline Phosphatase			
Grade 2	>2.5 to ≤5.0 X ULN	1%	1%
Grade 3	>5.0 to ≤10.0 X ULN	0%	<1%
Grade 4	>10.0 X ULN	0%	0%
Hyperbilirubinemia			
Grade 2	>1.5 to ≤2.5 X ULN	<1%	5%
Grade 3	>2.5 to ≤5.0 X ULN	<1%	<1%
Grade 4	>5.0 X ULN	0%	0%
Triglycerides			
Grade 2	5.65-8.48 mmol/L 500-750 mg/dL	3%	10%
Grade 3	8.49-13.56 mmol/L 751-1200 mg/dL	2%	5%
Grade 4	>13.56 mmol/L >1200 mg/dL	1%	1%
Total Cholesterol			
Grade 2	6.20-7.77 mmol/L 240-300 mg/dL	23%	27%
Grade 3	>7.77 mmol/L >300 mg/dL	1%	5%
Low-Density Lipoprotein Cholesterol			
Grade 2	4.13-4.90 mmol/L 160-190 mg/dL	14%	12%

Grade 3	≥4.91 mmol/L ≥191 mg/dL	9%	6%
Elevated Glucose Levels			
Grade 2	6.95-13.88 mmol/L 126-250 mg/dL	11%	10%
Grade 3	13.89-27.75 mmol/L 251-500 mg/dL	1%	<1%
Grade 4	>27.75 mmol/L >500 mg/dL	0%	0%
Pancreatic Lipase			
Grade 2	>1.5 to ≤3.0 X ULN	3%	2%
Grade 3	>3.0 to ≤5.0 X ULN	<1%	1%
Grade 4	>5.0 X ULN	0%	<1%
Pancreatic Amylase			
Grade 2	>1.5 to ≤2.0 X ULN	5%	2%
Grade 3	>2.0 to ≤5.0 X ULN	5%	4%
Grade 4	>5.0 X ULN	0%	<1%

N=total number of subjects per treatment group; FTC=emtricitabine; TDF=tenofovir disoproxil fumarate

\* Grade 4 data not applicable in Division of AIDS grading scale.

### Treatment-Experienced Adults: TMC114-C214

The safety assessment is based on all safety data from the Phase 3 trial TMC114-C214 comparing PREZISTA/ritonavir 600/100 mg twice daily versus lopinavir/ritonavir 400/100 mg twice daily in 595 antiretroviral treatment-experienced HIV-1-infected adult subjects. The total mean exposure for subjects in the PREZISTA/ritonavir 600/100 mg twice daily arm and in the lopinavir/ritonavir 400/100 mg twice daily arm was 80.7 and 76.4 weeks, respectively.

The majority of the ADRs reported during treatment with PREZISTA/ritonavir 600/100 mg twice daily were mild in severity. The most common clinical ADRs to PREZISTA/ritonavir 600/100 mg twice daily (greater than or equal to 5%) of at least moderate intensity (greater than or equal to Grade 2) were diarrhea, nausea, rash, abdominal pain and vomiting. 4.7% of subjects in the PREZISTA/ritonavir arm discontinued treatment due to ADRs.

ADRs to PREZISTA/ritonavir 600/100 mg twice daily of at least moderate intensity (greater than or equal to Grade 2) in antiretroviral treatment-experienced HIV-1-infected adult subjects are presented in Table 9 and subsequent text below the table.

**Table 9: Selected Clinical Adverse Drug Reactions to PREZISTA/ritonavir 600/100 mg Twice Daily\* of at Least Moderate Intensity (≥Grade 2) Occurring in ≥2% of Antiretroviral Treatment-Experienced HIV-1-Infected Adult Subjects (Trial TMC114-C214)**

System organ class, preferred term, %	PREZISTA/ritonavir 600/100 mg twice daily + OBR N=298	lopinavir/ritonavir 400/100 mg twice daily + OBR N=297
<b>Gastrointestinal Disorders</b>		
Abdominal distension	2%	<1%
Abdominal pain	6%	3%
Diarrhea	14%	20%
Dyspepsia	2%	1%
Nausea	7%	6%

Vomiting	5%	3%
<b>General Disorders and Administration Site Conditions</b>		
Asthenia	3%	1%
Fatigue	2%	1%
<b>Metabolism and Nutrition Disorders</b>		
Anorexia	2%	2%
Diabetes mellitus	2%	<1%
<b>Nervous System Disorders</b>		
Headache	3%	3%
<b>Skin and Subcutaneous Tissue Disorders</b>		
Rash	7%	3%

N=total number of subjects per treatment group; OBR=optimized background regimen

\* Excluding laboratory abnormalities reported as ADRs

### *Less Common Adverse Reactions*

Treatment-emergent ADRs of at least moderate intensity (greater than or equal to Grade 2) occurring in less than 2% of antiretroviral treatment-experienced subjects receiving PREZISTA/ritonavir 600/100 mg twice daily are listed below by body system:

*Gastrointestinal Disorders:* acute pancreatitis, flatulence

*Musculoskeletal and Connective Tissue Disorders:* myalgia

*Psychiatric Disorders:* abnormal dreams

*Skin and Subcutaneous Tissue Disorders:* pruritus, urticaria

### *Laboratory Abnormalities*

Selected Grade 2 to 4 laboratory abnormalities that represent a worsening from baseline observed in antiretroviral treatment-experienced adult subjects treated with PREZISTA/ritonavir 600/100 mg twice daily are presented in Table 10.

**Table 10: Grade 2 to 4 Laboratory Abnormalities Observed in Antiretroviral Treatment-Experienced HIV-1-Infected Adult Subjects\* (Trial TMC114-C214)**

Laboratory parameter, %	Limit	PREZISTA/ritonavir 600/100 mg twice daily + OBR	lopinavir/ritonavir 400/100 mg twice daily + OBR
<b>Biochemistry</b>			
Alanine Aminotransferase			
Grade 2	>2.5 to ≤5.0 X ULN	7%	5%
Grade 3	>5.0 to ≤10.0 X ULN	2%	2%
Grade 4	>10.0 X ULN	1%	2%
Aspartate Aminotransferase			
Grade 2	>2.5 to ≤5.0 X ULN	6%	6%
Grade 3	>5.0 to ≤10.0 X ULN	2%	2%
Grade 4	>10.0 X ULN	<1%	2%
Alkaline Phosphatase			
Grade 2	>2.5 to ≤5.0 X ULN	<1%	0%
Grade 3	>5.0 to ≤10.0 X ULN	<1%	<1%
Grade 4	>10.0 X ULN	0%	0%

Hyperbilirubinemia			
Grade 2	>1.5 to ≤2.5 X ULN	<1%	2%
Grade 3	>2.5 to ≤5.0 X ULN	<1%	<1%
Grade 4	>5.0 X ULN	<1%	0%
Triglycerides			
Grade 2	5.65-8.48 mmol/L 500-750 mg/dL	10%	11%
Grade 3	8.49-13.56 mmol/L 751-1200 mg/dL	7%	10%
Grade 4	>13.56 mmol/L >1200 mg/dL	3%	6%
Total Cholesterol			
Grade 2	6.20-7.77 mmol/L 240-300 mg/dL	25%	23%
Grade 3	>7.77 mmol/L >300 mg/dL	10%	14%
Low-Density Lipoprotein Cholesterol			
Grade 2	4.13-4.90 mmol/L 160-190 mg/dL	14%	14%
Grade 3	≥4.91 mmol/L ≥191 mg/dL	8%	9%
Elevated Glucose Levels			
Grade 2	6.95-13.88 mmol/L 126-250 mg/dL	10%	11%
Grade 3	13.89-27.75 mmol/L 251-500 mg/dL	1%	<1%
Grade 4	>27.75 mmol/L >500 mg/dL	<1%	0%
Pancreatic Lipase			
Grade 2	>1.5 to ≤3.0 X ULN	3%	4%
Grade 3	>3.0 to ≤5.0 X ULN	2%	<1%
Grade 4	>5.0 X ULN	<1%	0%
Pancreatic Amylase			
Grade 2	>1.5 to ≤2.0 X ULN	6%	7%
Grade 3	>2.0 to ≤5.0 X ULN	7%	3%
Grade 4	>5.0 X ULN	0%	0%

N=total number of subjects per treatment group; OBR=optimized background regimen

\* Grade 4 data not applicable in Division of AIDS grading scale

## Serious ADRs

The following serious ADRs of at least moderate intensity (greater than or equal to Grade 2) occurred in the Phase 2b and Phase 3 trials with PREZISTA/ritonavir: abdominal pain, acute hepatitis, acute pancreatitis, anorexia, asthenia, diabetes mellitus, diarrhea, fatigue, headache, hepatic enzyme increased, hypercholesterolemia, hyperglycemia, hypertriglyceridemia, immune reconstitution syndrome, low density lipoprotein increased, nausea, pancreatic enzyme increased, rash, Stevens-Johnson Syndrome, and vomiting.

## Patients Co-Infected with Hepatitis B and/or Hepatitis C Virus

In subjects co-infected with hepatitis B or C virus receiving PREZISTA/ritonavir, the incidence of adverse events and clinical chemistry abnormalities was not higher than in subjects receiving PREZISTA/ritonavir who were not co-infected, except for increased hepatic enzymes [see



*Warnings and Precautions (5.2)]*. The pharmacokinetic exposure in co-infected subjects was comparable to that in subjects without co-infection.

### **Clinical Trials Experience: Pediatric Patients**

PREZISTA/ritonavir has been studied in combination with other antiretroviral agents in 3 Phase 2 trials. TMC114-C212, in which 80 antiretroviral treatment-experienced HIV-1-infected pediatric subjects 6 to less than 18 years of age and weighing at least 20 kg were included, TMC114-C228, in which 21 antiretroviral treatment-experienced HIV-1-infected pediatric subjects 3 to less than 6 years of age and weighing at least 10 kg were included, and TMC114-C230 in which 12 antiretroviral treatment-naïve HIV-1 infected pediatric patients aged from 12 to less than 18 years and weighing at least 40 kg were included. The TMC114-C212 and C228 trials evaluated PREZISTA/ritonavir twice daily dosing and the TMC114-C230 trial evaluated PREZISTA/ritonavir once daily dosing [*see Use in Specific Populations (8.4) and Clinical Studies (14.4)]*.

Frequency, type, and severity of ADRs in pediatric subjects were comparable to those observed in adults.

#### ***TMC114-C212***

Clinical ADRs to PREZISTA/ritonavir (all grades, greater than or equal to 3%), were vomiting (13%), diarrhea (11%), abdominal pain (10%), headache (9%), rash (5%), nausea (4%), and fatigue (3%).

Grade 3 or 4 laboratory abnormalities were ALT increased (Grade 3: 3%; Grade 4: 1%), AST increased (Grade 3: 1%), pancreatic amylase increased (Grade 3: 4%, Grade 4: 1%), pancreatic lipase increased (Grade 3: 1%), total cholesterol increased (Grade 3: 1%), and LDL increased (Grade 3: 3%).

#### ***TMC114-C228***

Clinical ADRs to PREZISTA/ritonavir (all grades, greater than or equal to 5%), were diarrhea (24%), vomiting (19%), rash (19%), abdominal pain (5%), and anorexia (5%).

There were no Grade 3 or 4 laboratory abnormalities considered as ADRs in this trial.

#### ***TMC114-C230***

Clinical ADRs to PREZISTA/ritonavir (all grades, greater than or equal to 3%), were vomiting (33%), nausea (25%), diarrhea (16.7%), abdominal pain (8.3%), decreased appetite (8.3%), pruritus (8.3%), and rash (8.3%).

There were no Grade 3 or 4 laboratory abnormalities considered as ADRs in this trial.

## 6.2 Postmarketing Experience

The following events have been identified during post approval use of PREZISTA. Because these events are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Redistribution of body fat has been reported.

Rarely, rhabdomyolysis (associated with co-administration with HMG-CoA reductase inhibitors and PREZISTA/ritonavir) has been reported.

In addition, toxic epidermal necrolysis, acute generalized exanthematous pustulosis and drug rash with eosinophilia and systemic symptoms have been reported rarely [*see Warnings and Precautions (5.3)*].

## 7 DRUG INTERACTIONS

### 7.1 Potential for PREZISTA/ritonavir to Affect Other Drugs

PREZISTA co-administered with ritonavir is an inhibitor of CYP3A, CYP2D6, and P-gp. Co-administration of PREZISTA and ritonavir with drugs that are primarily metabolized by CYP3A and CYP2D6, or are transported by P-gp may result in increased plasma concentrations of such drugs, which could increase or prolong their therapeutic effect and adverse events (see Table 11).

### 7.2 Potential for Other Drugs to Affect Darunavir

Darunavir and ritonavir are metabolized by CYP3A. *In vitro* data indicate that darunavir may be a P-gp substrate. Drugs that induce CYP3A activity would be expected to increase the clearance of darunavir and ritonavir, resulting in lowered plasma concentrations of darunavir and ritonavir. Co-administration of darunavir and ritonavir and other drugs that inhibit CYP3A, or P-gp may decrease the clearance of darunavir and ritonavir and may result in increased plasma concentrations of darunavir and ritonavir (see Table 11).

### 7.3 Established and Other Potentially Significant Drug Interactions

Table 11 provides dosing recommendations as a result of drug interactions with PREZISTA/ritonavir. These recommendations are based on either drug interaction studies or predicted interactions due to the expected magnitude of interaction and potential for serious adverse events or loss of efficacy [*see Contraindications (4) and Clinical Pharmacology (12.3)*].

**Table 11: Established and Other Potentially Significant Drug Interactions: Alterations in Dose or Regimen May be Recommended Based on Drug Interaction Studies or Predicted Interaction**

(see *Contraindications (4)* for a complete list of contraindicated drugs)  
[see *Clinical Pharmacology (12.3)* for Magnitude of Interaction, Tables 16 and 17]

Concomitant Drug Class Drug Name	Effect on Concentration of Darunavir Or Concomitant Drug	Clinical Comment
<b>HIV-1-Antiviral Agents: Nucleoside Reverse Transcriptase Inhibitors (NRTIs)</b>		
didanosine	↔ darunavir ↔ didanosine	Didanosine should be administered one hour before or two hours after PREZISTA/ritonavir (which are administered with food).
<b>HIV-1-Antiviral Agents: HIV-Protease Inhibitors (PIs)</b>		
indinavir  (The reference regimen for indinavir was indinavir/ritonavir 800/100 mg twice daily.)	↑ darunavir ↑ indinavir	The appropriate dose of indinavir in combination with PREZISTA/ritonavir has not been established.
lopinavir/ritonavir	↓ darunavir ↔ lopinavir	Appropriate doses of the combination have not been established. Hence, it is not recommended to co-administer lopinavir/ritonavir and PREZISTA, with or without ritonavir.
saquinavir	↓ darunavir ↔ saquinavir	Appropriate doses of the combination have not been established. Hence, it is not recommended to co-administer saquinavir and PREZISTA, with or without ritonavir.
Other HIV protease inhibitors, except atazanavir [see <i>Drug Interactions(7.4)</i> ]		As coadministration with PREZISTA/ritonavir has not been studied, coadministration is not recommended.
<b>HIV-1-Antiviral Agents: CCR5 co-receptor antagonists</b>		
maraviroc	↑ maraviroc	When used in combination with PREZISTA/ritonavir, the dose of maraviroc should be 150 mg twice daily.
<b>Other Agents</b>		
<b>Antiarrhythmics:</b>  e.g. amiodarone, bepridil, disopyramide, flecainide, lidocaine (systemic), mexiletine, propafenone, quinidine  digoxin	↑ antiarrhythmics  ↑ digoxin	<b>For contraindicated antiarrhythmics, [see <i>Contraindications (4)</i>].</b>  Therapeutic concentration monitoring, if available, is recommended for antiarrhythmics when co-administered with PREZISTA/ritonavir.  The lowest dose of digoxin should initially be prescribed. The serum digoxin concentrations should be monitored and used for titration of digoxin dose to obtain the desired clinical effect.
<b>Antibacterial:</b> clarithromycin	↔ darunavir ↑ clarithromycin	No dose adjustment of the combination is required for patients with normal renal function. For co-administration of clarithromycin and

		<p>PREZISTA/ritonavir in patients with renal impairment, the following dose adjustments should be considered:</p> <ul style="list-style-type: none"> <li>For subjects with CLcr of 30-60 mL/min, the dose of clarithromycin should be reduced by 50%.</li> <li>For subjects with CLcr of &lt;30 mL/min, the dose of clarithromycin should be reduced by 75%.</li> </ul>
<p><b>Anticoagulants:</b> apixaban</p> <p>dabigatran etexilate</p> <p>rivaroxaban</p> <p>warfarin</p>	<p>↑ anticoagulant</p> <p>↓ warfarin ↔ darunavir</p>	<p>Concomitant use of apixaban and PREZISTA/ritonavir is not recommended.</p> <p>The combination of PREZISTA/ritonavir and dabigatran etexilate is not recommended in specific renal impairment groups, depending on the indication. Please see the dabigatran US prescribing information for specific recommendations.</p> <p>Co-administration of PREZISTA/ritonavir and rivaroxaban is not recommended.</p> <p>Warfarin concentrations are decreased when co-administered with PREZISTA/ritonavir. It is recommended that the international normalized ratio (INR) be monitored when warfarin is combined with PREZISTA/ritonavir.</p>
<p><b>Anticonvulsants:</b> carbamazepine</p> <p>phenobarbital, phenytoin</p>	<p>↔ darunavir ↑ carbamazepine</p> <p>↔ darunavir ↓ phenytoin ↓ phenobarbital</p>	<p>The dose of either PREZISTA/ritonavir or carbamazepine does not need to be adjusted when initiating co-administration with PREZISTA/ritonavir and carbamazepine. Clinical monitoring of carbamazepine concentrations and its dose titration is recommended to achieve the desired clinical response.</p> <p>Phenytoin and phenobarbital levels should be monitored when co-administering with PREZISTA/ritonavir.</p>
<p><b>Antidepressants:</b> <u>Selective Serotonin Reuptake Inhibitors (SSRIs):</u> paroxetine, sertraline</p> <p><u>Tricyclic Antidepressants (TCAs):</u> amitriptyline, desipramine, imipramine, nortriptyline</p> <p><u>Other:</u> trazodone</p>	<p>↓ paroxetine ↓ sertraline</p> <p>↑ amitriptyline ↑ desipramine ↑ imipramine ↑ nortriptyline</p> <p>↑ trazodone</p>	<p>If either sertraline or paroxetine is initiated in patients receiving PREZISTA/ritonavir, dose titrating the SSRI based on a clinical assessment of antidepressant response is recommended. Monitor for antidepressant response in patients on a stable dose of sertraline or paroxetine who start treatment with PREZISTA/ritonavir.</p> <p>Use a lower dose of the tricyclic antidepressants and trazodone due to potential increased adverse events such as nausea, dizziness, hypotension and syncope.</p>
<p><b>Antifungals:</b> itraconazole, ketoconazole, posaconazole</p>	<p>↑ darunavir ↑ itraconazole (not studied) ↑ ketoconazole</p>	<p>Monitor for increased PREZISTA/ritonavir adverse events with concomitant use of itraconazole, ketoconazole, or posaconazole. When co-administration is required, the daily dose of</p>

voriconazole	<p>↔ posaconazole (not studied)</p> <p>↓ voriconazole (not studied)</p>	<p>ketoconazole or itraconazole should not exceed 200 mg with monitoring for increased antifungal adverse events.</p> <p>Voriconazole is not recommended for patients receiving PREZISTA/ritonavir unless an assessment comparing predicted benefit to risk ratio justifies the use of voriconazole.</p>
<b>Anti-gout:</b> colchicine	↑ colchicine	<p><b>The coadministration of PREZISTA/ritonavir with colchicine in patients with renal or hepatic impairment is contraindicated [see Contraindications (4)].</b></p> <p><u>For patients without renal or hepatic impairment:</u></p> <ul style="list-style-type: none"> <li>• <u>Treatment of gout-flares – co-administration of colchicine in patients on PREZISTA/ritonavir:</u> 0.6 mg (1 tablet) × 1 dose, followed by 0.3 mg (half tablet) 1 hour later. Treatment course to be repeated no earlier than 3 days.</li> <li>• <u>Prophylaxis of gout-flares – co-administration of colchicine in patients on PREZISTA/ritonavir:</u> If the original regimen was 0.6 mg twice a day, the regimen should be adjusted to 0.3 mg once a day. If the original regimen was 0.6 mg once a day, the regimen should be adjusted to 0.3 mg once every other day.</li> <li>• <u>Treatment of familial Mediterranean fever – co-administration of colchicine in patients on PREZISTA/ritonavir:</u> maximum daily dose of 0.6 mg (may be given as 0.3 mg twice a day).</li> </ul>
<b>Antimalarial:</b> artemether/lumefantrine	<p>↓ artemether</p> <p>↓ dihydroartemisinin</p> <p>↑ lumefantrine</p> <p>↔ darunavir</p>	The combination of PREZISTA/ritonavir and artemether/lumefantrine can be used without dose adjustments. However, the combination should be used with caution as increased lumefantrine exposure may increase the risk of QT prolongation.
<b>Antimycobacterials:</b>  rifabutin  (The reference regimen for rifabutin was 300 mg once daily.)  rifapentine	<p>↑ darunavir</p> <p>↑ rifabutin</p> <p>↑ 25-O-desacetyl rifabutin</p> <p>↓ darunavir</p>	<p><b>For contraindicated antimycobacterials, [see Contraindications (4)].</b></p> <p>Dose reduction of rifabutin by at least 75% of the usual dose (300 mg once daily) is recommended (i.e., a maximum dose of 150 mg every other day). Increased monitoring for adverse events is warranted in patients receiving this combination and further dose reduction of rifabutin may be necessary.</p> <p>Co-administration of PREZISTA/ritonavir with rifapentine is not recommended.</p>
<b>Antineoplastics:</b> dasatinib, nilotinib	↑ antineoplastics	A decrease in the dosage or an adjustment of the dosing interval of dasatinib and nilotinib may be necessary for patients. Please refer to the dasatinib and nilotinib prescribing information for dosing instructions.



bosentan		<p><b>PREZISTA/ritonavir:</b> In patients who have been receiving PREZISTA/ritonavir for at least 10 days, start bosentan at 62.5 mg once daily or every other day based upon individual tolerability.</p> <p><u>Co-administration of PREZISTA/ritonavir in patients on bosentan:</u> Discontinue use of bosentan at least 36 hours prior to initiation of PREZISTA/ritonavir. After at least 10 days following the initiation of PREZISTA/ritonavir, resume bosentan at 62.5 mg once daily or every other day based upon individual tolerability.</p>
<p><b>Hepatitis C virus (HCV):</b> <u>Direct-Acting Antivirals:</u></p> <p>simeprevir</p>	<p>↑ simeprevir ↑ darunavir</p>	<p><b>For contraindicated HCV Direct-Acting Antivirals, [see Contraindications (4)].</b></p> <p>Co-administration of PREZISTA/ritonavir and simeprevir is not recommended</p>
<p><b>HMG-CoA reductase inhibitors:</b></p> <p>atorvastatin, pravastatin, rosuvastatin</p>	<p>↑ HMG-CoA reductase inhibitors</p>	<p><b>For contraindicated HMG-CoA reductase inhibitors, [see Contraindications (4)].</b></p> <p>Co-administration of PREZISTA/ritonavir with HMG-Co A reductase inhibitors may lead to adverse events such as myopathy. Titrate atorvastatin, pravastatin or rosuvastatin dose carefully and use the lowest necessary dose while monitoring for adverse events. Do not exceed atorvastatin 20 mg/day.</p>
<p><b>Immunosuppressants:</b> e.g. cyclosporine, tacrolimus, sirolimus</p> <p><b>Immunosuppressant/neoplastic:</b> everolimus</p>	<p>↑ immunosuppressants</p>	<p>Therapeutic concentration monitoring of the immunosuppressive agent is recommended when co-administered with PREZISTA/ritonavir.</p> <p>Co-administration of everolimus and PREZISTA/ritonavir is not recommended.</p>
<p><b>Inhaled beta agonist:</b> salmeterol</p>	<p>↑ salmeterol</p>	<p>Co-administration of salmeterol and PREZISTA/ritonavir is not recommended. The combination may result in increased risk of cardiovascular adverse events associated with salmeterol, including QT prolongation, palpitations and sinus tachycardia.</p>
<p><b>Narcotic analgesics/treatment of opioid dependence:</b> buprenorphine, buprenorphine/naloxone</p> <p>Methadone</p>	<p>↔ buprenorphine, naloxone ↑ norbuprenorphine (metabolite)</p> <p>↓ methadone</p>	<p>No dose adjustment for buprenorphine or buprenorphine/naloxone is required with concurrent administration of PREZISTA/ritonavir. Clinical monitoring is recommended if PREZISTA/ritonavir and buprenorphine or buprenorphine/naloxone are coadministered.</p> <p>No adjustment of methadone dosage is required when initiating co-administration of PREZISTA/ritonavir. However, clinical monitoring is recommended as the dose of methadone during maintenance therapy may need to be adjusted in some patients.</p>
<p><b>Oral contraceptives/estrogen:</b> ethinyl estradiol, norethindrone</p>	<p>↓ ethinyl estradiol</p>	<p>Effective alternative contraceptive method or barrier method of contraception is recommended [see Use in</p>

	↓ norethindrone	<i>Specific Populations (8.3)].</i>
<b>PDE-5 inhibitors:</b> e.g. avanafil, sildenafil, tadalafil, vardenafil	↑ PDE-5 inhibitors (only the use of sildenafil at doses used for treatment of erectile dysfunction has been studied with PREZISTA/ritonavir)	<p>Co-administration with PREZISTA/ritonavir may result in an increase in PDE-5 inhibitor-associated adverse events, including hypotension, syncope, visual disturbances and priapism.</p> <p><u>Use of PDE-5 inhibitors for pulmonary arterial hypertension (PAH):</u>  <b>Co-administration of PREZISTA/ritonavir and sildenafil for treatment of pulmonary arterial hypertension is contraindicated [see Contraindications (4)].</b>  The following dose adjustments are recommended for use of tadalafil with PREZISTA/ritonavir:</p> <ul style="list-style-type: none"> <li>• <u>Co-administration of tadalafil in patients on PREZISTA/ritonavir:</u>  In patients receiving PREZISTA/ritonavir for at least one week, start tadalafil at 20 mg once daily. Increase to 40 mg once daily based upon individual tolerability.</li> <li>• <u>Co-administration of PREZISTA/ritonavir in patients on tadalafil:</u>  Avoid use of tadalafil during the initiation of PREZISTA/ritonavir. Stop tadalafil at least 24 hours prior to starting PREZISTA/ritonavir. After at least one week following the initiation of PREZISTA/ritonavir, resume tadalafil at 20 mg once daily. Increase to 40 mg once daily based upon individual tolerability.</li> </ul> <p><u>Use of PDE-5 inhibitors for erectile dysfunction:</u>  Sildenafil at a single dose not exceeding 25 mg in 48 hours, vardenafil at a single dose not exceeding 2.5 mg dose in 72 hours, or tadalafil at a single dose not exceeding 10 mg dose in 72 hours can be used with increased monitoring for PDE-5 inhibitor-associated adverse events.</p> <p>Co-administration of PREZISTA/ritonavir and avanafil is not recommended.</p>
<b>Proton pump inhibitor:</b> omeprazole	↓ omeprazole ↔ darunavir	<p>When omeprazole is co-administered with PREZISTA/ritonavir, monitor patients for decreased efficacy of omeprazole. Consider increasing the omeprazole dose in patients whose symptoms are not well controlled; avoid use of more than 40 mg per day of omeprazole.</p>
<b>Sedatives/hypnotics:</b>  metabolized by CYP3A e.g. buspirone, diazepam, estazolam, zolpidem	↑ sedatives/hypnotics	<p><b>For contraindicated sedatives/hypnotics, [see Contraindications (4)].</b></p> <p>Titration is recommended when co-administering PREZISTA/ritonavir with sedatives/hypnotics metabolized by CYP3A and a lower dose of the sedatives/hypnotics should be considered with</p>



parenterally administered midazolam	<div data-bbox="883 195 1208 222" data-label="Text"> <p>monitoring for adverse events.</p> </div> <div data-bbox="883 258 1455 468" data-label="Text"> <p>Co-administration of parenteral midazolam should be done in a setting which ensures close clinical monitoring and appropriate medical management in case of respiratory depression and/or prolonged sedation. Dosage reduction for midazolam should be considered, especially if more than a single dose of midazolam is administered.</p> </div>
--	---

## 7.4 Drugs without Clinically Significant Interactions with PREZISTA

No dosage adjustments are recommended when PREZISTA/ritonavir is co-administered with the following medications: atazanavir, dolutegravir, efavirenz, etravirine, nevirapine, nucleoside reverse transcriptase inhibitors (abacavir, emtricitabine, lamivudine, stavudine, tenofovir disoproxil fumarate, zidovudine), pitavastatin, raltegravir, ranitidine, or rilpivirine.

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### Pregnancy Exposure Registry

There is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed to PREZISTA during pregnancy. Healthcare providers are encouraged to register patients by calling the Antiretroviral Pregnancy Registry (APR) 1-800-258-4263.

#### Risk Summary

Available limited data from the APR show no difference in rate of overall birth defects for darunavir (2.7%) compared with the background rate for major birth defects of 2.7% in a U.S. reference population of the Metropolitan Atlanta Congenital Defects Program (MACDP) [*see Data*]. The APR uses the MACDP as the U.S. reference population for birth defects in the general population. The MACDP evaluates women and infants from a limited geographic area and does not include outcomes for births that occurred at less than 20 weeks gestation.

The rate of miscarriage is not reported in the APR. The estimated background rate of miscarriage in clinically recognized pregnancies in the U.S. general population is 15-20%. The background risk of major birth defects and miscarriage for the indicated population is unknown.

Studies in animals did not show evidence of developmental toxicity. Exposures (based on AUC) in rats were 3-fold higher, whereas in mice and rabbits, exposures were lower (less than 1-fold) than human exposures at the recommended daily dose [*see Data*].

#### Clinical Considerations

The recommended dosage in pregnant patients is PREZISTA 600 mg taken with ritonavir 100 mg twice daily with food.

PREZISTA 800 mg taken with ritonavir 100 mg once daily should only be considered in certain pregnant patients who are already on a stable PREZISTA 800 mg with ritonavir 100 mg once daily regimen prior to pregnancy, are virologically suppressed (HIV-1 RNA less than 50 copies per mL), and in whom a change to twice daily PREZISTA 600 mg with ritonavir 100 mg may compromise tolerability or compliance [see *Dosage and Administration* (2.4) and *Clinical Pharmacology* (12.3)].

## **Data**

### ***Human Data***

PREZISTA/ritonavir (600/100 mg twice daily or 800/100 mg once daily) in combination with a background regimen was evaluated in a clinical trial of 34 pregnant women during the second and third trimesters, and postpartum. The pharmacokinetic data demonstrate that exposure to darunavir and ritonavir as part of an antiretroviral regimen was lower during pregnancy compared with postpartum (6-12 weeks). Virologic response was preserved throughout the trial period in both arms. No mother to child transmission occurred in the infants born to the 29 subjects who stayed on the antiretroviral treatment through delivery. PREZISTA/ritonavir was well tolerated during pregnancy and postpartum. There were no new clinically relevant safety findings compared with the known safety profile of PREZISTA/ritonavir in HIV-1-infected adults [see *Clinical Pharmacology* (12.3)].

Based on prospective reports to the APR of 615 live births following exposure to darunavir-containing regimens during pregnancy (including 385 exposed in the first trimester and 230 exposed in the second/third trimester), there was no difference in rate of overall birth defects for darunavir compared with the background rate for major birth defects in a U.S. reference population of the MACDP.

The prevalence of birth defects in live births was 2.6% (95% CI: 1.2% to 4.7%) with first trimester exposure to darunavir containing regimens and 1.7% (95% CI: 0.5% to 4.4%) with second/third trimester exposure to darunavir containing regimens.

### ***Animal Data***

Reproduction studies conducted with darunavir showed no embryotoxicity or teratogenicity in mice (doses up to 1000 mg/kg from gestation day (GD) 6-15 with darunavir alone) and rats (doses up to 1000 mg/kg from GD 7-19 in the presence or absence of ritonavir) as well as in rabbits (doses up to 1000 mg/kg/day from GD 8-20 with darunavir alone). In these studies, darunavir exposures (based on AUC) were higher in rats (3-fold), whereas in mice and rabbits, exposures were lower (less than 1-fold) compared to those obtained in humans at the recommended clinical dose of darunavir boosted with ritonavir.

## **8.2 Lactation**

### **Risk Summary**

The Centers for Disease Control and Prevention recommend that HIV-infected mothers not breastfeed their infants to avoid risking postnatal transmission of HIV.

There are no data on the presence of darunavir in human milk, the effects on the breastfed infant, or the effects on milk production. Darunavir is present in the milk of lactating rats [see *Data*]. Because of the potential for (1) HIV transmission (in HIV-negative infants), (2) developing viral resistance (in HIV-positive infants) and (3) serious adverse reactions in a breastfed infant, instruct mothers not to breastfeed if they are receiving PREZISTA [see *Use in Specific Populations* (8.4)].

## Data

### *Animal Data*

Studies in rats (with darunavir alone or with ritonavir) have demonstrated that darunavir is secreted in the milk. In the rat pre- and postnatal development study, a reduction in pup body weight gain was observed due to exposure of pups to drug substances via milk. The maximal maternal plasma exposures achieved with darunavir (up to 1000 mg/kg with ritonavir) were approximately 50% of those obtained in humans at the recommended clinical dose with ritonavir.

## **8.3 Females and Males of Reproductive Potential**

### Contraception

Use of PREZISTA may reduce the efficacy of combined hormonal contraceptives and the progestin only pill. Advise patients using combined hormonal contraceptives or the progestin only pill to use an effective alternative contraceptive method or add a barrier method of contraception [see *Drug Interactions* (7.3)].

## **8.4 Pediatric Use**

PREZISTA/ritonavir is not recommended in pediatric patients below 3 years of age because of toxicity and mortality observed in juvenile rats dosed with darunavir (from 20 mg/kg to 1000 mg/kg) up to days 23 to 26 of age [see *Warnings and Precautions* (5.10), *Use in Specific Populations* (8.1) and *Clinical Pharmacology* (12.3)].

The safety, pharmacokinetic profile, and virologic and immunologic responses of PREZISTA/ritonavir administered twice daily were evaluated in treatment-experienced HIV-1-infected pediatric subjects 3 to less than 18 years of age and weighing at least 10 kg. These subjects were evaluated in clinical trials TMC114-C212 (80 subjects, 6 to less than 18 years of age) and TMC114-228 (21 subjects, 3 to less than 6 years of age) [see *Adverse Reactions* (6.1), *Clinical Pharmacology* (12.3) and *Clinical Studies* (14.4)]. Frequency, type, and severity of adverse drug reactions in pediatric subjects were comparable to those observed in adults [see *Adverse Reactions* (6.1)]. Refer to *Dosage and Administration* (2.5) for twice-daily dosing recommendations for pediatric subjects 3 to less than 18 years of age and weighing at least 10 kg.

In clinical trial TMC114-C230, the safety, pharmacokinetic profile and virologic and immunologic responses of PREZISTA/ritonavir administered once daily were evaluated in treatment-naïve HIV-1 infected pediatric subjects 12 to less than 18 years of age (12 subjects)

[see *Adverse Reactions* (6.1), *Clinical Pharmacology* (12.3) and *Clinical Studies* (14.4)]. Frequency, type, and severity of adverse drug reactions in pediatric subjects were comparable to those observed in adults [see *Adverse Reactions* (6.1)]. Once daily dosing recommendations for pediatric patients 3 to less than 12 years of age were derived using population pharmacokinetic modeling and simulation. Although a PREZISTA/ritonavir once daily dosing pediatric trial was not conducted in children less than 12 years of age, there is sufficient clinical safety data to support the predicted PREZISTA exposures for the dosing recommendations in this age group [see *Clinical Pharmacology* (12.3)]. Please see *Dosage and Administration* (2.5) for once-daily dosing recommendations for pediatric subjects 3 to less than 18 years of age and weighing at least 10 kg.

### Juvenile Animal Data

In a juvenile toxicity study where rats were directly dosed with darunavir (up to 1000 mg/kg), deaths occurred from post-natal day 5 at plasma exposure levels ranging from 0.1 to 1.0 of the human exposure levels. In a 4-week rat toxicology study, when dosing was initiated on post-natal day 23 (the human equivalent of 2 to 3 years of age), no deaths were observed with a plasma exposure (in combination with ritonavir) 2 times the human plasma exposure levels.

## 8.5 Geriatric Use

Clinical studies of PREZISTA did not include sufficient numbers of patients aged 65 years and over to determine whether they respond differently from younger patients. In general, caution should be exercised in the administration and monitoring of PREZISTA in elderly patients, reflecting the greater frequency of decreased hepatic function, and of concomitant disease or other drug therapy [see *Clinical Pharmacology* (12.3)].

## 8.6 Hepatic Impairment

No dosage adjustment of PREZISTA/ritonavir is necessary for patients with either mild or moderate hepatic impairment. No pharmacokinetic or safety data are available regarding the use of PREZISTA/ritonavir in subjects with severe hepatic impairment. Therefore, PREZISTA/ritonavir is not recommended for use in patients with severe hepatic impairment [see *Dosage and Administration* (2.6) and *Clinical Pharmacology* (12.3)].

## 8.7 Renal Impairment

Population pharmacokinetic analysis showed that the pharmacokinetics of darunavir were not significantly affected in HIV-infected subjects with moderate renal impairment (CrCL between 30-60 mL/min, n=20). No pharmacokinetic data are available in HIV-1-infected patients with severe renal impairment or end stage renal disease; however, because the renal clearance of darunavir is limited, a decrease in total body clearance is not expected in patients with renal impairment. As darunavir and ritonavir are highly bound to plasma proteins, it is unlikely that they will be significantly removed by hemodialysis or peritoneal dialysis [see *Clinical Pharmacology* (12.3)].


## 10 OVERDOSAGE

Human experience of acute overdose with PREZISTA/ritonavir is limited. No specific antidote is available for overdose with PREZISTA. Treatment of overdose with PREZISTA consists of general supportive measures including monitoring of vital signs and observation of the clinical status of the patient. If indicated, elimination of unabsorbed active substance is to be achieved by emesis. Administration of activated charcoal may also be used to aid in removal of unabsorbed active substance. Since PREZISTA is highly protein bound, dialysis is unlikely to be beneficial in significant removal of the active substance.

## 11 DESCRIPTION

PREZISTA (darunavir) is an inhibitor of the human immunodeficiency virus (HIV-1) protease.

PREZISTA (darunavir), in the form of darunavir ethanolate, has the following chemical name: [(1S,2R)-3-[[[4-aminophenyl)sulfonyl](2-methylpropyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-carbamic acid (3R,3aS,6aR)-hexahydrofuro[2,3-b]furan-3-yl ester monoethanolate. Its molecular formula is  $C_{27}H_{37}N_3O_7S \cdot C_2H_5OH$  and its molecular weight is 593.73. Darunavir ethanolate has the following structural formula:



Darunavir ethanolate is a white to off-white powder with a solubility of approximately 0.15 mg per mL in water at 20°C.

PREZISTA 100 mg per mL oral suspension is available as a white to off-white opaque suspension for oral administration.

Each mL of the oral suspension contains darunavir ethanolate equivalent to 100 mg darunavir. In addition, each mL contains the inactive ingredients citric acid monohydrate, hydrochloric acid (for pH adjustment), hydroxypropyl cellulose, masking flavor, methylparaben sodium, microcrystalline cellulose, purified water, sodium carboxymethylcellulose, strawberry cream flavor and sucralose.

PREZISTA 75 mg tablets are available as white, caplet-shaped, film-coated tablets for oral administration. Each 75 mg tablet contains darunavir ethanolate equivalent to 75 mg of darunavir.

PREZISTA 150 mg tablets are available as white, oval-shaped, film-coated tablets for oral administration. Each 150 mg tablet contains darunavir ethanolate equivalent to 150 mg of darunavir.

PREZISTA 600 mg tablets are available as orange, oval-shaped, film-coated tablets for oral administration. Each 600 mg tablet contains darunavir ethanolate equivalent to 600 mg of darunavir.

PREZISTA 800 mg tablets are available as dark red, oval-shaped, film-coated tablets for oral administration. Each 800 mg tablet contains darunavir ethanolate equivalent to 800 mg of darunavir.

During storage, partial conversion from ethanolate to hydrate may occur; however, this does not affect product quality or performance. Each tablet also contains the inactive ingredients colloidal silicon dioxide, crospovidone, magnesium stearate, and microcrystalline cellulose. The 800 mg tablet also contains hypromellose. The 75 and 150 mg tablet film coating, OPADRY® White, contains polyethylene glycol 3350, polyvinyl alcohol-partially hydrolyzed, talc, and titanium dioxide. The 600 mg tablet film coating, OPADRY® Orange, contains FD&C Yellow No. 6, polyethylene glycol 3350, polyvinyl alcohol-partially hydrolyzed, talc, and titanium dioxide. The 800 mg tablet film coating, OPADRY® Dark Red, contains iron oxide red, polyethylene glycol 3350, polyvinyl alcohol-partially hydrolyzed, talc, and titanium dioxide.

All dosages for PREZISTA are expressed in terms of the free form of darunavir.

## **12 CLINICAL PHARMACOLOGY**

### **12.1 Mechanism of Action**

Darunavir is an HIV-1 antiviral drug [*see Microbiology (12.4)*].

### **12.2 Pharmacodynamics**

#### **Cardiac Electrophysiology**

In a thorough QT/QTc study in 40 healthy subjects, PREZISTA/ritonavir doses of 1.33 times the maximum recommended dose did not affect the QT/QTc interval.

### **12.3 Pharmacokinetics**

#### **Pharmacokinetics in Adults**

##### ***General***

Darunavir is primarily metabolized by CYP3A. Ritonavir inhibits CYP3A, thereby increasing the plasma concentrations of darunavir. When a single dose of PREZISTA 600 mg was given orally in combination with 100 mg ritonavir twice daily, there was an approximate 14-fold increase in the systemic exposure of darunavir. Therefore, PREZISTA should only be used in combination with 100 mg of ritonavir to achieve sufficient exposures of darunavir.

The pharmacokinetics of darunavir, co-administered with low dose ritonavir (100 mg), has been evaluated in healthy adult volunteers and in HIV-1-infected subjects. Table 12 displays the population pharmacokinetic estimates of darunavir after oral administration of PREZISTA/ritonavir 600/100 mg twice daily (based on sparse sampling in 285 patients in trial TMC114-C214, 278 patients in trial TMC114-C229 and 119 patients [integrated data] from trials TMC114-C202 and TMC114-C213) and PREZISTA/ritonavir 800/100 mg once daily (based on sparse sampling in 335 patients in trial TMC114-C211 and 280 patients in trial TMC114-C229) to HIV-1-infected patients.

**Table 12: Population Pharmacokinetic Estimates of Darunavir at PREZISTA/ritonavir 800/100 mg Once Daily (Trial TMC114-C211, 48-Week Analysis and Trial TMC114-C229, 48-Week Analysis) and PREZISTA/ritonavir 600/100 mg Twice Daily (Trial TMC114-C214, 48-Week Analysis, Trial TMC114-C229, 48-Week Analysis and Integrated Data from Trials TMC114-C213 and TMC114-C202, Primary 24-Week Analysis)**

Parameter	PREZISTA / ritonavir 800/100 mg once daily		PREZISTA / ritonavir 600/100 mg twice daily		
	TMC114-C211 N=335	TMC114-C229 N=280	TMC114-C214 N=285	TMC114-C229 N=278	TMC114-C213 + TMC114-C202 (integrated data) N=119
<b>AUC<sub>24h</sub> (ng.h/mL)*</b>					
Mean ± Standard Deviation	93026 ± 27050	93334 ± 28626	116796 ± 33594	114302 ± 32681	124698 ± 32286
Median (Range)	87854 (45000-219240)	87788 (45456-236920)	111632 (64874-355360)	109401 (48934-323820)	123336 (67714-212980)
<b>C<sub>0h</sub> (ng/mL)</b>					
Mean ± Standard Deviation	2282 ± 1168	2160 ± 1201	3490 ± 1401	3386 ± 1372	3578 ± 1151
Median (Range)	2041 (368-7242)	1896 (184-7881)	3307 (1517-13198)	3197 (250-11865)	3539 (1255-7368)

N=number of subjects with data

\* AUC<sub>24h</sub> is calculated as AUC<sub>12h</sub>\*2.

### Absorption and Bioavailability

Darunavir, co-administered with 100 mg ritonavir twice daily, was absorbed following oral administration with a T<sub>max</sub> of approximately 2.5-4 hours. The absolute oral bioavailability of a single 600 mg dose of darunavir alone and after co-administration with 100 mg ritonavir twice daily was 37% and 82%, respectively. *In vivo* data suggest that PREZISTA/ritonavir is an inhibitor of the P-glycoprotein (P-gp) transporters.

### Effects of Food on Oral Absorption

When PREZISTA tablets were administered with food, the C<sub>max</sub> and AUC of darunavir, co-administered with ritonavir, is approximately 40% higher relative to the fasting state. Within the range of meals studied, darunavir exposure is similar. The total caloric content of the various meals evaluated ranged from 240 Kcal (12 gms fat) to 928 Kcal (56 gms fat).

### *Distribution*

Darunavir is approximately 95% bound to plasma proteins. Darunavir binds primarily to plasma alpha 1-acid glycoprotein (AAG).

### *Metabolism*

*In vitro* experiments with human liver microsomes (HLMs) indicate that darunavir primarily undergoes oxidative metabolism. Darunavir is extensively metabolized by CYP enzymes, primarily by CYP3A. A mass balance study in healthy volunteers showed that after a single dose administration of 400 mg <sup>14</sup>C-darunavir, co-administered with 100 mg ritonavir, the majority of the radioactivity in the plasma was due to darunavir. At least 3 oxidative metabolites of darunavir have been identified in humans; all showed activity that was at least 90% less than the activity of darunavir against wild-type HIV-1.

### *Elimination*

A mass balance study in healthy volunteers showed that after single dose administration of 400 mg <sup>14</sup>C-darunavir, co-administered with 100 mg ritonavir, approximately 79.5% and 13.9% of the administered dose of <sup>14</sup>C-darunavir was recovered in the feces and urine, respectively. Unchanged darunavir accounted for approximately 41.2% and 7.7% of the administered dose in feces and urine, respectively. The terminal elimination half-life of darunavir was approximately 15 hours when co-administered with ritonavir. After intravenous administration, the clearance of darunavir, administered alone and co-administered with 100 mg twice daily ritonavir, was 32.8 L/h and 5.9 L/h, respectively.

### Special Populations

#### *Hepatic Impairment*

Darunavir is primarily metabolized by the liver. The steady-state pharmacokinetic parameters of darunavir were similar after multiple dose co-administration of PREZISTA/ritonavir 600/100 mg twice daily to subjects with normal hepatic function (n=16), mild hepatic impairment (Child-Pugh Class A, n=8), and moderate hepatic impairment (Child-Pugh Class B, n=8). The effect of severe hepatic impairment on the pharmacokinetics of darunavir has not been evaluated [see *Dosage and Administration (2.6) and Use in Specific Populations (8.6)*].

#### *Hepatitis B or Hepatitis C Virus Co-infection*

The 48-week analysis of the data from Studies TMC114-C211 and TMC114-C214 in HIV-1-infected subjects indicated that hepatitis B and/or hepatitis C virus co-infection status had no apparent effect on the exposure of darunavir.

#### *Renal Impairment*

Results from a mass balance study with <sup>14</sup>C-PREZISTA/ritonavir showed that approximately 7.7% of the administered dose of darunavir is excreted in the urine as unchanged drug. As darunavir and ritonavir are highly bound to plasma proteins, it is unlikely that they will be significantly removed by hemodialysis or peritoneal dialysis. Population pharmacokinetic analysis showed that the pharmacokinetics of darunavir were not significantly affected in HIV-



1-infected subjects with moderate renal impairment (CrCL between 30-60 mL/min, n=20). There are no pharmacokinetic data available in HIV-1-infected patients with severe renal impairment or end stage renal disease [see *Use in Specific Populations* (8.7)].

### *Gender*

Population pharmacokinetic analysis showed higher mean darunavir exposure in HIV-1-infected females compared to males. This difference is not clinically relevant.

### *Race*

Population pharmacokinetic analysis of darunavir in HIV-1-infected subjects indicated that race had no apparent effect on the exposure to darunavir.

### *Geriatric Patients*

Population pharmacokinetic analysis in HIV-1-infected subjects showed that darunavir pharmacokinetics are not considerably different in the age range (18 to 75 years) evaluated in HIV-1-infected subjects (n=12, age greater than or equal to 65) [see *Use in Specific Populations* (8.5)].

### *Pediatric Patients*

#### PREZISTA/ritonavir administered twice daily

The pharmacokinetics of darunavir in combination with ritonavir in 93 antiretroviral treatment-experienced HIV-1-infected pediatric subjects 3 to less than 18 years of age and weighing at least 10 kg showed that the administered weight-based dosages resulted in similar darunavir exposure when compared to the darunavir exposure achieved in treatment-experienced adults receiving PREZISTA/ritonavir 600/100 mg twice daily [see *Dosage and Administration* (2.5)].

#### PREZISTA/ritonavir administered once daily

The pharmacokinetics of darunavir in combination with ritonavir in 12 antiretroviral treatment-naïve HIV-1-infected pediatric subjects 12 to less than 18 years of age and weighing at least 40 kg receiving PREZISTA/ritonavir 800/100 mg once daily resulted in similar darunavir exposures when compared to the darunavir exposure achieved in treatment-naïve adults receiving PREZISTA/ritonavir 800/100 mg once daily [see *Dosage and Administration* (2.5)].

Based on population pharmacokinetic modeling and simulation, the proposed PREZISTA/ritonavir once daily dosing regimens for pediatric patients 3 to less than 12 years of age is predicted to result in similar darunavir exposures when compared to the darunavir exposures achieved in treatment-naïve adults receiving PREZISTA/ritonavir 800/100 mg once daily [see *Dosage and Administration* (2.5)].

The population pharmacokinetic parameters in pediatric subjects with PREZISTA/ritonavir administered once or twice daily are summarized in the table below:

**Table 13: Population Pharmacokinetic Estimates of Darunavir Exposure (Trials TMC114-C230, TMC114-C212 and TMC114-C228) Following Administration of Doses in Tables 2 and 3**

Parameter	PREZISTA/ritonavir once daily	PREZISTA/ritonavir twice daily		
	TMC114-C230 <sup>β</sup> N=12	TMC114-C212 N=74	TMC114-C228 <sup>*</sup>	
			10 to less than 15 kg <sup>‡</sup> N=10	15 to less than 20 kg <sup>§</sup> N=13
AUC <sub>24h</sub> (ng·h/mL) <sup>†</sup>				
Mean ± Standard Deviation	84390 ± 23587	126377 ± 34356	137896 ± 51420	157760 ± 54080
Median (Range)	86741 (35527–123325)	127340 (67054–230720)	124044 (89688–261090)	132698 (112310–294840)
C <sub>0h</sub> (ng/mL)				
Mean ± Standard Deviation	2141 ± 865	3948 ± 1363	4510 ± 2031	4848 ± 2143
Median (Range)	2234 (542–3776)	3888 (1836–7821)	4126 (2456–9361)	3927 (3046–10292)

N=number of subjects with data.

<sup>\*</sup> Subjects may have contributed pharmacokinetic data to both the 10 kg to less than 15 kg weight group and the 15 kg to less than 20 kg weight group.

<sup>†</sup> AUC<sub>24h</sub> is calculated as AUC<sub>12h</sub>\*2.

<sup>‡</sup> Calculated from individual pharmacokinetic parameters estimated for Week 2 and Week 4, based on the Week 48 analysis that evaluated a darunavir dose of 20 mg/kg twice daily with ritonavir 3 mg/kg twice daily.

<sup>§</sup> The 15 kg to less than 20 kg weight group received 380 mg (3.8 mL) PREZISTA oral suspension twice daily with 48 mg (0.6 mL) ritonavir oral solution twice daily in TMC114-C228. Calculated from individual pharmacokinetic parameters estimated for Week 2 post-dose adjustment visit; Week 24 and Week 48 based on the – Week 48 analysis that evaluated a darunavir dose of 380 mg twice daily.

<sup>β</sup> Summary statistics for population pharmacokinetic parameter estimates for DRV after administration of DRV/rtv at 800/100 mg once daily in treatment-naïve HIV-1 infected subjects from 12 to <18 years of age – Week-48 Analyses.

### Pregnancy and Postpartum

The exposure to total darunavir and ritonavir after intake of PREZISTA/ritonavir 600/100 mg twice daily and PREZISTA/ritonavir 800/100 mg once daily as part of an antiretroviral regimen was generally lower during pregnancy compared with postpartum (see Table 14, Table 15 and Figure 1).

**Table 14: Pharmacokinetic Results of Total Darunavir After Administration of PREZISTA/ritonavir at 600/100 mg Twice Daily as Part of an Antiretroviral Regimen, During the 2<sup>nd</sup> Trimester of Pregnancy, the 3<sup>rd</sup> Trimester of Pregnancy and Postpartum**

Pharmacokinetics of total darunavir (mean ± standard deviation)	2 <sup>nd</sup> Trimester of pregnancy (n=11) <sup>*</sup>	3 <sup>rd</sup> Trimester of pregnancy (n=11)	Postpartum (6-12 Weeks) (n=11)
C <sub>max</sub> , ng/mL	4601 ± 1125	5111 ± 1517	6499 ± 2411
AUC <sub>24h</sub> , ng·h/mL <sup>‡</sup>	77900 ± 20020	87400 ± 32800	110600 ± 54040
C <sub>min</sub> , ng/mL <sup>†</sup>	1980 ± 839.9	2498 ± 1193	2711 ± 2268

<sup>\*</sup> n=10 for AUC<sub>24h</sub>

<sup>†</sup> excluding C<sub>min</sub> value below LLOQ, n=10 for reference

<sup>‡</sup> AUC<sub>24h</sub> is calculated as AUC<sub>12h</sub>\*2.


**Table 15: Pharmacokinetic Results of Total Darunavir After Administration of PREZISTA/ritonavir at 800/100 mg Once Daily as Part of an Antiretroviral Regimen, During the 2<sup>nd</sup> Trimester of Pregnancy, the 3<sup>rd</sup> Trimester of Pregnancy and Postpartum**

Pharmacokinetics of total darunavir (mean ± standard deviation)	2 <sup>nd</sup> Trimester of pregnancy (n=16)	3 <sup>rd</sup> Trimester of pregnancy (n=14)	Postpartum (6-12 Weeks) (n=15)
C <sub>max</sub> , ng/mL	4988 ± 1551	5138 ± 1243	7445 ± 1674
AUC <sub>24h</sub> , ng.h/mL	61303 ± 16232	60439 ± 14052	94529 ± 28572
C <sub>min</sub> , ng/mL*	1193 ± 509	1098 ± 609	1572 ± 1108

\* N=12 for postpartum, N=15 for 2<sup>nd</sup> trimester and N=14 for 3<sup>rd</sup> trimester

Due to an increase in the unbound fraction of darunavir during pregnancy compared to postpartum, unbound darunavir exposures were less reduced during pregnancy as compared to postpartum. Exposure reductions during pregnancy were greater for the once daily regimen as compared to the twice daily regimen (see Figure 1).

**Figure 1: Pharmacokinetic Results (Within-Subject Comparison) of Total and Unbound Darunavir After Administration of PREZISTA/ritonavir at 600/100 mg Twice Daily or 800/100 mg Once Daily as Part of an Antiretroviral Regimen, During the 2<sup>nd</sup> and 3<sup>rd</sup> Trimester of Pregnancy Compared to Postpartum**



Legend: 90% CI: 90% confidence interval; GMR: geometric mean ratio. Solid vertical line: ratio of 1.0; dotted vertical lines: reference lines of 0.8 and 1.25.

## Drug Interactions

[See also *Contraindications (4)*, *Warnings and Precautions (5.5)* and *Drug Interactions (7)*.]

Darunavir co-administered with ritonavir is an inhibitor of CYP3A, CYP2D6, and P-gp. Co-administration of darunavir and ritonavir with drugs primarily metabolized by CYP3A and CYP2D6, or are transported by P-gp, may result in increased plasma concentrations of such drugs, which could increase or prolong their therapeutic effect and adverse events.

Darunavir and ritonavir are metabolized by CYP3A. *In vitro* data indicate that darunavir may be a P-gp substrate. Drugs that induce CYP3A activity would be expected to increase the clearance of darunavir and ritonavir, resulting in lowered plasma concentrations of darunavir and ritonavir. Co-administration of darunavir and ritonavir and other drugs that inhibit CYP3A or P-gp may

decrease the clearance of darunavir and ritonavir and may result in increased plasma concentrations of darunavir and ritonavir.

Drug interaction studies were performed with darunavir and other drugs likely to be co-administered and some drugs commonly used as probes for pharmacokinetic interactions. The effects of co-administration of darunavir on the AUC,  $C_{max}$ , and  $C_{min}$  values are summarized in Table 16 (effect of other drugs on darunavir) and Table 17 (effect of darunavir on other drugs). For information regarding clinical recommendations, *see Drug Interactions (7)*.

Several interaction studies have been performed with a dose other than the recommended dose of the co-administered drug or darunavir; however, the results are applicable to the recommended dose of the co-administered drug and/or darunavir.

**Table 16: Drug Interactions: Pharmacokinetic Parameters for Darunavir in the Presence of Co-Administered Drugs**

Co-administered drug	Dose/Schedule		N	PK	LS Mean ratio (90% CI) of <u>darunavir</u> Pharmacokinetic parameters with/without co-administered drug no effect =1.00		
	Co-administere d Drug	Darunavir/ ritonavir			C <sub>max</sub>	AUC	C <sub>min</sub>
Co-administration with other HIV protease inhibitors							
Atazanavir	300 mg q.d.*	400/100 mg b.i.d. <sup>†</sup>	13	↔	1.02 (0.96-1.09)	1.03 (0.94-1.12)	1.01 (0.88-1.16)
Indinavir	800 mg b.i.d.	400/100 mg b.i.d.	9	↑	1.11 (0.98-1.26)	1.24 (1.09-1.42)	1.44 (1.13-1.82)
Lopinavir/ritonavir	400/100 mg b.i.d.	1200/100 mg b.i.d. <sup>‡</sup>	14	↓	0.79 (0.67-0.92)	0.62 (0.53-0.73)	0.49 (0.39-0.63)
	533/133.3 mg b.i.d.	1200 mg b.i.d. <sup>‡</sup>	15	↓	0.79 (0.64-0.97)	0.59 (0.50-0.70)	0.45 (0.38-0.52)
Saquinavir hard gel capsule	1000 mg b.i.d.	400/100 mg b.i.d.	14	↓	0.83 (0.75-0.92)	0.74 (0.63-0.86)	0.58 (0.47-0.72)
Co-administration with other HIV antiretrovirals							
Didanosine	400 mg q.d.	600/100 mg b.i.d.	17	↔	0.93 (0.86-1.00)	1.01 (0.95-1.07)	1.07 (0.95-1.21)
Efavirenz	600 mg q.d.	300/100 mg b.i.d.	12	↓	0.85 (0.72-1.00)	0.87 (0.75-1.01)	0.69 (0.54-0.87)
Etravirine	200 mg b.i.d.	600/100 mg b.i.d.	15	↔	1.11 (1.01-1.22)	1.15 (1.05-1.26)	1.02 (0.90-1.17)
Nevirapine	200 mg b.i.d.	400/100 mg b.i.d.	8	↑	1.40 <sup>§</sup> (1.14-1.73)	1.24 <sup>§</sup> (0.97-1.57)	1.02 <sup>§</sup> (0.79-1.32)
Rilpivirine	150 mg q.d.	800/100 mg q.d.	15	↔	0.90 (0.81-1.00)	0.89 (0.81-0.99)	0.89 (0.68-1.16)
Tenofovir disoproxil fumarate	300 mg q.d.	300/100 mg b.i.d.	12	↑	1.16 (0.94-1.42)	1.21 (0.95-1.54)	1.24 (0.90-1.69)
Co-administration with HCV NS3-4A protease inhibitors							
Simeprevir	50 mg q.d. <sup>€</sup>	800 mg q.d.	25 <sup>+</sup>	↑	1.04 (0.99-1.10)	1.18 (1.11-1.25)	1.31 (1.13-1.52)

<b>Co-administration with other drugs</b>							
Artemether/lumefantrine	80/480 mg (6 doses at 0, 8, 24, 36, 48, and 60 hours)	600/100 mg b.i.d.	14	↔	1.00 (0.93-1.07)	0.96 (0.90-1.03)	0.87 (0.77-0.98)
Carbamazepine	200 mg b.i.d.	600/100 mg b.i.d.	16	↔	1.04 (0.93-1.16)	0.99 (0.90-1.08)	0.85 (0.73-1.00)
Clarithromycin	500 mg b.i.d.	400/100 mg b.i.d.	17	↔	0.83 (0.72-0.96)	0.87 (0.75-1.01)	1.01 (0.81-1.26)
Ketoconazole	200 mg b.i.d.	400/100 mg b.i.d.	14	↑	1.21 (1.04-1.40)	1.42 (1.23-1.65)	1.73 (1.39-2.14)
Omeprazole	20 mg q.d.	400/100 mg b.i.d.	16	↔	1.02 (0.95-1.09)	1.04 (0.96-1.13)	1.08 (0.93-1.25)
Paroxetine	20 mg q.d.	400/100 mg b.i.d.	16	↔	0.97 (0.92-1.02)	1.02 (0.95-1.10)	1.07 (0.96-1.19)
Pitavastatin	4 mg q.d.	800/100 mg q.d.	27	↔	1.06 (1.00-1.12)	1.03 (0.95-1.12)	NA
Ranitidine	150 mg b.i.d.	400/100 mg b.i.d.	16	↔	0.96 (0.89-1.05)	0.95 (0.90-1.01)	0.94 (0.90-0.99)
Rifabutin	150 mg q.o.d. <sup>¶</sup>	600/100 mg b.i.d.	11	↑	1.42 (1.21-1.67)	1.57 (1.28-1.93)	1.75 (1.28-2.37)
Sertraline	50 mg q.d.	400/100 mg b.i.d.	13	↔	1.01 (0.89-1.14)	0.98 (0.84-1.14)	0.94 (0.76-1.16)

N = number of subjects with data

\* q.d. = once daily

† b.i.d. = twice daily

‡ The pharmacokinetic parameters of darunavir in this study were compared with the pharmacokinetic parameters following administration of PREZISTA/ritonavir 600/100 mg twice daily.

§ Ratio based on between-study comparison.

¶ q.o.d. = every other day

€ The dose of simeprevir in this interaction study was 50 mg when co-administered in combination with PREZISTA/ritonavir compared to 150 mg once daily in the simeprevir alone treatment group.

+ Maximum number of subjects

**Table 17: Drug Interactions: Pharmacokinetic Parameters for Co-Administered Drugs in the Presence of PREZISTA/ritonavir**

Co-administered drug	Dose/Schedule		N	PK	LS Mean ratio (90% CI) of <u>co-administered drug</u> pharmacokinetic parameters with/without darunavir no effect =1.00		
	Co-administered drug	Darunavir/ ritonavir			C <sub>max</sub>	AUC	C <sub>min</sub>
Co-administration with other HIV protease inhibitors							
Atazanavir	300 mg q.d.* /100 mg ritonavir q.d. when administered alone  300 mg q.d. when administered with darunavir/ ritonavir	400/100 mg b.i.d.†	13	↔	0.89 (0.78-1.01)	1.08 (0.94-1.24)	1.52 (0.99-2.34)
Indinavir	800 mg b.i.d.	400/100 mg	9	↑	1.08	1.23	2.25

	/100 mg ritonavir b.i.d. when administered alone  800 mg b.i.d. when administered with darunavir/ ritonavir	b.i.d.			(0.95-1.22)	(1.06-1.42)	(1.63-3.10)
Lopinavir/ritonavir	400/100 mg b.i.d. <sup>‡</sup>	1200/100 m g b.i.d.	14	↔	0.98 (0.78-1.22)	1.09 (0.86-1.37)	1.23 (0.90-1.69)
	533/133.3 mg b.i.d. <sup>‡</sup>	1200 mg b.i.d.	15	↔	1.11 (0.96-1.30)	1.09 (0.96-1.24)	1.13 (0.90-1.42)
Saquinavir hard gel capsule	1000 mg b.i.d. /100 mg ritonavir b.i.d. when administered alone  1000 mg b.i.d. when administered with darunavir/ ritonavir	400/100 mg b.i.d.	12	↔	0.94 (0.78-1.13)	0.94 (0.76-1.17)	0.82 (0.52-1.30)
Co-administration with other HIV antiretrovirals							
Didanosine	400 mg q.d.	600/100 mg b.i.d.	17	↔	0.84 (0.59-1.20)	0.91 (0.75-1.10)	-
Dolutegravir	30 mg q.d	600/100 mg b.i.d.	15	↓	0.89 (0.83-0.97)	0.78 (0.72-0.85)	0.62 <sup>Ω</sup> (0.56-0.69)
Dolutegravir	50 mg q.d.	600/100 mg b.i.d. with 200 mg b.i.d. etravirine	9	↓	0.88 (0.78-1.00)	0.75 (0.69-0.81)	0.63 <sup>Ω</sup> (0.52-0.76)
Efavirenz	600 mg q.d.	300/100 mg b.i.d.	12	↑	1.15 (0.97-1.35)	1.21 (1.08-1.36)	1.17 (1.01-1.36)
Etravirine	100 mg b.i.d.	600/100 mg b.i.d.	14	↓	0.68 (0.57-0.82)	0.63 (0.54-0.73)	0.51 (0.44-0.61)
Nevirapine	200 mg b.i.d.	400/100 mg b.i.d.	8	↑	1.18 (1.02-1.37)	1.27 (1.12-1.44)	1.47 (1.20-1.82)
Rilpivirine	150 mg q.d.	800/100 mg q.d.	14	↑	1.79 (1.56-2.06)	2.30 (1.98-2.67)	2.78 (2.39-3.24)
Tenofovir disoproxil fumarate	300 mg q.d.	300/100 mg b.i.d.	12	↑	1.24 (1.08-1.42)	1.22 (1.10-1.35)	1.37 (1.19-1.57)
Maraviroc	150 mg b.i.d.	600/100 mg b.i.d.	12	↑	2.29 (1.46-3.59)	4.05 (2.94-5.59)	8.00 (6.35-10.1)
		600/100 mg b.i.d. with 200 mg b.i.d. etravirine	10	↑	1.77 (1.20-2.60)	3.10 (2.57-3.74)	5.27 (4.51-6.15)
Co-administration with HCV NS3-4A protease inhibitors							
Simeprevir	50 mg q.d. <sup>€</sup>	800/100 mg q.d.	25 <sup>+</sup>	↑	1.79 (1.55-2.06)	2.59 (2.15-3.11)	4.58 (3.54-5.92)
Co-administration with other drugs							
Atorvastatin	40 mg q.d. when administered alone	300/100 mg b.i.d.	15	↑	0.56 (0.48-0.67)	0.85 (0.76-0.97)	1.81 (1.37-2.40)

	10 mg q.d. when administered with darunavir/ritonavir						
Artemether Dihydroartemisinin	80 mg single dose	600/100 mg b.i.d.	15	↓	0.85 (0.68-1.05)	0.91 (0.78-1.06)	-
			15	↑	1.06 (0.82-1.39)	1.12 (0.96-1.30)	-
Artemether Dihydroartemisinin	artemether/lumefantrine 80/480 mg	600/100 mg b.i.d.	15	↓	0.82 (0.61-1.11)	0.84 (0.69-1.02)	0.97 (0.90-1.05)
Lumefantrine	(6 doses at 0, 8, 24, 36, 48, and 60 hours)		15	↓	0.82 (0.66-1.01)	0.82 (0.74-0.91)	1.00 (0.82-1.22)
			15	↑	1.65 (1.49-1.83)	2.75 (2.46-3.08)	2.26 (1.92-2.67)
Buprenorphine/Naloxone	8/2 mg to 16/4 mg q.d.	600/100 mg b.i.d.	17	↔	0.92 <sup>§</sup> (0.79-1.08)	0.89 <sup>§</sup> (0.78-1.02)	0.98 <sup>§</sup> (0.82-1.16)
Norbuprenorphine			17	↑	1.36 (1.06-1.74)	1.46 (1.15-1.85)	1.71 (1.29-2.27)
Carbamazepine	200 mg b.i.d.	600/100 mg b.i.d.	16	↑	1.43 (1.34-1.53)	1.45 (1.35-1.57)	1.54 (1.41-1.68)
Carbamazepine epoxide			16	↓	0.46 (0.43-0.49)	0.46 (0.44-0.49)	0.48 (0.45-0.51)
Clarithromycin	500 mg b.i.d.	400/100 mg b.i.d.	17	↑	1.26 (1.03-1.54)	1.57 (1.35-1.84)	2.74 (2.30-3.26)
Dextromethorphan	30 mg	600/100 mg b.i.d.	12	↑	2.27 (1.59-3.26)	2.70 (1.80-4.05)	-
Dextrorphan				↓	0.87 (0.77-0.98)	0.96 (0.90-1.03)	-
Digoxin	0.4 mg	600/100 mg b.i.d.	8	↑	1.15 (0.89-1.48)	1.36 (0.81-2.27)	-
Ethinyl estradiol (EE)	Ortho-Novum 1/35 (35 µg EE /1 mg NE)	600/100 mg b.i.d.	11	↓	0.68 (0.61-0.74)	0.56 (0.50-0.63)	0.38 (0.27-0.54)
Norethindrone (NE)			11	↓	0.90 (0.83-0.97)	0.86 (0.75-0.98)	0.70 (0.51-0.97)
Ketoconazole	200 mg b.i.d.	400/100 mg b.i.d.	15	↑	2.11 (1.81-2.44)	3.12 (2.65-3.68)	9.68 (6.44-14.55)
R-Methadone	55-150 mg q.d.	600/100 mg b.i.d.	16	↓	0.76 (0.71-0.81)	0.84 (0.78-0.91)	0.85 (0.77-0.94)
Omeprazole	40 mg single dose	600/100 mg b.i.d.	12	↓	0.66 (0.48-0.90)	0.58 (0.50-0.66)	-
5-hydroxy omeprazole				↓	0.93 (0.71-1.21)	0.84 (0.77-0.92)	-
Paroxetine	20 mg q.d.	400/100 mg b.i.d.	16	↓	0.64 (0.59-0.71)	0.61 (0.56-0.66)	0.63 (0.55-0.73)
Pitavastatin	4 mg q.d.	800/100 mg q.d.	27	↓	0.96 (0.84-1.09)	0.74 (0.69-0.80)	NA
Pravastatin	40 mg single dose	600/100 mg b.i.d.	14	↑	1.63 (0.95-2.82)	1.81 (1.23-2.66)	-
Rifabutin	150 mg	600/100 mg	11	↑	0.72	0.93	1.64

25- <i>O</i> -desacetyl-rifabutin	q.o.d. <sup>¶</sup> when administered with PREZISTA/ritonavir 300 mg q.d. when administered alone	b.i.d. <sup>#</sup>	11	↑	(0.55-0.93) 4.77 (4.04-5.63)	(0.80-1.09) 9.81 (8.09-11.9)	(1.48-1.81) 27.1 (22.2-33.2)
Sertraline	50 mg q.d.	400/100 mg b.i.d.	13	↓	0.56 (0.49-0.63)	0.51 (0.46-0.58)	0.51 (0.45-0.57)
Sildenafil	100 mg (single dose) administered alone  25 mg (single dose) when administered with darunavir/ritonavir	400/100 mg b.i.d.	16	↑	0.62 (0.55-0.70)	0.97 (0.86-1.09)	-
S-warfarin	10 mg single dose	600/100 mg b.i.d.	12	↓	0.92 (0.86-0.97)	0.79 (0.73-0.85)	-
7-OH-S-warfarin			12	↑	1.42 (1.24-1.63)	1.23 (0.97-1.57)	-

N = number of subjects with data; - = no information available

\* q.d. = once daily

† b.i.d. = twice daily

‡ The pharmacokinetic parameters of lopinavir in this study were compared with the pharmacokinetic parameters following administration of lopinavir/ritonavir 400/100 mg twice daily.

§ ratio is for buprenorphine; mean  $C_{max}$  and  $AUC_{24}$  for naloxone were comparable when buprenorphine/naloxone was administered with or without PREZISTA/ritonavir

¶ q.o.d. = every other day

# In comparison to rifabutin 300 mg once daily.

Ω Noted as  $C_{\tau}$  or  $C_{24}$  in the dolutegravir U.S. prescribing information

+ Maximum number of subjects

€ The dose of simeprevir in this interaction study was 50 mg when co-administered in combination with PREZISTA/ritonavir compared to 150 mg once daily in the simeprevir alone treatment group.

A cocktail study was conducted in 12 healthy volunteers to evaluate the effect of steady state pharmacokinetics of PREZISTA/ritonavir on the activity of CYP2D6 (using dextromethorphan as probe substrate), CYP2C9 (using warfarin as probe substrate), and CYP2C19 (using omeprazole as probe substrate). The pharmacokinetic results are shown in Table 17.

## 12.4 Microbiology

### Mechanism of Action

Darunavir is an inhibitor of the HIV-1 protease. It selectively inhibits the cleavage of HIV-1 encoded Gag-Pol polyproteins in infected cells, thereby preventing the formation of mature virus particles.

### Antiviral Activity

Darunavir exhibits activity against laboratory strains and clinical isolates of HIV-1 and laboratory strains of HIV-2 in acutely infected T-cell lines, human peripheral blood mononuclear cells and human monocytes/macrophages with median  $EC_{50}$  values ranging from 1.2 to 8.5 nM (0.7 to 5.0 ng/mL). Darunavir demonstrates antiviral activity in cell culture against a broad panel



of HIV-1 group M (A, B, C, D, E, F, G), and group O primary isolates with EC<sub>50</sub> values ranging from less than 0.1 to 4.3 nM. The EC<sub>50</sub> value of darunavir increases by a median factor of 5.4 in the presence of human serum. Darunavir did not show antagonism when studied in combination with the PIs amprenavir, atazanavir, indinavir, lopinavir, nelfinavir, ritonavir, saquinavir, or tipranavir, the N(t)RTIs abacavir, didanosine, emtricitabine, lamivudine, stavudine, tenofovir, zalcitabine, or zidovudine, the NNRTIs delavirdine, rilpivirine, efavirenz, etravirine, or nevirapine, and the fusion inhibitor enfuvirtide.

## Resistance

*Cell Culture:* HIV-1 isolates with a decreased susceptibility to darunavir have been selected in cell culture and obtained from subjects treated with PREZISTA/ritonavir. Darunavir-resistant virus derived in cell culture from wild-type HIV-1 had 21- to 88-fold decreased susceptibility to darunavir and developed 2 to 4 of the following amino acid substitutions S37D, R41E/T, K55Q, H69Q, K70E, T74S, V77I, or I85V in the protease. Selection in cell culture of darunavir resistant HIV-1 from nine HIV-1 strains harboring multiple PI resistance-associated mutations resulted in the overall emergence of 22 mutations in the protease gene, coding for amino acid substitutions L10F, V11I, I13V, I15V, G16E, L23I, V32I, L33F, S37N, M46I, I47V, I50V, F53L, L63P, A71V, G73S, L76V, V82I, I84V, T91A/S, and Q92R, of which L10F, V32I, L33F, S37N, M46I, I47V, I50V, L63P, A71V, and I84V were the most prevalent. These darunavir-resistant viruses had at least eight protease substitutions and exhibited 50- to 641-fold decreases in darunavir susceptibility with final EC<sub>50</sub> values ranging from 125 nM to 3461 nM.

*Clinical trials of PREZISTA/ritonavir in treatment-experienced subjects:* In a pooled analysis of the 600/100 mg PREZISTA/ritonavir twice daily arms of trials TMC114-C213, TMC114-C202, TMC114-C215, and the control arms of etravirine trials TMC125-C206 and TMC125-C216, the amino acid substitutions V32I and I54L or M developed most frequently on PREZISTA/ritonavir in 41% and 25%, respectively, of the treatment-experienced subjects who experienced virologic failure, either by rebound or by never being suppressed (less than 50 copies/mL). Other substitutions that developed frequently in PREZISTA/ritonavir virologic failure isolates occurred at amino acid positions V11I, I15V, L33F, I47V, I50V, and L89V. These amino acid substitutions were associated with decreased susceptibility to darunavir; 90% of the virologic failure isolates had a greater than 7-fold decrease in susceptibility to darunavir at failure. The median darunavir phenotype (fold change from reference) of the virologic failure isolates was 4.3-fold at baseline and 85-fold at failure. Amino acid substitutions were also observed in the protease cleavage sites in the Gag polyprotein of some PREZISTA/ritonavir virologic failure isolates. In trial TMC114-C212 of treatment-experienced pediatric subjects, the amino acid substitutions V32I, I54L and L89M developed most frequently in virologic failures on PREZISTA/ritonavir.

In the 96-week as-treated analysis of the Phase 3 trial TMC114-C214, the percent of virologic failures (never suppressed, rebounders and discontinued before achieving suppression) was 21% (62/298) in the group of subjects receiving PREZISTA/ritonavir 600/100 mg twice daily compared to 32% (96/297) of subjects receiving lopinavir/ritonavir 400/100 mg twice daily.

Examination of subjects who failed on PREZISTA/ritonavir 600/100 mg twice daily and had post-baseline genotypes and phenotypes showed that 7 subjects (7/43; 16%) developed PI substitutions on PREZISTA/ritonavir treatment resulting in decreased susceptibility to darunavir. Six of the 7 had baseline PI resistance-associated substitutions and baseline darunavir phenotypes greater than 7. The most common emerging PI substitutions in these virologic failures were V32I, L33F, M46I or L, I47V, I54L, T74P and L76V. These amino acid substitutions were associated with 59- to 839-fold decreased susceptibility to darunavir at failure. Examination of individual subjects who failed in the comparator arm on lopinavir/ritonavir and had post-baseline genotypes and phenotypes showed that 31 subjects (31/75; 41%) developed substitutions on lopinavir treatment resulting in decreased susceptibility to lopinavir (greater than 10-fold) and the most common substitutions emerging on treatment were L10I or F, M46I or L, I47V or A, I54V and L76V. Of the 31 lopinavir/ritonavir virologic failure subjects, 14 had reduced susceptibility (greater than 10-fold) to lopinavir at baseline.

In the 48-week analysis of the Phase 3 trial TMC114-C229, the number of virologic failures (including those who discontinued before suppression after Week 4) was 26% (75/294) in the group of subjects receiving PREZISTA/ritonavir 800/100 mg once daily compared to 19% (56/296) of subjects receiving PREZISTA/ritonavir 600/100 mg twice daily. Examination of isolates from subjects who failed on PREZISTA/ritonavir 800/100 mg once daily and had post-baseline genotypes showed that 8 subjects (8/60; 13%) had isolates that developed IAS-USA defined PI resistance-associated substitutions compared to 5 subjects (5/39; 13%) on PREZISTA/ritonavir 600/100 mg twice daily. Isolates from 2 subjects developed PI resistance associated substitutions associated with decreased susceptibility to darunavir; 1 subject isolate in the PREZISTA/ritonavir 800/100 mg once daily arm, developed substitutions V32I, M46I, L76V and I84V associated with a 24-fold decreased susceptibility to darunavir, and 1 subject isolate in the PREZISTA/ritonavir 600/100 mg twice daily arm developed substitutions L33F and I50V associated with a 40-fold decreased susceptibility to darunavir. In the PREZISTA/ritonavir 800/100 mg once daily and PREZISTA/ritonavir 600/100 mg twice daily groups, isolates from 7 (7/60; 12%) and 4 (4/42; 10%) virologic failures, respectively, developed decreased susceptibility to an NRTI included in the treatment regimen.

*Clinical trials of PREZISTA/ritonavir in treatment-naïve subjects:* In the 192-week as-treated analysis censoring those who discontinued before Week 4 of the Phase 3 trial TMC114-C211, the percentage of virologic failures (never suppressed, rebounders and discontinued before achieving suppression) was 22% (64/288) in the group of subjects receiving PREZISTA/ritonavir 800/100 mg once daily compared to 29% (76/263) of subjects receiving lopinavir/ritonavir 800/200 mg per day. In the PREZISTA/ritonavir arm, emergent PI resistance-associated substitutions were identified in 11 of the virologic failures with post-baseline genotypic data (n=43). However, none of the darunavir virologic failures had a decrease in darunavir susceptibility (greater than 7-fold change) at failure. In the comparator lopinavir/ritonavir arm, emergent PI resistance-associated substitutions were identified in 17 of the virologic failures with post-baseline genotypic data (n=53), but none of the lopinavir/ritonavir virologic failures had decreased susceptibility to lopinavir (greater than 10-

fold change) at failure. The reverse transcriptase M184V substitution and/or resistance to emtricitabine, which was included in the fixed background regimen, was identified in 4 virologic failures from the PREZISTA/ritonavir arm and 7 virologic failures in the lopinavir/ritonavir arm.

### Cross-resistance

Cross-resistance among PIs has been observed. Darunavir has a less than 10-fold decreased susceptibility in cell culture against 90% of 3309 clinical isolates resistant to amprenavir, atazanavir, indinavir, lopinavir, nelfinavir, ritonavir, saquinavir and/or tipranavir showing that viruses resistant to these PIs remain susceptible to darunavir.

Darunavir-resistant viruses were not susceptible to amprenavir, atazanavir, indinavir, lopinavir, nelfinavir, ritonavir or saquinavir in cell culture. However, six of nine darunavir-resistant viruses selected in cell culture from PI-resistant viruses showed a fold change in EC<sub>50</sub> values less than 3 for tipranavir, indicative of limited cross-resistance between darunavir and tipranavir. In trials TMC114-C213, TMC114-C202, and TMC114-C215, 34% (64/187) of subjects in the PREZISTA/ritonavir arm whose baseline isolates had decreased susceptibility to tipranavir (tipranavir fold change greater than 3) achieved less than 50 copies/mL serum HIV-1 RNA levels at Week 96. Of the viruses isolated from subjects experiencing virologic failure on PREZISTA/ritonavir 600/100 mg twice daily (greater than 7-fold change), 41% were still susceptible to tipranavir and 10% were susceptible to saquinavir while less than 2% were susceptible to the other protease inhibitors (amprenavir, atazanavir, indinavir, lopinavir or nelfinavir).

In trial TMC114-C214, the 7 PREZISTA/ritonavir virologic failures with reduced susceptibility to darunavir at failure were also resistant to the approved PIs (fos)amprenavir, atazanavir, lopinavir, indinavir, and nelfinavir at failure. Six of these 7 were resistant to saquinavir and 5 were resistant to tipranavir. Four of these virologic failures were already PI-resistant at baseline.

Cross-resistance between darunavir and nucleoside/nucleotide reverse transcriptase inhibitors, non-nucleoside reverse transcriptase inhibitors, fusion inhibitors, CCR5 co-receptor antagonists, or integrase inhibitors is unlikely because the viral targets are different.

### Baseline Genotype/Phenotype and Virologic Outcome Analyses

Genotypic and/or phenotypic analysis of baseline virus may aid in determining darunavir susceptibility before initiation of PREZISTA/ritonavir 600/100 mg twice daily therapy. The effect of baseline genotype and phenotype on virologic response at 96 weeks was analyzed in as-treated analyses using pooled data from the Phase 2b trials (Trials TMC114-C213, TMC114-C202, and TMC114-C215) (n=439). The findings were confirmed with additional genotypic and phenotypic data from the control arms of etravirine trials TMC125-C206 and TMC125-C216 at Week 24 (n=591).

Diminished virologic responses were observed in subjects with 5 or more baseline IAS-defined primary protease inhibitor resistance-associated substitutions (D30N, V32I, L33F, M46I/L, I47A/V, G48V, I50L/V, I54L/M, L76V, V82A/F/L/S/T, I84V, N88S, L90M) (see Table 18).

**Table 18: Response to PREZISTA/ritonavir 600/100 mg Twice Daily by Baseline Number of IAS-Defined Primary PI Resistance-Associated Substitutions: As-treated Analysis of Trials TMC114-C213, TMC114-C202, and TMC114-C215**

# IAS-defined primary PI substitutions	Proportion of subjects with <50 copies/mL at Week 96 N=439		
	Overall	<i>de novo</i> ENF	Re-used/No ENF
<b>All</b>	44% (192/439)	54% (61/112)	40% (131/327)
<b>0 – 4</b>	50% (162/322)	58% (49/85)	48% (113/237)
<b>5</b>	22% (16/74)	47% (9/19)	13% (7/55)
<b>≥6</b>	9% (3/32)	17% (1/6)	8% (2/26)

ENF=enfuvirtide

IAS Primary PI Substitutions (2008): D30N, V32I, L33F, M46I/L, I47A/V, G48V, I50L/V, I54L/M, L76V, V82A/F/L/S/T, I84V, N88S, L90M

The presence at baseline of two or more of the substitutions V11I, V32I, L33F, I47V, I50V, I54L or M, T74P, L76V, I84V or L89V was associated with a decreased virologic response to PREZISTA/ritonavir. In subjects not taking enfuvirtide *de novo*, the proportion of subjects achieving viral load less than 50 plasma HIV-1 RNA copies/mL at 96 weeks was 59%, 29%, and 12% when the baseline genotype had 0-1, 2 and greater than or equal to 3 of these substitutions, respectively.

Baseline darunavir phenotype (shift in susceptibility relative to reference) was shown to be a predictive factor of virologic outcome. Response rates assessed by baseline darunavir phenotype are shown in Table 19. These baseline phenotype groups are based on the select patient populations in the trials TMC114-C213, TMC114-C202, and TMC114-C215, and are not meant to represent definitive clinical susceptibility breakpoints for PREZISTA/ritonavir. The data are provided to give clinicians information on the likelihood of virologic success based on pre-treatment susceptibility to darunavir.

**Table 19: Response (HIV-1 RNA <50 copies/mL at Week 96) to PREZISTA/ritonavir 600/100 mg Twice Daily by Baseline Darunavir Phenotype and by Use of Enfuvirtide: As-treated Analysis of Trials TMC114-C213, TMC114-C202, and TMC114-C215**

Baseline DRV phenotype	Proportion of subjects with <50 copies/mL at Week 96 N=417		
	All	<i>de novo</i> ENF	Re-used/No ENF
<b>Overall</b>	175/417 (42%)	61/112 (54%)	131/327 (40%)
<b>0 – 7</b>	148/270 (55%)	44/65 (68%)	104/205 (51%)
<b>&gt;7 – 20</b>	16/53 (30%)	7/17 (41%)	9/36 (25%)
<b>&gt;20</b>	11/94 (12%)	6/23 (26%)	5/71 (7%)

ENF=enfuvirtide

## 13 NONCLINICAL TOXICOLOGY

### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

#### Carcinogenesis and Mutagenesis

Darunavir was evaluated for carcinogenic potential by oral gavage administration to mice and rats up to 104 weeks. Daily doses of 150, 450 and 1000 mg/kg were administered to mice and doses of 50, 150 and 500 mg/kg was administered to rats. A dose-related increase in the incidence of hepatocellular adenomas and carcinomas were observed in males and females of both species as well as an increase in thyroid follicular cell adenomas in male rats. The observed hepatocellular findings in rodents are considered to be of limited relevance to humans. Repeated administration of darunavir to rats caused hepatic microsomal enzyme induction and increased thyroid hormone elimination, which predispose rats, but not humans, to thyroid neoplasms. At the highest tested doses, the systemic exposures to darunavir (based on AUC) were between 0.4- and 0.7-fold (mice) and 0.7- and 1-fold (rats), relative to those observed in humans at the recommended therapeutic doses (600/100 mg twice daily or 800/100 mg once daily).

Darunavir was not mutagenic or genotoxic in a battery of *in vitro* and *in vivo* assays including bacterial reserve mutation (Ames), chromosomal aberration in human lymphocytes and *in vivo* micronucleus test in mice.

#### Impairment of Fertility

No effects on fertility or early embryonic development were observed with darunavir in rats.

## 14 CLINICAL STUDIES

### 14.1 Description of Adult Clinical Trials

The evidence of efficacy of PREZISTA/ritonavir is based on the analyses of 192-week data from a randomized, controlled open-label Phase 3 trial in treatment-naïve (TMC114-C211) HIV-1-infected adult subjects and 96-week data from a randomized, controlled, open-label Phase 3 trial in antiretroviral treatment-experienced (TMC114-C214) HIV-1-infected adult subjects. In addition, 96-week data are included from 2 randomized, controlled Phase 2b trials, TMC114-C213 and TMC114-C202, in antiretroviral treatment-experienced HIV-1-infected adult subjects.

### 14.2 Treatment-Naïve Adult Subjects

#### TMC114-C211

TMC114-C211 is a randomized, controlled, open-label Phase 3 trial comparing PREZISTA/ritonavir 800/100 mg once daily versus lopinavir/ritonavir 800/200 mg per day (given as a twice daily or as a once daily regimen) in antiretroviral treatment-naïve HIV-1-infected adult subjects. Both arms used a fixed background regimen consisting of tenofovir disoproxil fumarate 300 mg once daily (TDF) and emtricitabine 200 mg once daily (FTC).

HIV-1-infected subjects who were eligible for this trial had plasma HIV-1 RNA greater than or equal to 5000 copies/mL. Randomization was stratified by screening plasma viral load (HIV-1 RNA less than 100,000 copies/mL or greater than or equal to 100,000 copies/mL) and screening

CD4+ cell count (less than 200 cells/mm<sup>3</sup> or greater than or equal to 200 cells/mm<sup>3</sup>). Virologic response was defined as a confirmed plasma HIV-1 RNA viral load less than 50 copies/mL. Analyses included 689 subjects in trial TMC114-C211 who had completed 192 weeks of treatment or discontinued earlier.

Demographics and baseline characteristics were balanced between the PREZISTA/ritonavir arm and the lopinavir/ritonavir arm (see Table 20). Table 20 compares the demographic and baseline characteristics between subjects in the PREZISTA/ritonavir 800/100 mg once daily arm and subjects in the lopinavir/ritonavir 800/200 mg per day arm in trial TMC114-C211.

**Table 20: Demographic and Baseline Characteristics of Subjects in Trial TMC114-C211**

	<b>PREZISTA/ritonavir 800/100 mg once daily + TDF/FTC N=343</b>	<b>lopinavir/ritonavir 800/200 mg per day + TDF/FTC N=346</b>
<b>Demographic characteristics</b>		
Median age (years) (range, years)	34 (18-70)	33 (19-68)
Sex		
Male	70%	70%
Female	30%	30%
Race		
White	40%	45%
Black	23%	21%
Hispanic	23%	22%
Asian	13%	11%
<b>Baseline characteristics</b>		
Mean baseline plasma HIV-1 RNA (log <sub>10</sub> copies/mL)	4.86	4.84
Median baseline CD4+ cell count (cells/mm <sup>3</sup> ) (range, cells/mm <sup>3</sup> )	228 (4-750)	218 (2-714)
Percentage of patients with baseline viral load ≥100,000 copies/mL	34%	35%
Percentage of patients with baseline CD4+ cell count <200 cells/mm <sup>3</sup>	41%	43%

FTC=emtricitabine; TDF=tenofovir disoproxil fumarate

Week 192 outcomes for subjects on PREZISTA/ritonavir 800/100 mg once daily from trial TMC114-C211 are shown in Table 21.

**Table 21: Virologic Outcome of Randomized Treatment of Trial TMC114-C211 at 192 Weeks**

	<b>PREZISTA/ritonavir 800/100 mg once daily + TDF/FTC N=343</b>	<b>lopinavir/ritonavir 800/200 mg per day + TDF/FTC N=346</b>
Virologic success HIV-1 RNA <50 copies/mL	70% <sup>*</sup>	61%
Virologic failure <sup>†</sup>	12%	15%
No virologic data at Week 192 window <sup>‡</sup>		
Reasons		
Discontinued trial due to adverse event or death <sup>§</sup>	5%	13%
Discontinued trial for other reasons <sup>  </sup>	13%	12%
Missing data during window <sup>‡</sup> but on trial	<1%	0%

N = total number of subjects with data; FTC=emtricitabine; TDF=tenofovir disoproxil fumarate

\* 95% CI: 1.9; 16.1

† Includes patients who discontinued prior to Week 192 for lack or loss of efficacy and patients who are  $\geq 50$  copies in the 192-week window and patients who had a change in their background regimen that was not permitted by the protocol.

‡ Window 186-198 Weeks.

§ Includes patients who discontinued due to adverse event or death at any time point from Day 1 through the time window if this resulted in no virologic data on treatment during the specified window.

¶ Other includes: withdrew consent, loss to follow-up, etc., if the viral load at the time of discontinuation was  $< 50$  copies/mL

In trial TMC114-C211 at 192 weeks of treatment, the median increase from baseline in CD4+ cell counts was 258 cells/mm<sup>3</sup> in the PREZISTA/ritonavir 800/100 mg once daily arm and 263 cells/mm<sup>3</sup> in the lopinavir/ritonavir 800/200 mg per day arm. Of the PREZISTA/ritonavir subjects with a confirmed virologic response of  $< 50$  copies/mL at Week 48, 81% remained undetectable at Week 192 versus 68% with lopinavir/ritonavir. In the 192 week analysis, statistical superiority of the PREZISTA/ritonavir regimen over the lopinavir/ritonavir regimen was demonstrated for both ITT and OP populations.

### 14.3 Treatment-Experienced Adult Subjects

#### TMC114-C229

TMC114-C229 is a randomized, open-label trial comparing PREZISTA/ritonavir 800/100 mg once daily to PREZISTA/ritonavir 600/100 mg twice daily in treatment-experienced HIV-1-infected patients with screening genotype resistance test showing no darunavir resistance associated substitutions (i.e. V11I, V32I, L33F, I47V, I50V, I54L, I54M, T74P, L76V, I84V, L89V) and a screening viral load of greater than 1,000 HIV-1 RNA copies/mL. Both arms used an optimized background regimen consisting of greater than or equal to 2 NRTIs selected by the investigator.

HIV-1-infected subjects who were eligible for this trial were on a highly active antiretroviral therapy regimen (HAART) for at least 12 weeks. Virologic response was defined as a confirmed plasma HIV-1 RNA viral load less than 50 copies/mL. Analyses included 590 subjects who had completed 48 weeks of treatment or discontinued earlier.

Table 22 compares the demographic and baseline characteristics between subjects in the PREZISTA/ritonavir 800/100 mg once daily arm and subjects in the PREZISTA/ritonavir 600/100 mg twice daily arm in trial TMC114-C229. No imbalances between the 2 arms were noted.

**Table 22: Demographic and Baseline Characteristics of Subjects in Trial TMC114-C229**

	<b>PREZISTA/ritonavir 800/100 mg once daily + OBR N=294</b>	<b>PREZISTA/ritonavir 600/100 mg twice daily + OBR N=296</b>
<b>Demographic characteristics</b>		
Median age (years) (range, years)	40 (18-70)	40 (18-77)
Sex		
Male	61%	67%
Female	39%	33%

Race		
White	35%	37%
Black	28%	24%
Hispanic	16%	20%
Asian	16%	14%
<b>Baseline characteristics</b>		
Mean baseline plasma HIV-1 RNA (log <sub>10</sub> copies/mL)	4.19	4.13
Median baseline CD4+ cell count (cells/mm <sup>3</sup> ) (range, cells/mm <sup>3</sup> )	219 (24-1306)	236 (44-864)
Percentage of patients with baseline viral load ≥100,000 copies/mL	13%	11%
Percentage of patients with baseline CD4+ cell count <200 cells/mm <sup>3</sup>	43%	39%
Median darunavir fold change (range)*	0.50 (0.1-1.8)	0.50 (0.1-1.9)
Median number of resistance-associated†:		
PI mutations	3	4
NNRTI mutations	2	1
NRTI mutations	1	1
Percentage of subjects susceptible to all available PIs at baseline	88%	86%
Percentage of subjects with number of baseline primary protease inhibitor mutations‡:		
0	84%	84%
1	8%	9%
2	5%	4%
≥3	3%	2%
Median number of ARVs previously used‡:		
NRTIs	3	3
NNRTIs	1	1
PIs (excluding low-dose ritonavir)	1	1

OBR=optimized background regimen

\* Based on phenotype (Antivirogram®)

† Johnson VA, Brun-Vézinet F, Clotet B, et al. Update of the drug resistance mutations in HIV-1: December 2008. Top HIV Med 2008; 16(5): 138-145

‡ Only counting ARVs, excluding low-dose ritonavir

Week 48 outcomes for subjects on PREZISTA/ritonavir 800/100 mg once daily from trial TMC114-C229 are shown in Table 23.

**Table 23: Virologic Outcome of Randomized Treatment of Trial TMC114-C229 at 48 Weeks**

	<b>PREZISTA/ritonavir 800/100 mg once daily + OBR N=294</b>	<b>PREZISTA/ritonavir 600/100 mg twice daily + OBR N=296</b>
Virologic success HIV-1 RNA <50 copies/mL	69%	69%
Virologic failure*	26%	23%
No virologic data at Week 48 window†		
Reasons		
Discontinued trial due to adverse event or death‡	3%	4%
Discontinued trial for other reasons§	2%	3%
Missing data during window† but on trial	0%	<1%

N = total number of subjects with data; OBR=optimized background regimen

\* Includes patients who discontinued prior to Week 48 for lack or loss of efficacy, patients who are ≥50 copies in the 48-week window, patients who had a change in their background regimen that was not permitted in the protocol (provided the switch occurred before the



earliest onset of an AE leading to permanent stop of trial medication) and patients who discontinued for reasons other than AEs/death and lack or loss of efficacy (provided their last available viral load was detectable (HIV RNA  $\geq 50$  copies/mL).

† Window 42-54 Weeks

‡ Patients who discontinued due to adverse event or death at any time point from Day 1 through the time window if this resulted in no virologic data on treatment during the specified window.

§ Other includes: withdrew consent, loss to follow-up, etc., if the viral load at the time of discontinuation was  $<50$  copies/mL.

The mean increase from baseline in CD4+ cell counts was comparable for both treatment arms (108 cells/mm<sup>3</sup> and 112 cells/mm<sup>3</sup> in the PREZISTA/ritonavir 800/100 mg once daily arm and the PREZISTA/ritonavir 600/100 mg twice daily arm, respectively).

### **TMC114-C214**

TMC114-C214 is a randomized, controlled, open-label Phase 3 trial comparing PREZISTA/ritonavir 600/100 mg twice daily versus lopinavir/ritonavir 400/100 mg twice daily in antiretroviral treatment-experienced, lopinavir/ritonavir-naïve HIV-1-infected adult subjects. Both arms used an optimized background regimen consisting of at least 2 antiretrovirals (NRTIs with or without NNRTIs).

HIV-1-infected subjects who were eligible for this trial had plasma HIV-1 RNA greater than 1000 copies/mL and were on a highly active antiretroviral therapy regimen (HAART) for at least 12 weeks. Virologic response was defined as a confirmed plasma HIV-1 RNA viral load less than 400 copies/mL. Analyses included 595 subjects in trial TMC114-C214 who had completed 96 weeks of treatment or discontinued earlier.

Demographics and baseline characteristics were balanced between the PREZISTA/ritonavir arm and the lopinavir/ritonavir arm (see Table 24). Table 24 compares the demographic and baseline characteristics between subjects in the PREZISTA/ritonavir 600/100 mg twice daily arm and subjects in the lopinavir/ritonavir 400/100 mg twice daily arm in trial TMC114-C214.

**Table 24: Demographic and Baseline Characteristics of Subjects in Trial TMC114-C214**

	<b>PREZISTA/ritonavir 600/100 mg twice daily + OBR N=298</b>	<b>lopinavir/ritonavir 400/100 mg twice daily + OBR N=297</b>
<b>Demographic characteristics</b>		
Median age (years) (range, years)	40 (18-68)	41 (22-76)
Sex		
Male	77%	81%
Female	23%	19%
Race		
White	54%	57%
Black	18%	17%
Hispanic	15%	15%
Asian	9%	9%
<b>Baseline characteristics</b>		
Mean baseline plasma HIV-1 RNA (log <sub>10</sub> copies/mL)	4.33	4.28
Median baseline CD4+ cell count (cells/mm <sup>3</sup> ) (range, cells/mm <sup>3</sup> )	235 (3-831)	230 (2-1096)
Percentage of patients with baseline viral load	19%	17%

≥100,000 copies/mL		
Percentage of patients with baseline CD4+ cell count <200 cells/mm <sup>3</sup>	40%	40%
Median darunavir fold change (range)	0.60 (0.10-37.40)	0.60 (0.1-43.8)
Median lopinavir fold change (range)	0.70 (0.40-74.40)	0.80 (0.30-74.50)
Median number of resistance-associated <sup>*</sup> :		
PI mutations	4	4
NNRTI mutations	1	1
NRTI mutations	2	2
Percentage of subjects with number of baseline primary protease inhibitor mutations <sup>*</sup> :		
≤1	78%	80%
2	8%	9%
≥3	13%	11%
Median number of ARVs previously used <sup>†</sup> :		
NRTIs	4	4
NNRTIs	1	1
PIs (excluding low-dose ritonavir)	1	1
Percentage of subjects resistant <sup>‡</sup> to all available <sup>§</sup> PIs at baseline, excluding darunavir	2%	3%

OBR=optimized background regimen

\* Johnson VA, Brun-Vezinet F, Clotet B, et al. Update of the drug resistance mutations in HIV-1: Fall 2006. Top HIV Med 2006; 14(3): 125-130

† Only counting ARVs, excluding low-dose ritonavir

‡ Based on phenotype (Antivirogram<sup>®</sup>)

§ Commercially available PIs at the time of trial enrollment

Week 96 outcomes for subjects on PREZISTA/ritonavir 600/100 mg twice daily from trial TMC114-C214 are shown in Table 25.

**Table 25: Virologic Outcome of Randomized Treatment of Trial TMC114-C214 at 96 Weeks**

	<b>PREZISTA /ritonavir 600/100 mg twice daily + OBR N=298</b>	<b>lopinavir/ritonavir 400/100 mg twice daily + OBR N=297</b>
Virologic success HIV-1 RNA <50 copies/mL	58%	52%
Virologic failure <sup>*</sup>	26%	33%
No virologic data at Week 96 window <sup>†</sup>		
Reasons		
Discontinued trial due to adverse event or death <sup>‡</sup>	7%	8%
Discontinued trial for other reasons <sup>§</sup>	8%	7%
Missing data during window <sup>†</sup> but on trial	1%	<1%

N = total number of subjects with data; OBR=optimized background regimen

\* Includes patients who discontinued prior to Week 96 for lack or loss of efficacy and patients who are ≥50 copies in the 96-week window and patients who had a change in their OBR that was not permitted by the protocol.

† Window 90-102 Weeks

‡ Includes patients who discontinued due to adverse event or death at any time point from Day 1 through the time window if this resulted in no virologic data on treatment during the specified window.

§ Other includes: withdrew consent, loss to follow-up, etc., if the viral load at the time of discontinuation was <50 copies/mL.

In trial TMC114-C214 at 96 weeks of treatment, the median increase from baseline in CD4+ cell counts was 81 cells/mm<sup>3</sup> in the PREZISTA/ritonavir 600/100 mg twice daily arm and 93 cells/mm<sup>3</sup> in the lopinavir/ritonavir 400/100 mg twice daily arm.

## TMC114-C213 and TMC114-C202

TMC114-C213 and TMC114-C202 are randomized, controlled, Phase 2b trials in adult subjects with a high level of PI resistance consisting of 2 parts: an initial partially-blinded, dose-finding part and a second long-term part in which all subjects randomized to PREZISTA/ritonavir received the recommended dose of 600/100 mg twice daily.

HIV-1-infected subjects who were eligible for these trials had plasma HIV-1 RNA greater than 1000 copies/mL, had prior treatment with PI(s), NNRTI(s) and NRTI(s), had at least one primary PI mutation (D30N, M46I/L, G48V, I50L/V, V82A/F/S/T, I84V, L90M) at screening, and were on a stable PI-containing regimen at screening for at least 8 weeks. Randomization was stratified by the number of PI mutations, screening viral load, and the use of enfuvirtide.

The virologic response rate was evaluated in subjects receiving PREZISTA/ritonavir plus an OBR versus a control group receiving an investigator-selected PI(s) regimen plus an OBR. Prior to randomization, PI(s) and OBR were selected by the investigator based on genotypic resistance testing and prior ARV history. The OBR consisted of at least 2 NRTIs with or without enfuvirtide. Selected PI(s) in the control arm included: lopinavir in 36%, (fos)amprenavir in 34%, saquinavir in 35% and atazanavir in 17%; 98% of control subjects received a ritonavir boosted PI regimen out of which 23% of control subjects used dual-boosted PIs. Approximately 47% of all subjects used enfuvirtide, and 35% of the use was in subjects who were ENF-naïve. Virologic response was defined as a decrease in plasma HIV-1 RNA viral load of at least 1 log<sub>10</sub> versus baseline.

In the pooled analysis for TMC114-C213 and TMC114-C202, demographics and baseline characteristics were balanced between the PREZISTA/ritonavir arm and the comparator PI arm (see Table 26). Table 26 compares the demographic and baseline characteristics between subjects in the PREZISTA/ritonavir 600/100 mg twice daily arm and subjects in the comparator PI arm in the pooled analysis of trials TMC114-C213 and TMC114-C202.

**Table 26: Demographic and Baseline Characteristics of Subjects in the Trials TMC114-C213 and TMC114-C202 (Pooled Analysis)**

	<b>PREZISTA/ritonavir 600/100 mg twice daily + OBR N=131</b>	<b>Comparator PI(s) + OBR N=124</b>
<b>Demographic characteristics</b>		
Median age (years) (range, years)	43 (27-73)	44 (25-65)
Sex		
Male	89%	88%
Female	11%	12%
Race		
White	81%	73%
Black	10%	15%
Hispanic	7%	8%
<b>Baseline characteristics</b>		
Mean baseline plasma HIV-1 RNA (log <sub>10</sub> copies/mL)	4.61	4.49
Median baseline CD4+ cell count (cells/mm <sup>3</sup> ) (range,	153 (3-776)	163 (3-1274)

cells/mm <sup>3</sup> )		
Percentage of patients with baseline viral load >100,000 copies/mL	24%	29%
Percentage of patients with baseline CD4+ cell count <200 cells/mm <sup>3</sup>	67%	58%
Median darunavir fold change	4.3	3.3
Median number of resistance-associated <sup>*</sup> :		
PI mutations	12	12
NNRTI mutations	1	1
NRTI mutations	5	5
Percentage of subjects with number of baseline primary protease inhibitor mutations <sup>†</sup> :		
≤1	8%	9%
2	22%	21%
≥3	70%	70%
Median number of ARVs previously used <sup>‡</sup> :		
NRTIs	6	6
NNRTIs	1	1
PIs (excluding low-dose ritonavir)	5	5
Percentage of subjects resistant <sup>†</sup> to all available <sup>‡</sup> PIs at baseline, excluding tipranavir and darunavir	63%	61%
Percentage of subjects with prior use of enfuvirtide	20%	17%

OBR=optimized background regimen

\* Johnson VA, Brun-Vezinet F, Clotet B, et al. Update of the drug resistance mutations in HIV-1: Fall 2006. Top HIV Med 2006; 14(3): 125-130

† Based on phenotype (Antivirogram®)

‡ Commercially available PIs at the time of trial enrollment

Week 96 outcomes for subjects on the recommended dose PREZISTA/ritonavir 600/100 mg twice daily from the pooled trials TMC114-C213 and TMC114-C202 are shown in Table 27.

**Table 27: Outcomes of Randomized Treatment Through Week 96 of the Trials TMC114-C213 and TMC114-C202 (Pooled Analysis)**

	Randomized trials TMC114-C213 and TMC114-C202	
	PREZISTA/ritonavir 600/100 mg twice daily + OBR N=131	Comparator PI(s) + OBR N=124
Virologic responders confirmed at least 1 log <sub>10</sub> HIV-1 RNA below baseline through Week 96 (<50 copies/mL at Week 96)	57% (39%)	10% (9%)
Virologic failures	29%	80%
Lack of initial response <sup>*</sup>	8%	53%
Rebounder <sup>†</sup>	17%	19%
Never suppressed <sup>‡</sup>	4%	8%
Death or discontinuation due to adverse events	9%	3%
Discontinuation due to other reasons	5%	7%

OBR=optimized background regimen

\* Subjects who did not achieve at least a confirmed 0.5 log<sub>10</sub> HIV-1 RNA drop from baseline at Week 12

† Subjects with an initial response (confirmed 1 log<sub>10</sub> drop in viral load), but without a confirmed 1 log<sub>10</sub> drop in viral load at Week 96

‡ Subjects who never reached a confirmed 1 log<sub>10</sub> drop in viral load before Week 96

In the pooled trials TMC114-C213 and TMC114-C202 through 48 weeks of treatment, the proportion of subjects with HIV-1 RNA less than 400 copies/mL in the arm receiving PREZISTA/ritonavir 600/100 mg twice daily compared to the comparator PI arm was 55.0% and 14.5%, respectively. In addition, the mean changes in plasma HIV-1 RNA from baseline were – 1.69 log<sub>10</sub> copies/mL in the arm receiving PREZISTA/ritonavir 600/100 mg twice daily and – 0.37 log<sub>10</sub> copies/mL for the comparator PI arm. The mean increase from baseline in CD4+ cell counts was higher in the arm receiving PREZISTA/ritonavir 600/100 mg twice daily (103 cells/mm<sup>3</sup>) than in the comparator PI arm (17 cells/mm<sup>3</sup>).

#### **14.4 Pediatric Patients**

The pharmacokinetic profile, safety and antiviral activity of PREZISTA/ritonavir were evaluated in 3 randomized, open-label, multicenter studies.

##### **TMC114-C212**

Treatment-experienced pediatric subjects between the ages of 6 and less than 18 years and weighing at least 20 kg were stratified according to their weight (greater than or equal to 20 kg to less than 30 kg, greater than or equal to 30 kg to less than 40 kg, greater than or equal to 40 kg) and received PREZISTA tablets with either ritonavir capsules or oral solution plus background therapy consisting of at least two non-protease inhibitor antiretroviral drugs. Eighty patients were randomized and received at least one dose of PREZISTA/ritonavir. Pediatric subjects who were at risk of discontinuing therapy due to intolerance of ritonavir oral solution (e.g., taste aversion) were allowed to switch to the capsule formulation. Of the 44 pediatric subjects taking ritonavir oral solution, 23 subjects switched to the 100 mg capsule formulation and exceeded the weight-based ritonavir dose without changes in observed safety.

The 80 randomized pediatric subjects had a median age of 14 (range 6 to less than 18 years), and were 71% male, 54% Caucasian, 30% Black, 9% Hispanic and 8% other. The mean baseline plasma HIV-1 RNA was 4.64 log<sub>10</sub> copies/mL, and the median baseline CD4+ cell count was 330 cells/mm<sup>3</sup> (range: 6 to 1505 cells/mm<sup>3</sup>). Overall, 38% of pediatric subjects had baseline plasma HIV-1 RNA ≥100,000 copies/mL. Most pediatric subjects (79%) had previous use of at least one NNRTI and 96% of pediatric subjects had previously used at least one PI.

Seventy-seven pediatric subjects (96%) completed the 24-week period. Of the patients who discontinued, one patient discontinued treatment due to an adverse event. An additional 2 patients discontinued for other reasons, one patient due to compliance and another patient due to relocation.

The proportion of pediatric subjects with HIV-1 RNA less than 400 copies/mL and less than 50 copies/mL was 64% and 50%, respectively. The mean increase in CD4+ cell count from baseline was 117 cells/mm<sup>3</sup>.

### TMC114-C228

Treatment-experienced pediatric subjects between the ages of 3 and less than 6 years and weighing greater than or equal to 10 kg to less than 20 kg received PREZISTA oral suspension with ritonavir oral solution plus background therapy consisting of at least two active non-protease inhibitor antiretroviral drugs. Twenty-one subjects received at least one dose of PREZISTA/ritonavir.

The 21 subjects had a median age of 4.4 years (range 3 to less than 6 years), and were 48% male, 57% Black, 29% Caucasian and 14% other. The mean baseline plasma HIV-1 was 4.34 log<sub>10</sub> copies/mL, the median baseline CD4+ cell count was 927 × 10<sup>6</sup> cells/L (range: 209 to 2,429 × 10<sup>6</sup> cells/L) and the median baseline CD4+ percentage was 27.7% (range: 15.6% to 51.1%). Overall, 24% of subjects had a baseline plasma HIV-1 RNA greater than or equal to 100,000 copies/mL. All subjects had used greater than or equal to 2 NRTIs, 62% of subjects had used greater than or equal to 1 NNRTI and 76% had previously used at least one HIV PI.

Twenty subjects (95%) completed the 48 week period. One subject prematurely discontinued treatment due to vomiting assessed as related to ritonavir.

The proportion of subjects with HIV-1 RNA less than 50 copies/mL at Week 48 was 71%. The mean increase in CD4+ percentage from baseline was 4%. The mean change in CD4+ cell count from baseline was 187 × 10<sup>6</sup> cells/L.

### TMC114-C230

Treatment-naïve pediatric subjects between the ages of 12 and less than 18 years and weighing at least 40 kg received the adult recommended dose of PREZISTA/ritonavir 800/100 mg once daily plus background therapy consisting of at least two non-protease inhibitor antiretroviral drugs.

The 12 randomized pediatric subjects had a median age of 14.4 years (range 12.6 to 17.3 years), and were 33.3% male, 58.3% Caucasian and 41.7% Black. The mean baseline plasma HIV-1 RNA was 4.72 log<sub>10</sub> copies/mL, and the median baseline CD4+ cell count was 282 cells/mm<sup>3</sup> (range: 204 to 515 cells/mm<sup>3</sup>). Overall, 41.7% of pediatric subjects had baseline plasma HIV-1 RNA ≥100,000 copies/mL.

All subjects completed the 48 week treatment period.

The proportion of subjects with HIV-1 RNA less than 50 copies/mL and less than 400 copies/mL was 83.3% and 91.7%, respectively. The mean increase in CD4+ cell count from baseline was 221 × 10<sup>6</sup> cells/L.

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

PREZISTA (darunavir) 100 mg per mL oral suspension is a white to off-white opaque liquid supplied in amber-colored multiple-dose bottles containing darunavir ethanolate equivalent to 100 mg of darunavir per mL packaged with a 6 mL oral dosing syringe with 0.2 mL gradations.

PREZISTA (darunavir) 75 mg tablets are supplied as white, caplet-shaped, film-coated tablets containing darunavir ethanolate equivalent to 75 mg of darunavir per tablet. Each tablet is debossed with “75” on one side and “TMC” on the other side.

PREZISTA (darunavir) 150 mg tablets are supplied as white, oval-shaped, film-coated tablets containing darunavir ethanolate equivalent to 150 mg of darunavir per tablet. Each tablet is debossed with “150” on one side and “TMC” on the other side.

PREZISTA (darunavir) 600 mg tablets are supplied as orange, oval-shaped, film-coated tablets containing darunavir ethanolate equivalent to 600 mg of darunavir per tablet. Each tablet is debossed with “600MG” on one side and “TMC” on the other side.

PREZISTA (darunavir) 800 mg tablets are supplied as dark red, oval-shaped, film-coated tablets containing darunavir ethanolate equivalent to 800 mg of darunavir per tablet. Each tablet is debossed with “800” on one side and “T” on the other side.

PREZISTA is packaged in bottles in the following configuration:

- 100 mg/mL oral suspension – 200 mL bottles (NDC 59676-565-01)
- 75 mg tablets — bottles of 480 (NDC 59676-563-01)
- 150 mg tablets — bottles of 240 (NDC 59676-564-01)
- 600 mg tablets — bottles of 60 (NDC 59676-562-01)
- 800 mg tablets — bottles of 30 (NDC 59676-566-30)

### Storage

#### *PREZISTA Oral Suspension*

- Store at 25°C (77°F); with excursions permitted to 15°-30°C (59°-86°F).
- Do not refrigerate or freeze. Avoid exposure to excessive heat.
- Store in the original container.
- Shake well before each usage.

#### *PREZISTA Tablets*

- Store at 25°C (77°F); with excursions permitted to 15°-30°C (59°-86°F).

## **17 PATIENT COUNSELING INFORMATION**

Advise the patient to read the FDA-approved patient labeling (Patient Information and Instruction for Use).

### Instructions for Use

Advise patients to take PREZISTA and ritonavir with food every day on a regular dosing schedule, as missed doses can result in development of resistance. PREZISTA must always be used with ritonavir in combination with other antiretroviral drugs. Advise patients not to alter the

dose of either PREZISTA or ritonavir, discontinue ritonavir, or discontinue therapy with PREZISTA without consulting their physician [*see Dosage and Administration (2)*].

### **Hepatotoxicity**

Inform patients that drug-induced hepatitis (e.g., acute hepatitis, cytolytic hepatitis) has been reported with PREZISTA co-administered with 100 mg of ritonavir. Advise patients about the signs and symptoms of liver problems [*see Warnings and Precautions (5.2)*].

### **Severe Skin Reactions**

Inform patients that skin reactions ranging from mild to severe, including Stevens-Johnson Syndrome, drug rash with eosinophilia and systemic symptoms, and toxic epidermal necrolysis, have been reported with PREZISTA co-administered with 100 mg of ritonavir. Advise patients to discontinue PREZISTA/ritonavir immediately if signs or symptoms of severe skin reactions develop. These can include but are not limited to severe rash or rash accompanied with fever, general malaise, fatigue, muscle or joint aches, blisters, oral lesions, conjunctivitis, hepatitis and/or eosinophilia [*see Warnings and Precautions (5.3)*].

### **Drug Interactions**

PREZISTA/ritonavir may interact with many drugs; therefore, advise patients to report to their healthcare provider the use of any other prescription or nonprescription medication or herbal products, including St. John's wort [*see Contraindications (4), Warnings and Precautions (5.4, 5.5) and Drug Interactions (7)*].

Instruct patients receiving combined hormonal contraception or the progestin only pill to use an effective alternative contraceptive method or add a barrier method during therapy with PREZISTA/ritonavir because hormonal levels may decrease [*see Drug Interactions (7.3) and Use in Specific Populations (8.3)*].

### **Fat Redistribution**

Inform patients that redistribution or accumulation of body fat may occur in patients receiving antiretroviral therapy, including PREZISTA/ritonavir, and that the cause and long-term health effects of these conditions are not known at this time [*see Warnings and Precautions (5.7)*].

### **Immune Reconstitution Syndrome**

Advise patients to inform their healthcare provider immediately of any symptoms of infection, as in some patients with advanced HIV infection (AIDS), signs and symptoms of inflammation from previous infections may occur soon after anti-HIV treatment is started [*see Warnings and Precautions (5.8)*].

### **Pregnancy Registry**

Inform patients that there is an antiretroviral pregnancy registry to monitor fetal outcomes of pregnant women exposed to PREZISTA [*see Use in Specific Populations (8.1)*].



### Lactation

Instruct women with HIV-1 infection not to breastfeed because HIV-1 can be passed to the baby in breast milk [*see Use in Specific Populations (8.2)*].

Product of Ireland

Manufactured by:  
PREZISTA oral suspension  
Janssen Pharmaceutica, NV  
Beerse, Belgium

PREZISTA tablets  
Janssen Ortho LLC,  
Gurabo,  
PR 00778

Manufactured for:  
Janssen Therapeutics,  
Division of Janssen Products, LP,  
Titusville NJ 08560

PREZISTA<sup>®</sup> is a registered trademark of Janssen Pharmaceuticals.

© 2006 Janssen Pharmaceutical Companies

PATIENT INFORMATION	
PREZISTA <sup>®</sup> (pre-ZIS-ta) (darunavir) oral suspension	PREZISTA <sup>®</sup> (pre-ZIS-ta) (darunavir) tablet
<p>Read this Patient Information before you start taking PREZISTA and each time you get a refill. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or your treatment.</p> <p>Also read the Patient Information leaflet for ritonavir.</p>	
<p><b>What is the most important information I should know about PREZISTA?</b></p> <ul style="list-style-type: none"> <li>• <b>Ask your healthcare provider or pharmacist about medicines that should not be taken with PREZISTA. For more information, see “Who should not take PREZISTA?” and “What should I tell my healthcare provider before taking PREZISTA?”</b></li> <li>• <b>PREZISTA may cause liver problems.</b> Some people taking PREZISTA in combination with ritonavir have developed liver problems, which may be life-threatening. Your healthcare provider should do blood tests before and during your PREZISTA and ritonavir combination treatment. If you have chronic hepatitis B or C infection, your healthcare provider should check your blood tests more often because you have an increased chance of developing liver problems. Tell your healthcare provider if you have any of the below signs and symptoms of liver problems. <ul style="list-style-type: none"> <li>○ dark (tea colored) urine</li> <li>○ yellowing of your skin or whites of your eyes</li> <li>○ pale colored stools (bowel movements)</li> <li>○ nausea</li> <li>○ vomiting</li> <li>○ pain or tenderness on your right side below your ribs</li> <li>○ loss of appetite</li> <li>○ tiredness</li> </ul> </li> <li>• <b>PREZISTA may cause severe or life-threatening skin reactions or rash.</b> Sometimes these skin reactions and skin rashes can become severe and require treatment in a hospital. Tell your healthcare provider right away if you develop a rash. <b>Stop</b> taking PREZISTA and ritonavir combination treatment and tell your healthcare provider right away if you have any skin changes with symptoms below: <ul style="list-style-type: none"> <li>○ fever</li> <li>○ tiredness</li> <li>○ muscle or joint pain</li> <li>○ blisters or skin lesions</li> <li>○ mouth sores or ulcers</li> <li>○ red or inflamed eyes, like “pink eye” (conjunctivitis)</li> </ul> </li> </ul> <p>Rash occurred more often in people taking PREZISTA and raltegravir together than with either drug separately, but was generally mild.</p> <p>See “<b>What are the possible side effects of PREZISTA?</b>” for more information about side effects.</p>	
<p><b>What is PREZISTA?</b></p> <p>PREZISTA is a prescription HIV-1 (Human Immunodeficiency Virus-type 1) medicine used with ritonavir and other antiretroviral medicines to treat HIV-1 infection in adults and children 3 years of age and older. HIV is the virus that causes AIDS (Acquired Immune Deficiency Syndrome).</p> <p>PREZISTA should not be used in children under 3 years of age.</p> <p>When used with other antiretroviral medicines to treat HIV-1 infection, PREZISTA may help:</p> <ul style="list-style-type: none"> <li>• reduce the amount of HIV-1 in your blood. This is called “viral load”.</li> <li>• increase the number of CD4+ (T) cells in your blood that help fight off other infections.</li> </ul> <p>Reducing the amount of HIV-1 and increasing the CD4+ (T) cells in your blood may improve your immune system. This may reduce your risk of death or getting infections that can happen when your immune system is weak (opportunistic infections).</p> <p><b>PREZISTA does not cure HIV-1 infection or AIDS.</b> You must keep taking HIV-1 medicines to control HIV-1 infection and decrease HIV-related illnesses.</p> <p><b>Avoid doing things that can spread HIV-1 infection to others:</b></p> <ul style="list-style-type: none"> <li>• Do not share or re-use needles or other injection equipment.</li> <li>• Do not share personal items that can have blood or body fluids on them, like toothbrushes and razor blades.</li> <li>• Do not have any kind of sex without protection. Always practice safe sex by using a latex or polyurethane condom to lower the chance of sexual contact with semen, vaginal secretions, or blood.</li> </ul> <p>Ask your healthcare provider if you have any questions on how to prevent passing HIV to other people.</p>	
<p><b>Who should not take PREZISTA?</b></p> <p><b>Do not take PREZISTA</b> with any of the following medicines:</p> <ul style="list-style-type: none"> <li>• alfuzosin (UROXATRAL<sup>®</sup>)</li> </ul>	

- elbasvir and grazoprevir (ZEPATIER<sup>®</sup>)
- ergot-containing medicines:
  - dihydroergotamine (D.H.E. 45<sup>®</sup>, EMBOLEX<sup>®</sup>, MIGRANAL<sup>®</sup>)
  - ergotamine tartrate (CAFERGOT<sup>®</sup>, ERGOMAR<sup>®</sup>, ERGOSTAT<sup>®</sup>, MEDIHALER ERGOTAMINE<sup>®</sup>, MIGERGOT<sup>®</sup>, WIGRAINE<sup>®</sup>, WIGRETTES<sup>®</sup>)
  - methylergonovine (ERGOTRATE<sup>®</sup>, METHERGINE<sup>®</sup>)
- cisapride (PROPULSID<sup>®</sup>)
- colchicine (COLCRYS<sup>®</sup>, MITIGARE<sup>®</sup>), if you have liver or kidney problems
- dronedarone (MULTAQ<sup>®</sup>)
- lurasidone (LATUDA<sup>®</sup>)
- pimozide (ORAP<sup>®</sup>)
- midazolam (VERSED<sup>®</sup>), when taken by mouth
- ranolazine (RANEXA<sup>®</sup>)
- St. John's wort (*Hypericum perforatum*) or a product that contains St. John's wort
- lovastatin or a product that contains lovastatin (ALTOPREV<sup>®</sup>, ADVICOR<sup>®</sup>, MEVACOR<sup>®</sup>)
- simvastatin or a product that contains simvastatin (SIMCOR<sup>®</sup>, VYTORIN<sup>®</sup>, ZOCOR<sup>®</sup>)
- rifampin or a product that contains rifampin (RIFADIN<sup>®</sup>, RIFATER<sup>®</sup>, RIFAMATE<sup>®</sup>, RIMACTANE<sup>®</sup>)
- sildenafil (REVATIO<sup>®</sup>), when used for the treatment of pulmonary arterial hypertension (PAH)
- triazolam (HALCION<sup>®</sup>)

Serious problems can happen if you or your child take any of these medicines with PREZISTA.

### What should I tell my healthcare provider before taking PREZISTA?

#### Before taking PREZISTA, tell your healthcare provider if you:

- have liver problems, including hepatitis B or hepatitis C
- are allergic to sulfa medicines
- have high blood sugar (diabetes)
- have hemophilia
- have any other medical conditions
- are pregnant or plan to become pregnant. Tell your healthcare provider if you become pregnant while taking PREZISTA.
  - **Pregnancy Registry:** There is a pregnancy registry for women who take antiretroviral medicines during pregnancy. The purpose of this registry is to collect information about the health of you and your baby. Talk to your healthcare provider about how you can take part in this registry.
- are breastfeeding or plan to breastfeed. Do not breastfeed if you take PREZISTA.
  - You should not breastfeed if you have HIV-1 because of the risk of passing HIV-1 to your baby.
  - It is not known if PREZISTA can pass into your breast milk.
  - Talk to your healthcare provider about the best way to feed your baby.

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Some medicines interact with PREZISTA. **Keep a list of your medicines to show your healthcare provider and pharmacist.**

- You can ask your healthcare provider or pharmacist for a list of medicines that interact with PREZISTA.
- **Do not start taking a new medicine without telling your healthcare provider.** Your healthcare provider can tell you if it is safe to take PREZISTA with other medicines.

### How should I take PREZISTA?

- Take PREZISTA exactly as your healthcare provider tells you.
- You must take ritonavir at the same time as PREZISTA.
- Do not change your dose or stop treatment with PREZISTA without talking to your healthcare provider.
- Take PREZISTA and ritonavir with food.
- If you have difficulty swallowing PREZISTA tablets, PREZISTA oral suspension is also available. Your healthcare provider will help decide whether PREZISTA tablets or oral suspension is right for you.
- If your child is taking PREZISTA, your child's healthcare provider will decide the right dose based on your child's weight. Your child's healthcare provider will tell you how much PREZISTA (tablets or oral suspension) and how much ritonavir (capsules, tablets or solution) your child should take. Your child should take PREZISTA with ritonavir with food. If your child does not tolerate ritonavir oral solution, ask your child's healthcare provider for advice.
- PREZISTA oral suspension should be given with the supplied oral dosing syringe. Shake the suspension well before each use. **See the "Instructions for Use" that come with PREZISTA oral suspension for information about the right way to prepare and take a dose.**
- It is important that you do not miss or skip doses of PREZISTA during treatment.

- If you take too much PREZISTA, call your healthcare provider or go to the nearest hospital emergency room right away.

### What are the possible side effects of PREZISTA?

PREZISTA may cause serious side effects, including:

- See “**What is the most important information I should know about PREZISTA?**”
- **Diabetes and high blood sugar (hyperglycemia).** Some people who take protease inhibitors including PREZISTA can get high blood sugar, develop diabetes, or your diabetes can get worse. Tell your healthcare provider if you notice an increase in thirst or urinate often while taking PREZISTA.
- **Changes in body fat** can happen in people who take HIV-1 medicines. The changes may include an increased amount of fat in the upper back and neck (“buffalo hump”), breast, and around the middle of your body (trunk). Loss of fat from the legs, arms, and face may also happen. The exact cause and long-term health effects of these conditions are not known.
- **Changes in your immune system (Immune Reconstitution Syndrome)** can happen when you start taking HIV-1 medicines. Your immune system may get stronger and begin to fight infections that have been hidden in your body for a long time. Tell your healthcare provider right away if you start having new symptoms after starting your HIV-1 medicine.
- **Increased bleeding for hemophiliacs.** Some people with hemophilia have increased bleeding with protease inhibitors including PREZISTA.

The most common side effects of PREZISTA include:

- |            |                                 |
|------------|---------------------------------|
| • diarrhea | • headache                      |
| • nausea   | • stomach-area (abdominal) pain |
| • rash     | • vomiting                      |

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all of the possible side effects of PREZISTA.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

### How should I store PREZISTA?

- Store PREZISTA oral suspension and tablets at room temperature 77°F (25°C).
- Do not refrigerate or freeze PREZISTA oral suspension.
- Keep PREZISTA oral suspension away from high heat.
- PREZISTA oral suspension should be stored in the original container.

**Keep PREZISTA and all medicines out of the reach of children.**

### General information about the safe and effective use of PREZISTA.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use PREZISTA for a condition for which it was not prescribed. Do not give PREZISTA to other people even if they have the same condition you have. It may harm them.

This leaflet summarizes the most important information about PREZISTA. If you would like more information, talk to your healthcare provider. You can ask your healthcare provider or pharmacist for information about PREZISTA that is written for health professionals. For more information, call 1-800-526-7736.

### What are the ingredients in PREZISTA?

Active ingredient: darunavir

Inactive ingredients:

**PREZISTA oral suspension:** citric acid monohydrate, hydrochloric acid (for pH adjustment), hydroxypropyl cellulose, masking flavor, methylparaben sodium, microcrystalline cellulose, purified water, sodium carboxymethylcellulose, strawberry cream flavor, and sucralose.

**PREZISTA 75 mg and 150 mg tablets:** colloidal silicon dioxide, crospovidone, magnesium stearate, microcrystalline cellulose. The film coating contains: OPADRY® White (polyethylene glycol 3350, polyvinyl alcohol-partially hydrolyzed, talc, titanium dioxide).

**PREZISTA 600 mg tablets:** colloidal silicon dioxide, crospovidone, magnesium stearate, microcrystalline cellulose. The film coating contains: OPADRY® Orange (FD&C Yellow No. 6, polyethylene glycol 3350, polyvinyl alcohol-partially hydrolyzed, talc, titanium dioxide).

**PREZISTA 800 mg tablets:** colloidal silicon dioxide, crospovidone, magnesium stearate, microcrystalline cellulose, hypromellose. The film coating contains: OPADRY® Dark Red (iron oxide red, polyethylene glycol 3350, polyvinyl alcohol-partially hydrolyzed, talc, titanium dioxide).

Product of Ireland

Manufactured by: PREZISTA oral suspension, Janssen Pharmaceutica NV, Beerse, Belgium

PREZISTA tablets, Janssen Ortho LLC, Gurabo, PR 00778

Manufactured for: Janssen Therapeutics, Division of Janssen Products, LP, Titusville NJ 08560




**INSTRUCTIONS FOR USE**  
**PREZISTA (pre-ZIS-ta)**  
**(darunavir)**  
**oral suspension**

**Be sure that you read, understand, and follow these Instructions for Use so that you measure and take PREZISTA oral suspension correctly. Ask your healthcare provider if you are not sure.**

Each PREZISTA oral suspension carton contains:

- 1 bottle of PREZISTA oral suspension
- Oral dosing syringe
- 1 oral syringe adapter



**Important information for use:**

- **Shake PREZISTA oral suspension well before each use.**
- **PREZISTA oral suspension should be given with the oral dosing syringe provided to make sure you measure the right amount. The oral dosing syringe provided with your PREZISTA oral suspension should not be used for any other medicines.**

**1. Shake the bottle.**

- Shake the bottle well before each use (See Figure A).


**Figure A: Shake the bottle**



**2. Open the bottle of PREZISTA oral suspension.**

- Open the bottle by pushing downward on the cap and twisting it in the direction of the arrow (counter-clockwise) (See Figure B).


**Figure B: Opening the bottle**



**3. The first time a bottle of PREZISTA oral suspension is used:**

- Insert the oral syringe adapter into the bottle. Press on the adapter until it is even with the top of the bottle (See Figure C).
- **Do not** remove the oral syringe adapter from the bottle once inserted.


**Figure C: Inserting the adapter**



**4. Insert the oral dosing syringe.**

- Fully push down (depress) the plunger of the syringe.
- Insert the syringe into the opening of the oral syringe adapter until it is firmly in place (See Figure D).


**Figure D: Inserting the syringe**



**5. Withdraw the prescribed amount of PREZISTA oral suspension.**


- Turn the bottle upside down. Gently pull back the plunger of the syringe until the bottom of the plunger is even with the markings on the syringe for the prescribed dose (See Figure E). If you see air bubbles in the syringe, push the plunger in to empty the oral suspension back into the bottle. Then repeat steps 4 and 5.
- If you or your child's dose of PREZISTA oral suspension is more than 6 mL, you will need to divide the dose. Follow the instructions given to you by your healthcare provider or pharmacist about how to divide the dose.

**Figure E: Withdrawing the Oral Suspension**



**6. Turn the bottle upright and remove the syringe from the bottle by pulling straight up on the oral dosing syringe (See Figure F).**

**Figure F: Removing the syringe**






**7. Take the dose of PREZISTA.**

- Place the tip of the oral dosing syringe in the mouth.
- Press on the plunger of the syringe towards the mouth (See Figure G).

If you or your child's dose is more than 6 mL you will need to divide the dose. Follow the instructions given to you by your healthcare provider or pharmacist about how to divide the dose, and repeat steps 4 through 7.


**Figure G: Taking the dose of PREZISTA**



**8. Close the bottle with the cap after use.**


- Close the bottle by twisting the cap in the direction of the arrow (clockwise) (See Figure H).

**Figure H: Closing the bottle**




- 9. Remove the plunger from the barrel by pulling the plunger and barrel away from each other (See Figure I). Rinse both parts of the syringe with water and allow to air dry after each use.**

**Figure I: Removing the plunger from the barrel**



10. After air drying, put the oral dosing syringe back together by inserting the plunger into the barrel (See Figure J). Store the oral dosing syringe with PREZISTA oral suspension.

**Figure J: Putting the syringe back together**



**How should I store PREZISTA?**

- Store PREZISTA oral suspension and the oral dosing syringe at room temperature 77°F (25°C).
- Do not refrigerate or freeze PREZISTA oral suspension.
- Keep PREZISTA oral suspension away from high heat.
- Store PREZISTA oral suspension in the original container.

**Keep PREZISTA and all medicines out of the reach of children.**

This Instruction for Use has been approved by the U.S. Food and Drug Administration.

Product of Ireland

Manufactured by:  
Janssen Pharmaceutica, N.V.  
Beerse, Belgium

Manufactured for:  
Janssen Therapeutics, Division of Janssen Products, LP, Titusville NJ 08560

Revised: June 2016  
© Janssen Pharmaceuticals, Inc. 2006