

MiraLAX
Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax

PATIENT SAMPLES NOT FOR SALE

MiraLAX
Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax

- Relieves Occasional Constipation/Irregularity
- Softens Stool

24 ONCE-DAILY DOSES



Unflavored Powder
Dissolves in ANY
Beverage!

UNFLAVORED
POWDER
GRIT FREE

24 PACKETS
NET WT 0.5 OZ
(17g) EACH

MiraLAX
Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax



Travel Size

CONTAINS 24 ONCE-DAILY DOSES
EACH PACKET CONTAINS ONE DOSE, NET WT 0.5 OZ (17g)

24
ONCE-DAILY DOSES

24 PACKETS
NET WT 0.5 OZ (17g)

OPEN HERE

PATIENT SAMPLES NOT FOR SALE

MiraLAX
Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax

Drug Facts	Drug Facts (continued)
Active ingredient (in each dose) Purpose Polyethylene Glycol 3350, 17 g..... Osmotic Laxative	If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Use ■ relieves occasional constipation (irregularity) ■ generally produces a bowel movement in 1 to 3 days	Directions ■ do not take more than directed unless advised by your doctor ■ adults and children 17 years of age and older: ■ stir and dissolve one packet of powder (17 g) in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink ■ use once a day ■ use no more than 7 days ■ children 16 years of age or under: ask a doctor
Warnings Allergy alert: Do not use if you are allergic to polyethylene glycol Do not use if you have kidney disease, except under the advice and supervision of a doctor Ask a doctor before use if you have ■ nausea, vomiting or abdominal pain ■ a sudden change in bowel habits that lasts over 2 weeks ■ irritable bowel syndrome Ask a doctor or pharmacist before use if you are taking a prescription drug	Other information ■ store at 20° - 25°C (68° - 77°F) ■ tamper-evident: do not use if foil is open or broken
When using this product you may have loose, watery, more frequent stools	Inactive ingredients none
Stop use and ask a doctor if ■ you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition. ■ you get diarrhea ■ you need to use a laxative for longer than 1 week	Questions or comments? 1-800-MiraLAX (1-800-647-2529)

Bayer, the Bayer Cross, and MiraLAX are registered trademarks of Bayer.
© 2016 Bayer.
Dist by: Bayer HealthCare LLC, Whippany, NJ 07981
XXXXX

Bayer

No Varnish
No Copy
Lot/Exp Area

PATIENT SAMPLES NOT FOR SALE

MiraLAX
Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax

• Unflavored • Sugar Free • Dissolves in Any Beverage



CONTAINS 24 ONCE-DAILY DOSES
EACH PACKET CONTAINS ONE DOSE, NET WT 0.5 OZ (17g)

Actual Size 100%

DRUG FACTS INFO.			
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	9 pt
Drug Facts	N/A	8 pt	8 pt
(continued):	N/A	8 pt	8 pt
Headings:	N/A	8 pt	8 pt
Subheadings:	N/A	6 pt	7 pt
Text:	N/A	6 pt	7 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	7.5 pt
Characters Per Inch:	N/A	No more than 39	≤ 39

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

KAREN M MAHONEY
10/04/2017