

Directions (continued)

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

BACK OF TOP PLY

DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED" OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

Treats Frequent Heartburn

ask

PREVAcid

Lansoprazole delayed-release capsules 15 mg / acid reducer

- May take 1 to 4 days for full effect
- Sodium free

14 CAPSULES

ONE 14-DAY COURSE OF TREATMENT

NOT FOR INDIVIDUAL SALE

IMPORTANT: THIS BOTTLE DOES NOT CONTAIN FULL PRODUCT INFORMATION. SEE CARTON FOR COMPLETE LABELING. READ ALL WARNINGS ON CARTON BEFORE USE. DO NOT DISCARD CARTON.

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

PEEL CORNER FOR ADDITIONAL DIRECTIONS

1.63 0.27 -1.14 L

FRONT OF TOP PLY

Directions (continued)

Store at 20-25°C (68-77°F). Keep product out of high heat and humidity.

Sodium Free

Close cap tightly after use.

Questions or comments?
1-800-462-0051

1-800-462-0051

Distributed by:
Watershed Healthcare
Warren, NJ 07059

Trademarks are owned by or licensed to the GSK group of companies.

©2016 GSK group of companies or its licensor. All rights reserved.

PREVAcid® is a registered trademark of Takeda Pharmaceutical North America, Inc., and is used under license.

163077-14L

FRONT OF BASE PLY

FOR FDA REVIEW ONLY

46165708
FPO

Drug Facts (continued)

Other information

- read the directions, warnings and package insert before use
- keep the carton and package insert.
- They contain important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture
- close cap tightly after use

Inactive ingredients colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid and ethyl acrylate copolymer, polyethylene glycol, polysorbate 80, starch, sucrose, sugar sphere, talc, titanium dioxide

Questions or comments?
1-800-452-0051

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

TAMPER-EVIDENT BOTTLE.
DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED FOR YOUR PROTECTION" OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

Treats Frequent Heartburn

NDC 0067-6286-14

PREVACID[®]

Lansoprazole delayed-release capsules 15 mg / acid reducer

• May take 1 to 4 days for full effect

• Sodium Free

14 CAPSULES
ONE 14-DAY COURSE OF TREATMENT

gsk

1 BOTTLE INSIDE

14 CAPSULES TOTAL



Actual Size

ONE 14-DAY COURSE OF TREATMENT

46165708
FPO

1 BOTTLE INSIDE

PREVACID[®]

Lansoprazole delayed-release capsules 15 mg / acid reducer

14 CAPSULES
ONE 14-DAY COURSE OF TREATMENT

<p>Drug Facts</p> <p>Active ingredient (in each capsule) Purpose Lansoprazole 15 mg Acid Reducer</p> <p>Use</p> <ul style="list-style-type: none"> • treats frequent heartburn (occurs 2 or more days a week) • not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect <p>Warnings</p> <p>Allergy alert: Do not use if you are allergic to lansoprazole</p> <p>Do not use</p> <ul style="list-style-type: none"> • if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> • liver disease • had heartburn over 3 months. This may be a sign of a more serious condition. • heartburn with lightheadedness, sweating or dizziness • chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness • frequent chest pain • frequent wheezing, particularly with heartburn • unexplained weight loss • nausea or vomiting • stomach pain <p>Ask a doctor or pharmacist before use if you are taking</p> <ul style="list-style-type: none"> • warfarin (blood-thinning medicine) • prescription antifungal or anti-yeast medicines • digoxin (heart medicine) • theophylline (asthma medicine) • tacrolimus or mycophenolate mofetil (immune system medicines) 	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> • atazanavir (medicine for HIV infection) • methotrexate (arthritis medication) <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> • your heartburn continues or worsens • you need to take this product for more than 14 days • you need to take more than 1 course of treatment every 4 months • you get diarrhea <p>If pregnant or breast-feeding, ask a health care professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> • adults 18 years of age and older • this product is to be used once a day (every 24 hours), every day for 14 days • it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours <p>14-Day Course of Treatment</p> <ul style="list-style-type: none"> • swallow 1 capsule with a glass of water before eating in the morning • take every day for 14 days • do not take more than 1 capsule a day • swallow whole. Do not crush or chew capsules. • do not use for more than 14 days unless directed by your doctor <p>Repeated 14-Day Courses (if needed)</p> <ul style="list-style-type: none"> • you may repeat a 14-day course every 4 months • do not take for more than 14 days or more often than every 4 months unless directed by a doctor <ul style="list-style-type: none"> • children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.
---	--

1-800-452-0051

Distributed by GSK Consumer Health, Warren, NJ 07059

Trade names are owned by or licensed to the GSK group of companies. ©2016 GSK group of companies or its licensor. All rights reserved. PREVACID[®] is a registered trademark of Takeda Pharmaceuticals North America, Inc. and is used under license.

1807-10


3 0067-6286-14 3

FPO

FOR FDA REVIEW ONLY



FOR FDA REVIEW ONLY





FOR FDA REVIEW ONLY

Drug Facts (continued)

Other information

- read the directions, warnings and package insert before use
- keep the carton and package insert. They contain important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture
- close cap tightly after use

Inactive ingredients colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid and ethyl acrylate copolymer, polyethylene glycol, polysorbate 80, starch, sucrose, sugar sphere, talc, titanium dioxide

Questions or comments?
1-800-452-0051

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

TAMPER-EVIDENT BOTTLE. DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED FOR YOUR PROTECTION" OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

Treats Frequent Heartburn

NDC 0067-6286-28

24HR

PREVACID

Lansoprazole delayed-release capsules 15 mg / acid reducer

- May take 1 to 4 days for full effect
- Sodium Free

28 CAPSULES

TWO 14-DAY COURSES OF TREATMENT

SAVE ON YOUR NEXT PURCHASE!
LOOK FOR YOUR COUPON INSIDE.

2 BOTTLES INSIDE

28 CAPSULES TOTAL



PREVACID 24HR
Lansoprazole delayed-release capsules 15 mg / acid reducer
Actual Size
ONE 14-DAY COURSE OF TREATMENT

PREVACID 24HR

2 BOTTLES INSIDE

28 CAPSULES

TWO 14-DAY COURSES OF TREATMENT

Drug Facts	Drug Facts (continued)
<p>Active ingredient (in each capsule) Lansoprazole 15 mg</p> <p>Purpose Acid Reducer</p> <p>Use</p> <ul style="list-style-type: none"> • treats frequent heartburn (occurs 2 or more days a week) • not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect <p>Warnings</p> <p>Allergy alert: Do not use if you are allergic to lansoprazole</p> <p>Do not use</p> <ul style="list-style-type: none"> • if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> • liver disease • had heartburn over 3 months. This may be a sign of a more serious condition. • heartburn with lightheadedness, sweating or dizziness • chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness • frequent chest pain • frequent wheezing, particularly with heartburn • unexplained weight loss • nausea or vomiting • stomach pain <p>Ask a doctor or pharmacist before use if you are taking</p> <ul style="list-style-type: none"> • warfarin (blood-thinning medicine) • prescription antifungal or anti-yeast medicines • digoxin (heart medicine) • theophylline (asthma medicine) • tacrolimus or mycophenolate mofetil (immune system medicines) 	<ul style="list-style-type: none"> • atazanavir (medicine for HIV infection) • methotrexate (arthritis medication) <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> • your heartburn continues or worsens • you need to take this product for more than 14 days • you need to take more than 1 course of treatment every 4 months • you get diarrhea <p>If pregnant or breast-feeding, ask a health care professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> • adults 18 years of age and older • this product is to be used once a day (every 24 hours), every day for 14 days • it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours <p>14-Day Course of Treatment</p> <ul style="list-style-type: none"> • swallow 1 capsule with a glass of water before eating in the morning • take every day for 14 days • do not take more than 1 capsule a day • swallow whole. Do not crush or chew capsules. <p>Repeat 14-Day Course (if needed)</p> <ul style="list-style-type: none"> • you may repeat a 14-day course every 4 months • do not take for more than 14 days or more often than every 4 months unless directed by a doctor <p>• children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p>

FOR FDA REVIEW ONLY

SAVE \$2.00

MANUFACTURER'S COUPON | EXPIRES 12/31/18

ONE (1) PREVACID 24HR 28 CAPSULES

For more information and restrictions, visit www.gsk.com. See back of coupon for details. This coupon is valid only in the United States. © 2018 GSK. All rights reserved.



Lansoprazole delayed-release capsules 15 mg / acid reducer

42165812

FPO

3 BOTTLES INSIDE
42 CAPSULES TOTAL



Drug Facts

Active ingredient (in each capsule) Purpose
Lansoprazole 15 mg Acid Reducer

Use

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to lansoprazole

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss • nausea or vomiting • stomach pain

Ask a doctor or pharmacist before use if you are taking

- warfarin (blood-thinning medicine)
- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)
- theophylline (asthma medicine)
- tacrolimus or mycophenolate mofetil (immune system medicines)

Drug Facts (continued)

• atazanavir (medicine for HIV infection) • methotrexate (arthritis medication)

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea

If pregnant or breast-feeding, ask a health care professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Drug Facts (continued)

Other information

- read the directions, warnings and package insert before use
- Keep the carton and package insert.
- They contain important information.
- store at 20-25°C (68-77°F)
- Keep product out of high heat and humidity
- protect product from moisture
- close cap tightly after use

Inactive ingredients colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid and ethyl acrylate copolymer, polyethylene glycol, polysorbate 80, starch, sucrose, sugar sphere, talc, titanium dioxide

Questions or comments?
1-800-452-0051

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

TAMPER-EVIDENT BOTTLE.
DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED FOR YOUR PROTECTION" OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

Treats Frequent Heartburn

NDC 0067-6286-42

gsk

PREVACID[®] 24HR

Lansoprazole delayed-release capsules 15 mg / acid reducer

- May take 1 to 4 days for full effect
- Sodium Free

42 CAPSULES
THREE 14-DAY COURSES OF TREATMENT

42165812

FPO

1897-EC

3 0067-6286-42 6

FPO

Distributed by: GSK Consumer Healthcare, Warren, NJ 07059
Trade mark owned by or licensed to the GSK group of companies.
©2016 GSK group of companies or its affiliate. All rights reserved.
PREVACID[®] is a registered trademark of Takeda Pharmaceutical
North America, Inc. and is used under license.

42165812

FPO

FOR FDA REVIEW ONLY



3 BOTTLES INSIDE
CAPSULES
THREE 14-DAY COURSES OF TREATMENT

Lansoprazole delayed-release capsules 15 mg / acid reducer



043



3 BOTTLES INSIDE
42 CAPSULES TOTAL

PREVACID 24HR
Lansoprazole delayed-release capsules 15 mg / acid reducer
Actual Size
ONE 14-DAY COURSE OF TREATMENT

NDC 0067-6286-29 gsk

Treats Frequent Heartburn

PREVACID 24HR

Lansoprazole delayed-release capsules 15 mg / acid reducer

- May take 1 to 4 days for full effect
- Sodium Free

42  **28**

CAPSULES

THREE 14-DAY COURSES OF TREATMENT

BONUS 14 FREE Capsules!

(42 capsules for the price of 28)

Drug Facts

Active ingredient (in each capsule)
Lansoprazole 15 mg

Purpose
Acid Reducer

Use

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to lansoprazole

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss • nausea or vomiting • stomach pain

Ask a doctor or pharmacist before use if you are taking

- warfarin (blood-thinning medicine)
- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)
- theophylline (asthma medicine)
- tacrolimus or mycophenolate mofetil (immune system medicines)

Drug Facts (continued)

- atazanavir (medicine for HIV infection)
- methotrexate (arthritis medication)

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Drug Facts (continued)

Other information

- read the directions, warnings and package insert before use
- keep the carton and package insert.
- They contain important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture

Inactive ingredients colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid and ethyl acrylate copolymer, polyethylene glycol, polysorbate 80, starch, sucrose, sugar sphere, talc, titanium dioxide

Questions or comments?
1-800-452-0061

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

TAMPER-EVIDENT BOTTLE. DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED FOR YOUR PROTECTION" OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.



FPO

1-800-452-0061

©2016 GSK group of companies or its licensee. All rights reserved. Trademarks are owned by or licensed to the GSK group of companies.

PREVACID is a registered trademark of Helsus Pharmaceuticals North America, Inc. and is used under license.

LOT: _____

EXP: _____

MADE IN: _____



3 0067-6286-29 7

FPO

16307 2014

FOR FDA REVIEW ONLY

NDC 0067-6286-42



Treats Frequent Heartburn

PREVACID[®] / 24HR

Lansoprazole delayed-release capsules 15 mg / acid reducer

- May take 1 to 4 days for full effect
- Sodium Free

3 14-DAY COURSES OF TREATMENT

42 CAPSULES

163077 FC

FOR FDA REVIEW ONLY

TAMPER-EVIDENT BOTTLE.
DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED for YOUR PROTECTION"
OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

Drug Facts	Purpose
Active ingredient (in each capsule) Lansoprazole 15 mg	Acid Reducer
Use • treats frequent heartburn (occurs 2 or more days a week) • not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect	
Warnings Allergy alert: Do not use if you are allergic to lansoprazole Do not use • if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. Ask a doctor before use if you have • liver disease • had heartburn over 3 months. This may be a sign of a more serious condition. • heartburn with lightheadedness, sweating or dizziness • chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness • frequent chest pain • frequent wheezing, particularly with heartburn • unexplained weight loss • nausea or vomiting • stomach pain Ask a doctor or pharmacist before use if you are taking • warfarin (blood-thinning medicine) • prescription antifungal or anti-yeast medicines • digoxin (heart medicine) • theophylline (asthma medicine) • tacrolimus or mycophenolate mofetil (immune system medicines) • atazanavir (medicine for HIV infection) • methotrexate (arthritis medication) Stop use and ask a doctor if • your heartburn continues or worsens • you need to take this product for more than 14 days • you need to take more than 1 course of treatment every 4 months • you get diarrhea If pregnant or breast-feeding , ask a health care professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	

Drug Facts (continued)
Directions • adults 18 years of age and older • this product is to be used once a day (every 24 hours), every day for 14 days • it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours 14-Day Course of Treatment • swallow 1 capsule with a glass of water before eating in the morning • take every day for 14 days • do not take more than 1 capsule a day • swallow whole. Do not crush or chew capsules. • do not use for more than 14 days unless directed by your doctor Repeated 14-Day Courses (if needed) • you may repeat a 14-day course every 4 months • do not take for more than 14 days or more often than every 4 months unless directed by a doctor • children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.
Other information • read the directions, warnings and all information before use • keep the package. It contains important information. • store at 20-25°C (68-77°F) • keep product out of high heat and humidity • protect product from moisture • close cap tightly after use
Inactive ingredients colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid and ethyl acrylate copolymer, polyethylene glycol, polysorbate 80, starch, sucrose, sugar sphere, talc, titanium dioxide
Questions or comments? 1-800-452-0051



What you should know about PREVACID®24HR
Please read all information before taking PREVACID®24HR.
Save for future reference.

How PREVACID®24HR Treats Your Frequent Heartburn
PREVACID®24HR stops acid production at the source – the pumps that release acid into the stomach. PREVACID®24HR is taken once a day (every 24 hours), every day for 14 days.

What You Can Expect When Taking PREVACID®24HR
Frequent heartburn can occur anytime during the 24-hour period (day or night). Take PREVACID®24HR in the morning before eating. PREVACID®24HR is clinically proven to treat frequent heartburn. Although some people get complete relief of symptoms within 24 hours, it may take 1 to 4 days for full effect. Make sure you take PREVACID®24HR every day for 14 days to treat your frequent heartburn.

Who Should Take PREVACID®24HR
Adults (18 years and older) with frequent heartburn – when you have heartburn 2 or more days a week.

Who Should NOT Take PREVACID®24HR
People who have one episode of heartburn a week or less, or who want immediate relief of heartburn.

How to Take PREVACID®24HR
14-DAY Course of Treatment
• Swallow 1 capsule with a glass of water before eating in the morning.
• Take every day for 14 days.
• Do not take more than 1 capsule a day.
• Swallow whole. Do not crush or chew capsules.
• Do not use for more than 14 days unless directed by your doctor.

When to Take PREVACID®24HR Again
You may repeat a 14-day course of therapy every 4 months.
When to Talk to Your Doctor
Do not take for more than 14 days or more often than every 4 months unless directed by a doctor.

Warnings and When to Ask Your Doctor
Allergy alert: Do not use if you are allergic to lansoprazole
Do not use
• if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have
• liver disease
• had heartburn over 3 months. This may be a sign of a more serious condition.
• heartburn with lightheadedness, sweating or dizziness
• chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
• frequent chest pain
• frequent wheezing, particularly with heartburn
• unexplained weight loss
• nausea or vomiting
• stomach pain

Ask a doctor or pharmacist before use if you are taking
• warfarin (blood-thinning medicine)
• prescription antifungal or anti-yeast medicines
• digoxin (heart medicine)
• theophylline (asthma medicine)
• tacrolimus or mycophenolate mofetil (immune system medicines)
• atazanavir (medicine for HIV infection)
• methotrexate (arthritis medication)

Stop use and ask a doctor if
• your heartburn continues or worsens
• you need to take this product for more than 14 days
• you need to take more than 1 course of treatment every 4 months
• you get diarrhea

If pregnant or breast-feeding, ask a health care professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Tips for Managing Heartburn
• Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
• Eat slowly and do not eat big meals.
• Do not eat late at night or just before bedtime.
• Do not lie flat or bend over soon after eating.
• Raise the head of your bed.
• Wear loose-fitting clothing around your stomach.
• If you are overweight, lose weight.
• If you smoke, quit smoking.

Clinical studies prove PREVACID®24HR effectively treats frequent heartburn
In three clinical studies, PREVACID®24HR was shown to be significantly better than placebo in treating frequent heartburn.
How PREVACID®24HR is Sold
PREVACID®24HR is available in 14 capsule, 28 capsule and 42 capsule sizes. These sizes contain one, two and three 14-day courses of treatment, respectively. Do not use for more than 14 days in a row unless directed by your doctor. For the 28 count (two 14-day courses) and the 42 count (three 14-day courses), you may repeat a 14-day course every 4 months.

For Questions or Comments About PREVACID®24HR
Call 1-800-452-0051

LOT:
EXP:
MADE IN:

1-800-452-0051

Distributed by: GSK Consumer Healthcare, Warren, NJ 07059
Trademarks are owned by or licensed to the GSK group of companies. ©2016 GSK group of companies or its licensor. All rights reserved.
PREVACID® is a registered trademark of Takeda Pharmaceuticals North America, Inc., and is used under license.

163077-8C

FOR FDA REVIEW ONLY

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

THERESA M MICHELE
02/24/2017