



DRUG FACT FORMATTING: STANDARD		
DRUG FACTS TEXT	TYPE SIZE	TYPE STYLE
DRUG FACTS TITLE	9 pt.	Helvetica Cond Bold Obl
DRUG FACTS CONT.	8 pt.	Helvetica Cond Bold Obl
HEADINGS	8 pt.	Helvetica Cond Bold Obl
SUBHEADS	6 pt.	Helvetica Cond Bold
BODY TEXT	6 pt.	Helvetica Cond
TRACKING	0	
CONDENSATION	85%	
LEADING	6.5 pt.	
BARLINES, HAIRLINES	1.5, 0.5 pt.	
CHARACTERS/INCH	<39	
BULLETS	5 pt.	

Drug Facts (continued)

Other information

- read the directions, warnings and package insert before use
- keep the carton and package insert. They contain important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture
- close cap tightly after use

Inactive ingredients colloidal silicon dioxide, D&C red No. 28, F35C blue No. 1, F35G green No. 3, F35G red No. 40, gelatin, hydroxypropyl cellulose, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid and ethyl acrylate copolymer, polyethylene glycol, polysorbate 80, starch, sucrose, sugar sphere, talc, titanium dioxide

Questions or comments?
1-800-452-0061

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

TAMPER-EVIDENT BOTTLE.
DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED FOR YOUR PROTECTION" OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

Treats Frequent Heartburn

NDC 0067-6286-14

24HR

PREVACID

Lansoprazole delayed-release capsules 15 mg / acid reducer

• May take 1 to 4 days for full effect

• Sodium Free

14 CAPSULES
ONE 14-DAY COURSE OF TREATMENT

SAVE ON YOUR NEXT PURCHASE!
LOOK FOR YOUR COUPON INSIDE.

1 BOTTLE INSIDE
14 CAPSULES TOTAL

PREVACID
Lansoprazole delayed-release capsules 15 mg / acid reducer
Actual Size
ONE 14-DAY COURSE OF TREATMENT

Drug Facts

Active ingredient (in each capsule) Purpose
Lansoprazole 15 mg Acid Reducer

Use

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to lansoprazole

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- liver disease
- heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking

- warfarin (blood-thinning medicine)
- prescription antifungal or anti-yeast medicines
- disipon (heart medicine)
- theophylline (asthma medicine)
- tacrolimus or mycophenolate mofetil (immune system medicines)

CODE 128 FPO

169961466

DRUG FACT FORMATTING- STANDARD		
DRUG FACTS TEXT	TYPE SIZE	TYPE STYLE
DRUG FACTS TITLE	9 pt.	Helvetica Cond Bold Out
DRUG FACTS CONT.	8 pt.	Helvetica Cond Bold Out
HEADINGS	8 pt.	Helvetica Cond Bold Out
SUBHEADS	6 pt.	Helvetica Cond Bold
BODY TEXT	6 pt.	Helvetica Cond
TRACKING	0	
CONDENSATION	85%	
LEADING	6.5 pt.	
BARLINES, HORIZONTALS	1.5, 0.5 pt.	
CHARACTERS/INCH	<39	
BULLETS	5 pt.	

Drug Facts (continued)

Other information

- read the directions, warnings and package insert before use
- keep the carton and package insert. They contain important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture
- close cap tightly after use

Inactive ingredients colloidal silicon dioxide, D&C red No. 28, F35C blue No. 1, F35G green No. 3, F35G red No. 40, gelatin, hydroxypropyl cellulose, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid and ethyl acrylate copolymer, polyethylene glycol, polysorbate 80, starch, sucrose, sugar sphere, talc, titanium dioxide

Questions or comments?
1-800-452-0061

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

TAMPER-EVIDENT BOTTLE.
DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED FOR YOUR PROTECTION" OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

Treats Frequent Heartburn

NDC 0067-6286-14

24HR

PREVACID

Lansoprazole delayed-release capsules 15 mg / acid reducer

• May take 1 to 4 days for full effect

• Sodium Free

14 CAPSULES
ONE 14-DAY COURSE OF TREATMENT

SAVE \$1.50 on any one (1)
MANUFACTURER'S COUPON | EXPRES XXXXXX

1 BOTTLE INSIDE
14 CAPSULES TOTAL

PREVACID
Lansoprazole delayed-release capsules 15 mg / acid reducer
Actual Size
ONE 14-DAY COURSE OF TREATMENT

Drug Facts

Active ingredient (in each capsule) Purpose
Lansoprazole 15 mg Acid Reducer

Use

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to lansoprazole

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- liver disease
- heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking

- warfarin (blood-thinning medicine)
- prescription antifungal or anti-yeast medicines
- disipon (heart medicine)
- theophylline (asthma medicine)
- tacrolimus or mycophenolate mofetil (immune system medicines)

CODE 128 FPO

169961466



DRUG FACT FORMATTING: STANDARD		
DRUG FACTS TEXT	TYPE SIZE	TYPE STYLE
DRUG FACTS TITLE	9 pt.	Helvetica Cond Bold Obl
DRUG FACTS CONT.	8 pt.	Helvetica Cond Bold Obl
HEADINGS	8 pt.	Helvetica Cond Bold Obl
SUBHEADS	6 pt.	Helvetica Cond Bold
BODY TEXT	6 pt.	Helvetica Cond
TRACKING	0	
CONDENSATION	85%	
LEADING	6.5 pt.	
BARLINES, HAIRLINES	1.5, 0.5 pt.	
CHARACTERS/INCH	<39	
BULLETS	5 pt.	

Drug Facts (continued)

Other information

- read the directions, warnings and package insert before use
- keep the carton and package insert. They contain important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture
- close cap tightly after use

Inactive ingredients colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid and ethyl acrylate copolymer, polyethylene glycol, polyisobutyl 80, starch, sucrose, sugar sphere, talc, titanium dioxide

Questions or comments?
1-800-452-0051

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

TAMPER-EVIDENT BOTTLE. DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED FOR YOUR PROTECTION" OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

Treats Frequent Heartburn

NDC 0067-6286-28

24HR

PREVACID

Lansoprazole delayed-release capsules 15 mg / acid reducer

- May take 1 to 4 days for full effect
- Sodium Free

28 CAPSULES

TWO 14-DAY COURSES OF TREATMENT

SAVE ON YOUR NEXT PURCHASE!
LOOK FOR YOUR COUPON INSIDE.

2 BOTTLES INSIDE

28 CAPSULES TOTAL



PREVACID 24HR
Lansoprazole delayed-release capsules 15 mg / acid reducer
Actual Size
14-DAY COURSE OF TREATMENT

PREVACID 24HR
Lansoprazole delayed-release capsules 15 mg / acid reducer
28 CAPSULES
TWO 14-DAY COURSES OF TREATMENT

CODE 128 FP0

168906286C

CODE 128 FP0

168906286C

1-800-452-0051

Distributed by: GSK Consumer Healthcare, Warren, NJ 07059

Parameters are owned by or licensed to the GSK group of companies.

PREVACID® is a registered trademark of Helsus Pharmaceuticals.

Lansoprazole is a registered trademark of Helsus Pharmaceuticals.

©2015 Helsus Pharmaceuticals, Inc. and is used under license.

DRUG FACT FORMATTING: STANDARD

DRUG FACTS TEXT	TYPE SIZE	TYPE STYLE
DRUG FACTS TITLE	9 pt.	Helvetica Cond Bold Obli
DRUG FACTS CONT.	8 pt.	Helvetica Cond Bold Obli
HEADINGS	8 pt.	Helvetica Cond Bold Obli
SUBHEADS	6 pt.	Helvetica Cond Bold
BODY TEXT	6 pt.	Helvetica Cond
TRACKING	0	
CONDENSATION	85%	
LEADING	6.5 pt.	
BAR LINES, HAR LINES	1.5, 0.5 pt.	
CHARACTERS/INCH	<=39	
BULLETS	5 pt.	

Drug Facts (continued)

Other information

- read the directions, warnings and package insert before use
- keep the carton and package insert. They contain important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture
- close cap tightly after use

Inactive ingredients colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid and ethyl acrylate copolymer, polyethylene glycol, polyisobutyl 80, starch, sucrose, sugar sphere, talc, titanium dioxide

Questions or comments?
1-800-452-0051

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

TAMPER-EVIDENT BOTTLE. DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED FOR YOUR PROTECTION" OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

Treats Frequent Heartburn

NDC 0067-6286-28

24HR

PREVACID

Lansoprazole delayed-release capsules 15 mg / acid reducer

- May take 1 to 4 days for full effect
- Sodium Free

28 CAPSULES

TWO 14-DAY COURSES OF TREATMENT

SAVE ON YOUR NEXT PURCHASE!
LOOK FOR YOUR COUPON INSIDE.

2 BOTTLES INSIDE

28 CAPSULES TOTAL



PREVACID 24HR
Lansoprazole delayed-release capsules 15 mg / acid reducer
Actual Size
14-DAY COURSE OF TREATMENT

PREVACID 24HR
Lansoprazole delayed-release capsules 15 mg / acid reducer
28 CAPSULES
TWO 14-DAY COURSES OF TREATMENT

CODE 128 FP0

168906286C

CODE 128 FP0

168906286C

1-800-452-0051

Distributed by: GSK Consumer Healthcare, Warren, NJ 07059

Parameters are owned by or licensed to the GSK group of companies.

PREVACID® is a registered trademark of Helsus Pharmaceuticals.

Lansoprazole is a registered trademark of Helsus Pharmaceuticals.

©2015 Helsus Pharmaceuticals, Inc. and is used under license.



SAVE \$2.00

 ONE (1)

 PREVACID 24HR

 28 CAPSULES

 090067-105113

MANUFACTURER'S COUPON | EXPIRES XXXXXX



3 BOTTLES INSIDE
CAPSULES
OF TREATMENT

Drug Facts

Active ingredient (in each capsule) Purpose
Lansoprazole 15 mg Acid Reducer

Use

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to lansoprazole

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss • nausea or vomiting • stomach pain

Ask a doctor or pharmacist before use if you are taking

- warfarin (blood-thinning medicine)
- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)
- theophylline (asthma medicine)
- tacrolimus or mycophenolate mofetil (immune system medicines)

Drug Facts (continued)

- atazanavir (medicine for HIV infection) • methotrexate (arthritis medication)

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea • you develop a rash or joint pain

if pregnant or breast-feeding, ask a health care professional before use. Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Drug Facts (continued)

Other information

- read the directions, warnings and package insert before use
- keep the carton and package insert. They contain important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture
- close cap tightly after use

Inactive ingredients colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid and ethyl acrylate copolymer, polyethylene glycol, polysorbate 80, starch, sucrose, sugar sphere, talc, titanium dioxide

Questions or comments?
1-800-452-0051

KEEP THE CARTON AND PACKAGE INSERT.
THEY CONTAIN IMPORTANT INFORMATION.

TAMPER-EVIDENT BOTTLE.
DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED FOR YOUR PROTECTION" OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

Treats Frequent Heartburn

NDC 0067-6286-42



PREVACID[®]

Lansoprazole delayed-release capsules 15 mg / acid reducer

- May take 1 to 4 days for full effect
- Sodium Free

42 CAPSULES

THREE 14-DAY COURSES OF TREATMENT



189906-426

3 0067-6286-426

FPO

North America, Inc., and is used under license.
PREVACID[®] is a registered trademark of Takeda Pharmaceuticals
©2016 GSK group of companies or its licensor. All rights reserved.
Trademarks are owned by or licensed to the GSK group of companies.
Distributed by GSK Consumer Healthcare, Warren, NJ 07059
1-800-452-0051

468986426

CODE 128 FPO

CODE 128 FPO

468986426

DRUG FACT FORMATTING: STANDARD		
DRUG FACTS TEXT	TYPE SIZE	TYPE STYLE
DRUG FACTS TITLE	9 pt.	Helvetica Cond Bold Obl
DRUG FACTS CONT.	8 pt.	Helvetica Cond Bold Obl
HEADINGS	8 pt.	Helvetica Cond Bold Obl
SUBHEADS	6 pt.	Helvetica Cond Bold
BODY TEXT	6 pt.	Helvetica Cond
TRACKING	0	
CONDENSATION	85%	
LEADING	6.5 pt.	
BARLINES, HAIRLINES	1.5, 0.5 pt.	
CHARACTERS/INCH	<39	
BULLETS	5 pt.	



Lansoprazole delayed-release capsules 15 mg / acid reducer
CAPSULES
THREE 14-DAY COURSES
OF TREATMENT
3 BOTTLES INSIDE

1-899-062814

CODE 128 FP0

1-899-062814

CODE 128 FP0

NDC 0067-6286-29 gsk

Treats Frequent Heartburn

PREVACID[®] 24HR

Lansoprazole delayed-release capsules 15 mg / acid reducer

- May take 1 to 4 days for full effect
- Sodium Free

42

28

CAPSULES

THREE 14-DAY COURSES OF TREATMENT

BONUS 14 FREE Capsules!

(42 capsules for the price of 28)



3 BOTTLES INSIDE

42 CAPSULES TOTAL

ONE 14-DAY COURSE OF TREATMENT

168906-28-14

7 0067-6286-29 3

FP0

1-800-452-0051

Distributed by: GSK Consumer Healthcare, Warren, NJ 07059
©2016 GSK group of companies or its licensee. All rights reserved.
PREVACID[®] is a registered trademark of Takeda Pharmaceuticals North America, Inc., and is used under license.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

TAMPER-EVIDENT BOTTLE. DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED FOR YOUR PROTECTION" OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

168906-2814

CODE 128 FP0

Drug Facts

Active ingredient (in each capsule)
Lansoprazole 15 mg

Purpose
Acid Reducer

Use

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to lansoprazole

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking

- warfarin (blood-thinning medicine)
- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)
- theophylline (asthma medicine)
- tacrolimus or mycophenolate mofetil (immune system medicines)

Drug Facts (continued)

- atazanavir (medicine for HIV infection)
- methotrexate (arthritis medication)

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

if pregnant or breast-feeding, ask a health care professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Drug Facts (continued)

Other information

- read the directions, warnings and package insert before use
- keep the carton and package insert.
- They contain important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture
- close cap tightly after use

Inactive ingredients colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid and ethyl acrylate copolymer, polyethylene glycol, polysorbate 80, starch, sucrose, sugar sphere, talc, titanium dioxide

Questions or comments?
1-800-452-0051

DRUG FACT FORMATTING: STANDARD

DRUG FACTS TEXT	TYPE SIZE	TYPE STYLE
DRUG FACTS TITLE	9 pt.	Helvetica Cond Bold Obl
DRUG FACTS CONT.	8 pt.	Helvetica Cond Bold Obl
HEADINGS	8 pt.	Helvetica Cond Bold Obl
SUBHEADS	6 pt.	Helvetica Cond Bold
BODY TEXT	6 pt.	Helvetica Cond
TRACKING	0	
CONDENSATION	85%	
LEADING	6.5 pt.	
BARLINES, HAIRLINES	1.5, 0.5 pt.	
CHARACTERS/INCH	<39	
BULLETS	5 pt.	



PREVACID[®] 24HR
Lansoprazole delayed-release capsules 15 mg / acid reducer

3 BOTTLES INSIDE
42 CAPSULES
THREE 14-DAY COURSES OF TREATMENT

1-800-664-2262

CODE 128 FP0



NDC 0067-6286-42

Drug Facts

Active ingredient (in each capsule) **Purpose**
Lansoprazole 15 mg Acid Reducer

Use
• treats frequent heartburn (occurs 2 or more days a week)
• not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to lansoprazole

Do not use
• if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have
• liver disease
• had heartburn over 3 months. This may be a sign of a more serious condition.
• heartburn with lightheadedness, sweating or dizziness
• chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness
• frequent chest pain
• frequent wheezing, particularly with heartburn
• unexplained weight loss • nausea or vomiting • stomach pain

Ask a doctor or pharmacist before use if you are taking
• warfarin (blood-thinning medicine)
• prescription antifungal or anti-yeast medicines
• digoxin (heart medicine)
• theophylline (asthma medicine)
• tacrolimus or mycophenolate mofetil (immune system medicines)

Drug Facts (continued)

• atazanavir (medicine for HIV infection) • methotrexate (arthritis medication)

Stop use and ask a doctor if
• your heartburn continues or worsens
• you need to take this product for more than 14 days
• you need to take more than 1 course of treatment every 4 months
• you get diarrhea • you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use. Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
• adults 18 years of age and older
• this product is to be used once a day (every 24 hours), every day for 14 days
• it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
• swallow 1 capsule with a glass of water before eating in the morning
• take every day for 14 days
• do not take more than 1 capsule a day
• swallow whole. Do not crush or chew capsules.
• do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)
• you may repeat a 14-day course every 4 months
• do not take for more than 14 days or more often than every 4 months unless directed by a doctor

• children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Drug Facts (continued)

Other information
• read the directions, warnings and package insert before use
• keep the carton and package insert. They contain important information.
• store at 20-25°C (68-77°F)
• keep product out of high heat and humidity
• protect product from moisture
• close cap tightly after use

Inactive ingredients colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, magnesium carbonate, methacrylic acid and ethyl acrylate copolymer, polyethylene glycol, polyborate 80, starch, sucrose, sugar sphere, talc, titanium dioxide

Questions or comments?
1-800-452-0051

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

TAMPER-EVIDENT BOTTLE.
DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED FOR YOUR PROTECTION" OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

Treats Frequent Heartburn

24HR

PREVACID[®]

Lansoprazole delayed-release capsules 15 mg / acid reducer

• May take 1 to 4 days for full effect
• Sodium Free

SAVE ON YOUR NEXT PURCHASE! LOOK FOR YOUR COUPON INSIDE!

42 CAPSULES

THREE 14-DAY COURSES OF TREATMENT

1-800-664-2262

CODE 128 FP0



3 BOTTLES INSIDE
42 CAPSULES TOTAL

PREVACID[®] 24HR
Lansoprazole delayed-release capsules 15 mg / acid reducer
Actual Size
ONE 14-DAY COURSE OF TREATMENT


1-800-664-2262

CODE 128 FP0

DRUG FACT FORMATTING: STANDARD		
DRUG FACTS TEXT	TYPE SIZE	TYPE STYLE
DRUG FACTS TITLE	9 pt.	Helvetica Cond Bold Obl
DRUG FACTS CONT.	8 pt.	Helvetica Cond Bold Obl
HEADINGS	8 pt.	Helvetica Cond Bold Obl
SUBHEADS	6 pt.	Helvetica Cond Bold
BODY TEXT	6 pt.	Helvetica Cond
TRACKING	0	
CONDENSATION	85%	
LEADING	6.5 pt.	
BARLINES, HARLINES	1.5, 0.5 pt.	
CHARACTERS/INCH	<39	
BULLETS	5 pt.	

1-800-664-2262

CODE 128 FP0



3 BOTTLES INSIDE
42 CAPSULES TOTAL

PREVACID[®] 24HR
Lansoprazole delayed-release capsules 15 mg / acid reducer
Actual Size
ONE 14-DAY COURSE OF TREATMENT

1-800-664-2262

CODE 128 FP0

Treats Frequent Heartburn

24HR

PREVACID[®]

Lansoprazole delayed-release capsules 15 mg / acid reducer

• May take 1 to 4 days for full effect
• Sodium Free

SAVE ON YOUR NEXT PURCHASE! LOOK FOR YOUR COUPON INSIDE!

42 CAPSULES

THREE 14-DAY COURSES OF TREATMENT

1-800-664-2262

CODE 128 FP0



3 BOTTLES INSIDE
42 CAPSULES TOTAL

PREVACID[®] 24HR
Lansoprazole delayed-release capsules 15 mg / acid reducer
Actual Size
ONE 14-DAY COURSE OF TREATMENT

1-800-664-2262

CODE 128 FP0



SAVE \$3.00 ON ONE (1) PREVACID[®] 24HR 42 ct.

MANUFACTURER'S COUPON | EXP. RES. XXXXXX
 Redeemable at retail drug and mass merchandise stores. COMPLETES 1st coupon purchase of indicated product. You pay any sales tax. 7874263 528 with numbers you take value of this coupon plus 124 ending five digits. Valid in US only. Not valid in Canada, Mexico, Puerto Rico, Alaska, Hawaii, or other territories. Void where prohibited. ©2015 GSK. All rights reserved.

NDC 0067-6286-42



Treats Frequent Heartburn

PREVACID[®] / 24HR

Lansoprazole delayed-release capsules 15 mg / acid reducer

- May take 1 to 4 days for full effect
- Sodium Free

3 14-DAY COURSES OF TREATMENT

42 CAPSULES

163077-FC

TAMPER-EVIDENT BOTTLE.
DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED for YOUR PROTECTION"
OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

Drug Facts

Active ingredient (in each capsule)
 Lansoprazole 15 mg

Purpose
 Acid Reducer

Use

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to lansoprazole

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
 - frequent wheezing, particularly with heartburn
- unexplained weight loss
 - nausea or vomiting
 - stomach pain

Ask a doctor or pharmacist before use if you are taking

- warfarin (blood-thinning medicine)
 - prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)
 - theophylline (asthma medicine)
- tacrolimus or mycophenolate mofetil (immune system medicines)
- atazanavir (medicine for HIV infection)
 - methotrexate (arthritis medication)

Stop use and ask a doctor if

- your heartburn continues or worsens
 - you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
 - you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.

Keep out of reach of children.
 In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor**

Other information

- read the directions, warnings and package insert before use
- keep the package. It contains important information.
- store at 20-25°C (68-77°F)
 - keep product out of high heat and humidity
- protect product from moisture
 - close cap tightly after use

Inactive ingredients colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid and ethyl acrylate copolymer, polyethylene glycol, polysorbate 80, starch, sucrose, sugar sphere, talc, titanium dioxide

Questions or comments? 1-800-452-0051



What you should know about PREVACID®24HR
 Please read all information before taking PREVACID®24HR.
 Save for future reference.

How PREVACID®24HR Treats Your Frequent Heartburn

PREVACID®24HR stops acid production at the source – the pumps that release acid into the stomach. PREVACID®24HR is taken once a day (every 24 hours), every day for 14 days.

What You Can Expect When Taking PREVACID®24HR

Frequent heartburn can occur anytime during the 24-hour period (day or night). Take PREVACID®24HR in the morning before eating. PREVACID®24HR is clinically proven to treat frequent heartburn. Although some people get complete relief of symptoms within 24 hours, it may take 1 to 4 days for full effect. Make sure you take PREVACID®24HR every day for 14 days to treat your frequent heartburn.

Who Should Take PREVACID®24HR

Adults (18 years and older) with frequent heartburn – when you have heartburn 2 or more days a week.

Who Should NOT Take PREVACID®24HR

People who have one episode of heartburn a week or less, or who want immediate relief of heartburn.

How to Take PREVACID®24HR

14-DAY Course of Treatment

- Swallow 1 capsule with a glass of water before eating in the morning.
- Take every day for 14 days.
- Do not take more than 1 capsule a day.
- Swallow whole. Do not crush or chew capsules.
- Do not use for more than 14 days unless directed by your doctor.

When to Take PREVACID®24HR Again

You may repeat a 14-day course of therapy every 4 months.

When to Talk to Your Doctor

Do not take for more than 14 days or more often than every 4 months unless directed by a doctor.

Warnings and When to Ask Your Doctor

Allergy alert: Do not use if you are allergic to lansoprazole

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
 - frequent wheezing, particularly with heartburn
- unexplained weight loss
 - nausea or vomiting
 - stomach pain

Ask a doctor or pharmacist before use if you are taking

- warfarin (blood-thinning medicine)
 - prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)
 - theophylline (asthma medicine)
- tacrolimus or mycophenolate mofetil (immune system medicines)
- atazanavir (medicine for HIV infection)
 - methotrexate (arthritis medication)

Stop use and ask a doctor if

- your heartburn continues or worsens
 - you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
 - you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

Clinical studies prove PREVACID®24HR effectively treats frequent heartburn

In three clinical studies, PREVACID®24HR was shown to be significantly better than placebo in treating frequent heartburn.

How PREVACID®24HR is Sold

PREVACID®24HR is available in 14 capsule, 28 capsule and 42 capsule sizes. These sizes contain one, two and three 14-day courses of treatment, respectively. Do not use for more than 14 days in a row unless directed by your doctor. For the 28 count (two 14-day courses) and the 42 count (three 14-day courses), you may repeat a 14-day course every 4 months.

For Questions or Comments About PREVACID®24HR
 Call 1-800-452-0051

LOT:
 EXP:
 MADE IN:

1-800-452-0051

Distributed by: GSK Consumer Healthcare, Warren, NJ 07059
 Trademarks are owned by or licensed to the GSK group of companies. ©2016 GSK group of companies or its licensor. All rights reserved.
 PREVACID® is a registered trademark of Takeda Pharmaceuticals North America, Inc., and is used under license.

168906-BC

DRUG FACT FORMATTING: STANDARD		
DRUG FACTS TEXT	TYPE SIZE	TYPE STYLE
DRUG FACTS TITLE	9 pt.	Helvetica Cond Bold Obl
DRUG FACTS CONT.	8 pt.	Helvetica Cond Bold Obl
HEADINGS	8 pt.	Helvetica Cond Bold Obl
SUBHEADS	6 pt.	Helvetica Cond Bold
BODY TEXT	6 pt.	Helvetica Cond
TRACKING	0	
CONDENSATION	85%	
LEADING	6.5 pt.	
BARLINES, HAIRLINES	1.5, 0.5 pt.	
CHARACTERS/INCH	<39	
BULLETS	5 pt.	

Treats Frequent Heartburn

PREVACID[®] 24HR

Lansoprazole delayed-release capsules 15 mg / acid reducer

- May take 1 to 4 days for full effect
- Sodium Free



QUIET ZONE

Please read the entire package insert before taking PREVACID[®]24HR. Save for future reference.

How PREVACID[®]24HR Treats Your Frequent Heartburn

PREVACID[®]24HR stops acid production at the source – the **pumps** that release acid into the stomach. PREVACID[®]24HR is taken once a day (every 24 hours), every day for 14 days.

What You Can Expect When Taking PREVACID[®]24HR

Frequent heartburn can occur anytime during the 24-hour period (day or night). Take PREVACID[®]24HR in the morning before eating. PREVACID[®]24HR is clinically proven to treat frequent heartburn. Although some people get complete relief of symptoms within 24 hours, it may take 1 to 4 days for full effect. Make sure you take PREVACID[®]24HR every day for 14 days to treat your frequent heartburn.

Who Should Take PREVACID[®]24HR

Adults (18 years and older) with **frequent heartburn** – when you have heartburn 2 or more days a week.

Who Should NOT Take PREVACID[®]24HR

People who have one episode of heartburn a week or less, or who want immediate relief of heartburn.

How to Take PREVACID[®]24HR

14-DAY Course of Treatment

- Swallow 1 capsule with a glass of water before eating in the morning.
- Take every day for 14 days.
- Do not take more than 1 capsule a day.
- Swallow whole. Do not crush or chew capsules.
- Do not use for more than 14 days unless directed by your doctor.

When to Take PREVACID[®]24HR Again

You may repeat a 14-day course of therapy every 4 months.

When to Talk to Your Doctor

Do not take for more than 14 days or more often than every 4 months unless directed by a doctor.

Warnings and When to Ask Your Doctor

Allergy alert: Do not use if you are allergic to lansoprazole

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking

- warfarin (blood-thinning medicine)
- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)
- theophylline (asthma medicine)
- tacrolimus or mycophenolate mofetil (immune system medicines)
- atazanavir (medicine for HIV infection)
- methotrexate (arthritis medication)

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

Clinical studies prove PREVACID[®]24HR effectively treats frequent heartburn

In three clinical studies, PREVACID[®]24HR was shown to be significantly better than placebo in treating frequent heartburn.

How PREVACID[®]24HR is Sold

PREVACID[®]24HR is available in 14 capsule, 28 capsule and 42 capsule sizes. These sizes contain one, two and three 14-day courses of treatment, respectively. Do not use for more than 14 days in a row unless directed by your doctor. For the 28 count (two 14-day courses) and the 42 count (three 14-day courses), you may repeat a 14-day course every 4 months.

For Questions or Comments About PREVACID[®]24HR

Call 1-800-452-0051

Distributed by: **GSK Consumer Healthcare**, Warren, NJ 07059
 Trademarks are owned by or licensed to the GSK group of companies.
 ©2016 GSK group of companies or its licensors. All rights reserved.
 PREVACID[®] is a registered trademark of Takeda Pharmaceuticals North America, Inc., and is used under license. 168906-PI-L

(b) (4)

Reference ID: 4187124

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

VALERIE S PRATT
11/28/2017