

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use MOXEZA® solution safely and effectively. See full prescribing information for MOXEZA®.

**MOXEZA® (moxifloxacin ophthalmic solution) 0.5%
Sterile topical ophthalmic solution
Initial U.S. Approval: 1999**

INDICATIONS AND USAGE

MOXEZA® solution is a topical fluoroquinolone anti-infective indicated for the treatment of bacterial conjunctivitis caused by susceptible strains of the following organisms:

*Aerococcus viridans**, *Corynebacterium macginleyi**,
*Enterococcus faecalis**, *Micrococcus luteus**,
*Staphylococcus arlettae**, *Staphylococcus aureus*,
Staphylococcus capitis, *Staphylococcus epidermidis*,
Staphylococcus haemolyticus, *Staphylococcus hominis*,
*Staphylococcus saprophyticus**, *Staphylococcus warneri**,
*Streptococcus mitis**, *Streptococcus pneumoniae*,
*Streptococcus parasanguinis**, *Escherichia coli**,
Haemophilus influenzae, *Klebsiella pneumoniae**,
Propionibacterium acnes, *Chlamydia trachomatis**

*Efficacy for this organism was studied in fewer than 10 infections. (1)

DOSAGE AND ADMINISTRATION

Instill 1 drop in the affected eye(s) 2 times daily for 7 days. (2)

DOSAGE FORMS AND STRENGTHS

4 mL bottle filled with 3 mL sterile ophthalmic solution of moxifloxacin 0.5%. (3)

CONTRAINDICATIONS

None. (4)

WARNINGS AND PRECAUTIONS

- Topical ophthalmic use only. (5.1)
- Hypersensitivity and anaphylaxis have been reported with systemic use of moxifloxacin. (5.2)
- Prolonged use may result in overgrowth of non-susceptible organisms, including fungi. (5.3)
- Patients should not wear contact lenses if they have signs or symptoms of bacterial conjunctivitis. (5.4)

ADVERSE REACTIONS

The most common adverse reactions reported in 1 to 2% of patients were eye irritation, pyrexia, and conjunctivitis. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Alcon Laboratories, Inc. at 1-800-757-9195 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

See 17 for PATIENT COUNSELING INFORMATION.

Revised: XX/2016

FULL PRESCRIBING INFORMATION: CONTENTS*

- 1 INDICATIONS AND USAGE
- 2 DOSAGE AND ADMINISTRATION
- 3 DOSAGE FORMS AND STRENGTHS
- 4 CONTRAINDICATIONS
- 5 WARNINGS AND PRECAUTIONS
 - 5.1 Topical Ophthalmic Use Only
 - 5.2 Hypersensitivity Reactions
 - 5.3 Growth of Resistant Organisms with Prolonged Use
 - 5.4 Avoidance of Contact Lens Wear
- 6 ADVERSE REACTIONS
- 8 USE IN SPECIFIC POPULATIONS
 - 8.1 Pregnancy
 - 8.3 Nursing Mothers
 - 8.4 Pediatric Use
 - 8.5 Geriatric Use

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.3 Pharmacokinetics
- 12.4 Microbiology

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

- 17.1 Avoid Contamination of the Product
- 17.2 Avoid Contact Lens Wear
- 17.3 Hypersensitivity Reactions

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

MOXEZA® solution is indicated for the treatment of bacterial conjunctivitis caused by susceptible strains of the following organisms:

*Aerococcus viridans**
*Corynebacterium macginleyi**
*Enterococcus faecalis**
*Micrococcus luteus**
*Staphylococcus arlettae**
Staphylococcus aureus
Staphylococcus capitis
Staphylococcus epidermidis
Staphylococcus haemolyticus
Staphylococcus hominis
*Staphylococcus saprophyticus**
*Staphylococcus warneri**
*Streptococcus mitis**
Streptococcus pneumoniae
*Streptococcus parasanguinis**
*Escherichia coli**
Haemophilus influenzae
*Klebsiella pneumoniae**
Propionibacterium acnes
*Chlamydia trachomatis**

*Efficacy for this organism was studied in fewer than 10 infections.

2 DOSAGE AND ADMINISTRATION

Instill 1 drop in the affected eye(s) 2 times daily for 7 days.

3 DOSAGE FORMS AND STRENGTHS

4 mL bottle filled with 3 mL of sterile ophthalmic solution of moxifloxacin 0.5%.

4 CONTRAINDICATIONS

None.

5 WARNINGS AND PRECAUTIONS

5.1 Topical Ophthalmic Use Only

NOT FOR INJECTION. MOXEZA® solution is for topical ophthalmic use only and should not be injected subconjunctivally or introduced directly into the anterior chamber of the eye.

5.2 Hypersensitivity Reactions

In patients receiving systemically administered quinolones, including moxifloxacin, serious and occasionally fatal hypersensitivity (anaphylactic) reactions have been reported, some following the first dose. Some reactions were accompanied by cardiovascular collapse, loss of consciousness, angioedema (including laryngeal, pharyngeal or facial edema), airway obstruction, dyspnea, urticaria, and itching. If an allergic reaction to moxifloxacin occurs, discontinue use of the drug. Serious acute hypersensitivity reactions may require immediate emergency treatment. Oxygen and airway management should be administered as clinically indicated.

5.3 Growth of Resistant Organisms with Prolonged Use

As with other anti-infectives, prolonged use may result in overgrowth of non-susceptible organisms, including fungi. If superinfection occurs, discontinue use and institute alternative therapy. Whenever clinical judgment dictates, the patient should be examined with the aid of magnification, such as slit-lamp biomicroscopy, and, where appropriate, fluorescein staining.

5.4 Avoidance of Contact Lens Wear

Patients should be advised not to wear contact lenses if they have signs or symptoms of bacterial conjunctivitis.

6 ADVERSE REACTIONS

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to the rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The data described below reflect exposure to MOXEZA® solution in 1263 patients, between 4 months and 92 years of age, with signs and symptoms of bacterial conjunctivitis. The most frequently reported adverse reactions were eye irritation, pyrexia and conjunctivitis, reported in 1 to 2% of patients.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category C. Moxifloxacin was not teratogenic when administered to pregnant rats during organogenesis at oral doses as high as 500 mg/kg/day (approximately 25,000 times the highest recommended total daily human ophthalmic dose); however, decreased fetal body weights and slightly delayed fetal skeletal development were observed. There was no evidence of teratogenicity when pregnant Cynomolgus monkeys were given oral doses as high as 100 mg/kg/day (approximately 5,000 times the highest recommended total daily human ophthalmic dose). An increased incidence of smaller fetuses was observed at 100 mg/kg/day.

Since there are no adequate and well-controlled studies in pregnant women, MOXEZA® solution should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

8.3 Nursing Mothers

Moxifloxacin has not been measured in human milk, although it can be presumed to be excreted in human milk. Caution should be exercised when MOXEZA® solution is administered to a nursing mother.

8.4 Pediatric Use

The safety and effectiveness of MOXEZA® solution in infants below 4 months of age have not been established.

There is no evidence that the ophthalmic administration of moxifloxacin has any effect on weight bearing joints, even though oral administration of some quinolones has been shown to cause arthropathy in immature animals.


8.5 Geriatric Use

No overall differences in safety and effectiveness have been observed between elderly and younger patients.

11 DESCRIPTION

MOXEZA® is a sterile solution for topical ophthalmic use.

Moxifloxacin hydrochloride is an 8-methoxy fluoroquinolone anti-infective, with a diazabicyclononyl ring at the C7 position.



$C_{21}H_{24}FN_3O_4 \cdot HCl$

Mol Wt 437.9

Chemical Name:

1-Cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-7-[(4aS,7aS)-octahydro-6H-pyrrolo[3,4-b]pyridin-6-yl]-4-oxo-3-quinolinecarboxylic acid, monohydrochloride.

Each mL of MOXEZA® solution contains 5.45 mg moxifloxacin hydrochloride, equivalent to 5 mg moxifloxacin base.

Inactives: Sodium chloride, xanthan gum, boric acid, sorbitol, tyloxapol, purified water, and hydrochloric acid and/or sodium hydroxide to adjust pH.

MOXEZA® is a greenish-yellow, isotonic solution with an osmolality of 300-370 mOsm/kg and a pH of approximately 7.4. Moxifloxacin hydrochloride is a slightly yellow to yellow crystalline powder.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Moxifloxacin is a member of the fluoroquinolone class of anti-infective drugs [see [Clinical Pharmacology \(12.4\)](#)].

12.3 Pharmacokinetics

Moxifloxacin steady-state plasma pharmacokinetics were evaluated in healthy adult male and female subjects who were administered multiple, bilateral, topical ocular doses of MOXEZA® solution two times daily for four days with a final dose on day 5. The average steady-state AUC₀₋₁₂ was 8.17 ± 5.31 ng*h/mL. Moxifloxacin C_{max} following twice-daily bilateral ophthalmic administration of moxifloxacin 0.5% for 5 days is approximately 0.02% of that achieved with the oral formulation of moxifloxacin hydrochloride (C_{max} following oral dosing of 400 mg AVELOX*, 4.5 ± 0.5 mcg/mL).

12.4 Microbiology

The antibacterial action of moxifloxacin results from inhibition of the topoisomerase II (DNA gyrase) and topoisomerase IV. DNA gyrase is an essential enzyme that is involved in the replication, transcription and repair of bacterial DNA. Topoisomerase IV is an enzyme known to play a key role in the partitioning of the chromosomal DNA during bacterial cell division.

The mechanism of action for quinolones, including moxifloxacin, is different from that of macrolides, aminoglycosides, or tetracyclines. Therefore, moxifloxacin may be active against pathogens that are resistant to these antibiotics and these antibiotics may be active against pathogens that are resistant to moxifloxacin. There is no cross-resistance between moxifloxacin and the aforementioned classes of antibiotics. Cross-resistance has been observed between systemic moxifloxacin and some other quinolones.

In vitro resistance to moxifloxacin develops via multiple-step mutations. Resistance to moxifloxacin occurs *in vitro* at a general frequency of between 1.8 x 10⁻⁹ to < 1x10⁻¹¹ for Gram-positive bacteria.

Moxifloxacin has been shown to be active against most strains of the following microorganisms, both *in vitro* and in clinical infections as described in the INDICATIONS AND USAGE section:

*Aerococcus viridans**
*Corynebacterium macginleyi**
*Enterococcus faecalis**
*Micrococcus luteus**
*Staphylococcus arlettae**
Staphylococcus aureus
Staphylococcus capitis
Staphylococcus epidermidis
Staphylococcus haemolyticus
Staphylococcus hominis
*Staphylococcus saprophyticus**
*Staphylococcus warneri**

*Streptococcus mitis**
Streptococcus pneumoniae
*Streptococcus parasanguinis**
*Escherichia coli**
Haemophilus influenzae
*Klebsiella pneumoniae**
Propionibacterium acnes
*Chlamydia trachomatis**

*Efficacy for this organism was studied in fewer than 10 infections.

The following *in vitro* data are available, but their clinical significance in ophthalmic infections is unknown. The safety and effectiveness of MOXEZA® solution in treating ophthalmic infections due to these organisms have not been established in adequate and well-controlled trials.

Moxifloxacin has been shown to be active *in vitro* against most strains of the microorganisms listed below. These organisms are considered susceptible when evaluated using systemic breakpoints; however, a correlation between the *in vitro* systemic breakpoint and ophthalmologic efficacy has not been established. The list of organisms is provided as guidance only in assessing the potential treatment of conjunctival infections. Moxifloxacin exhibits *in vitro* minimal inhibitory concentrations (MICs) of 2 mcg/mL or less (systemic susceptible breakpoint) against most ($\geq 90\%$) strains of the following ocular pathogens.

Aerobic Gram-positive microorganisms:

Staphylococcus caprae
Staphylococcus cohnii
Staphylococcus lugdunensis
Staphylococcus pasteurii
Streptococcus agalactiae
Streptococcus milleri group
Streptococcus oralis
Streptococcus pyogenes
Streptococcus salivarius
Streptococcus sanguis

Aerobic Gram-negative microorganisms:

Acinetobacter baumannii
Acinetobacter calcoaceticus
Acinetobacter junii
Enterobacter aerogenes
Enterobacter cloacae
Haemophilus parainfluenzae
Klebsiella oxytoca
Moraxella catarrhalis
Moraxella osloensis
Morganella morganii
Neisseria gonorrhoeae
Neisseria meningitidis
Pantoea agglomerans
Proteus vulgaris
Pseudomonas stutzeri
Serratia liquefaciens
Serratia marcescens
Stenotrophomonas maltophilia

Anaerobic microorganisms:

Clostridium perfringens

Peptostreptococcus anaerobius
Peptostreptococcus magnus
Peptostreptococcus micros
Peptostreptococcus prevotii

Other microorganisms:

Mycobacterium tuberculosis
Mycobacterium avium
Mycobacterium kansasii
Mycobacterium marinum

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term studies in animals to determine the carcinogenic potential of moxifloxacin have not been performed.

Moxifloxacin was not mutagenic in four bacterial strains used in the Ames *Salmonella* reversion assay. As with other quinolones, the positive response observed with moxifloxacin in strain TA 102 using the same assay may be due to the inhibition of DNA gyrase. Moxifloxacin was not mutagenic in the CHO/HGPRT mammalian cell gene mutation assay. An equivocal result was obtained in the same assay when v79 cells were used. Moxifloxacin was clastogenic in the v79 chromosome aberration assay, but it did not induce unscheduled DNA synthesis in cultured rat hepatocytes. There was no evidence of genotoxicity *in vivo* in a micronucleus test or a dominant lethal test in mice.

Moxifloxacin had no effect on fertility in male and female rats at oral doses as high as 500 mg/kg/day, approximately 25,000 times the highest recommended total daily human ophthalmic dose. At 500 mg/kg orally there were slight effects on sperm morphology (head-tail separation) in male rats and on the estrous cycle in female rats.

14 CLINICAL STUDIES

In one randomized, double-masked, multicenter, vehicle-controlled clinical trial in which patients with bacterial conjunctivitis were dosed with MOXEZA® solution 2 times a day, MOXEZA® was superior to its vehicle for both clinical and microbiological outcomes. Clinical cure achieved on Day 4 was 63% (265/424) in MOXEZA® solution treated patients, versus 51% (214/423) in vehicle treated patients. Microbiologic success (eradication of baseline pathogens) was achieved on Day 4 in 75% (316/424) of MOXEZA® solution treated patients versus 56% (237/423) of vehicle treated patients. Microbiologic eradication does not always correlate with clinical outcome in anti-infective trials.

16 HOW SUPPLIED/STORAGE AND HANDLING

MOXEZA® solution is supplied as a sterile ophthalmic solution in the Alcon DROP-TAINER® dispensing system consisting of a natural low density polyethylene bottle and dispensing plug and tan polypropylene closure. Tamper evidence is provided with a shrink band around the closure and neck area of the package.

3 mL in a 4 mL bottle - NDC 0065-0006-03

Storage: Store at 2°C to 25°C (36°F to 77°F).

17 PATIENT COUNSELING INFORMATION

17.1 Avoid Contamination of the Product

Advise patients not to touch the dropper tip to any surface to avoid contaminating the contents.

17.2 Avoid Contact Lens Wear

Advise patients not to wear contact lenses if they have signs and symptoms of bacterial conjunctivitis.

17.3 Hypersensitivity Reactions

Systemically administered quinolones, including moxifloxacin, have been associated with hypersensitivity reactions, even following a single dose. Advise patients to discontinue use immediately and contact their physician at the first sign of a rash or allergic reaction.

Licensed to Alcon by Bayer Pharma AG.

©2010-2012, 2016 Novartis

*AVELOX is a registered trademark of Bayer AG.

Alcon®
a Novartis company

Distributed by:
ALCON LABORATORIES, INC.
6201 South Freeway
Fort Worth, Texas 76134 USA
alcon.medinfo@alcon.com

XXXXXX-MMY