

pharmacokinetics and accumulated in plasma up to 36% following multiple doses of 100 mg and 300 mg.

Absorption

The mean absolute oral bioavailability of canagliflozin is approximately 65%. Co-administration of a high-fat meal with canagliflozin had no effect on the pharmacokinetics of canagliflozin; therefore, INVOKANA may be taken with or without food. However, based on the potential to reduce postprandial plasma glucose excursions due to delayed intestinal glucose absorption, it is recommended that INVOKANA be taken before the first meal of the day [*see Dosage and Administration (2.1)*].

Distribution

The mean steady-state volume of distribution of canagliflozin following a single intravenous infusion in healthy subjects was 83.5 L, suggesting extensive tissue distribution. Canagliflozin is extensively bound to proteins in plasma (99%), mainly to albumin. Protein binding is independent of canagliflozin plasma concentrations. Plasma protein binding is not meaningfully altered in patients with renal or hepatic impairment.

Metabolism

O-glucuronidation is the major metabolic elimination pathway for canagliflozin, which is mainly glucuronidated by UGT1A9 and UGT2B4 to two inactive *O*-glucuronide metabolites.

CYP3A4-mediated (oxidative) metabolism of canagliflozin is minimal (approximately 7%) in humans.

Excretion

Following administration of a single oral [¹⁴C] canagliflozin dose to healthy subjects, 41.5%, 7.0%, and 3.2% of the administered radioactive dose was recovered in feces as canagliflozin, a hydroxylated metabolite, and an *O*-glucuronide metabolite, respectively. Enterohepatic circulation of canagliflozin was negligible.

Approximately 33% of the administered radioactive dose was excreted in urine, mainly as *O*-glucuronide metabolites (30.5%). Less than 1% of the dose was excreted as unchanged canagliflozin in urine. Renal clearance of canagliflozin 100 mg and 300 mg doses ranged from 1.30 to 1.55 mL/min.

Mean systemic clearance of canagliflozin was approximately 192 mL/min in healthy subjects following intravenous administration.

Specific Populations

Renal Impairment

A single-dose, open-label study evaluated the pharmacokinetics of canagliflozin 200 mg in subjects with varying degrees of renal impairment (classified using the MDRD-eGFR formula) compared to healthy subjects.

Renal impairment did not affect the C_{\max} of canagliflozin. Compared to healthy subjects (N=3; eGFR greater than or equal to 90 mL/min/1.73 m²), plasma AUC of canagliflozin was increased by approximately 15%, 29%, and 53% in subjects with mild (N=10), moderate (N=9), and severe (N=10) renal impairment, respectively, (eGFR 60 to less than 90, 30 to less than 60 and 15 to less than 30 mL/min/1.73 m², respectively), but was similar for ESRD (N=8) subjects and healthy subjects.

Increases in canagliflozin AUC of this magnitude are not considered clinically relevant. The pharmacodynamic response to canagliflozin declines with increasing severity of renal impairment [see *Contraindications (4) and Warnings and Precautions (5.4)*].

Canagliflozin was negligibly removed by hemodialysis.

Hepatic Impairment

Relative to subjects with normal hepatic function, the geometric mean ratios for C_{\max} and AUC_{∞} of canagliflozin were 107% and 110%, respectively, in subjects with Child-Pugh class A (mild hepatic impairment) and 96% and 111%, respectively, in subjects with Child-Pugh class B (moderate hepatic impairment) following administration of a single 300 mg dose of canagliflozin.

These differences are not considered to be clinically meaningful. There is no clinical experience in patients with Child-Pugh class C (severe) hepatic impairment [see *Use in Specific Populations (8.7)*].

Pharmacokinetic Effects of Age, Body Mass Index (BMI)/Weight, Gender and Race

Based on the population PK analysis with data collected from 1526 subjects, age, body mass index (BMI)/weight, gender, and race do not have a clinically meaningful effect on the pharmacokinetics of canagliflozin [see *Use in Specific Populations (8.5)*].

Pediatric

Studies characterizing the pharmacokinetics of canagliflozin in pediatric patients have not been conducted.

Drug Interaction Studies

In Vitro Assessment of Drug Interactions

Canagliflozin did not induce CYP450 enzyme expression (3A4, 2C9, 2C19, 2B6, and 1A2) in cultured human hepatocytes. Canagliflozin did not inhibit the CYP450 isoenzymes (1A2, 2A6, 2C19, 2D6, or 2E1) and weakly inhibited CYP2B6, CYP2C8, CYP2C9, and CYP3A4 based on *in vitro* studies with human hepatic microsomes. Canagliflozin is a weak inhibitor of P-gp.

Canagliflozin is also a substrate of drug transporters P-glycoprotein (P-gp) and MRP2.

In Vivo Assessment of Drug Interactions

Table 7: Effect of Co-Administered Drugs on Systemic Exposures of Canagliflozin

Co-Administered Drug	Dose of Co-Administered Drug*	Dose of Canagliflozin*	Geometric Mean Ratio (Ratio With/Without Co-Administered Drug) No Effect=1.0	
			AUC [†] (90% CI)	C _{max} (90% CI)
See Drug Interactions (7.1) for the clinical relevance of the following:				
Rifampin	600 mg QD for 8 days	300 mg	0.49 (0.44; 0.54)	0.72 (0.61; 0.84)
No dose adjustments of INVOKANA required for the following:				
Cyclosporine	400 mg	300 mg QD for 8 days	1.23 (1.19; 1.27)	1.01 (0.91; 1.11)
Ethinyl estradiol and levonorgestrel	0.03 mg ethinyl estradiol and 0.15 mg levonorgestrel	200 mg QD for 6 days	0.91 (0.88; 0.94)	0.92 (0.84; 0.99)
Hydrochlorothiazide	25 mg QD for 35 days	300 mg QD for 7 days	1.12 (1.08; 1.17)	1.15 (1.06; 1.25)
Metformin	2,000 mg	300 mg QD for 8 days	1.10 (1.05; 1.15)	1.05 (0.96; 1.16)
Probenecid	500 mg BID for 3 days	300 mg QD for 17 days	1.21 (1.16; 1.25)	1.13 (1.00; 1.28)

* Single dose unless otherwise noted

† AUC_{inf} for drugs given as a single dose and AUC_{24h} for drugs given as multiple doses

QD = once daily; BID = twice daily

Table 8: Effect of Canagliflozin on Systemic Exposure of Co-Administered Drugs

Co-Administered Drug	Dose of Co-Administered Drug*	Dose of Canagliflozin*	Geometric Mean Ratio (Ratio With/Without Co-Administered Drug) No Effect = 1.0	
			AUC [†] (90% CI)	C _{max} (90% CI)

See Drug Interactions (7.2) for the clinical relevance of the following:					
Digoxin	0.5 mg QD first day followed by 0.25 mg QD for 6 days	300 mg QD for 7 days	digoxin	1.20 (1.12; 1.28)	1.36 (1.21; 1.53)
No dose adjustments of co-administered drug required for the following:					
Acetaminophen	1,000 mg	300 mg BID for 25 days	acetaminophen	1.06 [†] (0.98; 1.14)	1.00 (0.92; 1.09)
Ethinyl estradiol and levonorgestrel	0.03 mg ethinyl estradiol and 0.15 mg levonorgestrel	200 mg QD for 6 days	ethinyl estradiol	1.07 (0.99; 1.15)	1.22 (1.10; 1.35)
			levonorgestrel	1.06 (1.00; 1.13)	1.22 (1.11; 1.35)
Glyburide	1.25 mg	200 mg QD for 6 days	glyburide	1.02 (0.98; 1.07)	0.93 (0.85; 1.01)
			3-cis-hydroxy-glyburide	1.01 (0.96; 1.07)	0.99 (0.91; 1.08)
			4-trans-hydroxy-glyburide	1.03 (0.97; 1.09)	0.96 (0.88; 1.04)
Hydrochlorothiazide	25 mg QD for 35 days	300 mg QD for 7 days	hydrochlorothiazide	0.99 (0.95; 1.04)	0.94 (0.87; 1.01)
Metformin	2,000 mg	300 mg QD for 8 days	metformin	1.20 (1.08; 1.34)	1.06 (0.93; 1.20)
Simvastatin	40 mg	300 mg QD for 7 days	simvastatin	1.12 (0.94; 1.33)	1.09 (0.91; 1.31)
			simvastatin acid	1.18 (1.03; 1.35)	1.26 (1.10; 1.45)
Warfarin	30 mg	300 mg QD for 12 days	(R)-warfarin	1.01 (0.96; 1.06)	1.03 (0.94; 1.13)
			(S)-warfarin	1.06 (1.00; 1.12)	1.01 (0.90; 1.13)
			INR	1.00 (0.98; 1.03)	1.05 (0.99; 1.12)

* Single dose unless otherwise noted

[†] AUC_{inf} for drugs given as a single dose and AUC_{24h} for drugs given as multiple doses

[‡] AUC_{0-12h}

QD = once daily; BID = twice daily; INR = International Normalized Ratio

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Carcinogenicity was evaluated in 2-year studies conducted in CD1 mice and Sprague-Dawley rats. Canagliflozin did not increase the incidence of tumors in mice dosed at 10, 30, or 100 mg/kg (less than or equal to 14 times exposure from a 300 mg clinical dose).

Testicular Leydig cell tumors, considered secondary to increased luteinizing hormone (LH), increased significantly in male rats at all doses tested (10, 30, and 100 mg/kg). In a 12-week clinical study, LH did not increase in males treated with canagliflozin.

Renal tubular adenoma and carcinoma increased significantly in male and female rats dosed at 100 mg/kg, or approximately 12-times exposure from a 300 mg clinical dose. Also, adrenal pheochromocytoma increased significantly in males and numerically in females dosed at 100 mg/kg. Carbohydrate malabsorption associated with high doses of canagliflozin was considered a necessary proximal event in the emergence of renal and adrenal tumors in rats. Clinical studies have not demonstrated carbohydrate malabsorption in humans at canagliflozin doses of up to 2-times the recommended clinical dose of 300 mg.

Mutagenesis

Canagliflozin was not mutagenic with or without metabolic activation in the Ames assay. Canagliflozin was mutagenic in the *in vitro* mouse lymphoma assay with but not without metabolic activation. Canagliflozin was not mutagenic or clastogenic in an *in vivo* oral micronucleus assay in rats and an *in vivo* oral Comet assay in rats.

Impairment of Fertility

Canagliflozin had no effects on the ability of rats to mate and sire or maintain a litter up to the high dose of 100 mg/kg (approximately 14 times and 18 times the 300 mg clinical dose in males and females, respectively), although there were minor alterations in a number of reproductive parameters (decreased sperm velocity, increased number of abnormal sperm, slightly fewer corpora lutea, fewer implantation sites, and smaller litter sizes) at the highest dosage administered.

14 CLINICAL STUDIES

INVOKANA (canagliflozin) has been studied as monotherapy, in combination with metformin, sulfonylurea, metformin and sulfonylurea, metformin and sitagliptin, metformin and a thiazolidinedione (i.e., pioglitazone), and in combination with insulin (with or without other antihyperglycemic agents). The efficacy of INVOKANA was compared to a dipeptidyl peptidase-4 (DPP-4) inhibitor (sitagliptin), both as add-on combination therapy with metformin and sulfonylurea, and a sulfonylurea (glimepiride), both as add-on combination therapy with metformin. INVOKANA was also evaluated in adults 55 to 80 years of age and patients with moderate renal impairment.

In patients with type 2 diabetes, treatment with INVOKANA produced clinically and statistically significant improvements in HbA_{1C} compared to placebo. Reductions in HbA_{1C} were observed across subgroups including age, gender, race, and baseline body mass index (BMI).

14.1 Monotherapy

A total of 584 patients with type 2 diabetes inadequately controlled on diet and exercise participated in a 26-week, double-blind, placebo-controlled study to evaluate the efficacy and safety of INVOKANA. The mean age was 55 years, 44% of patients were men, and the mean baseline eGFR was 87 mL/min/1.73 m². Patients taking other antihyperglycemic agents (N=281) discontinued the agent and underwent an 8-week washout followed by a 2-week, single-blind, placebo run-in period. Patients not taking oral antihyperglycemic agents (N=303) entered the 2-week, single-blind, placebo run-in period directly. After the placebo run-in period, patients were randomized to INVOKANA 100 mg, INVOKANA 300 mg, or placebo, administered once daily for 26 weeks.

At the end of treatment, INVOKANA 100 mg and 300 mg once daily resulted in a statistically significant improvement in HbA_{1C} (p<0.001 for both doses) compared to placebo. INVOKANA 100 mg and 300 mg once daily also resulted in a greater proportion of patients achieving an HbA_{1C} less than 7%, in significant reduction in fasting plasma glucose (FPG), in improved postprandial glucose (PPG), and in percent body weight reduction compared to placebo (see Table 9). Statistically significant (p<0.001 for both doses) mean changes from baseline in systolic blood pressure relative to placebo were -3.7 mmHg and -5.4 mmHg with INVOKANA 100 mg and 300 mg, respectively.

Table 9: Results from 26-Week Placebo-Controlled Clinical Study with INVOKANA as Monotherapy*

Efficacy Parameter	Placebo (N=192)	INVOKANA 100 mg (N=195)	INVOKANA 300 mg (N=197)
HbA_{1C} (%)			
Baseline (mean)	7.97	8.06	8.01
Change from baseline (adjusted mean)	0.14	-0.77	-1.03
Difference from placebo (adjusted mean) (95% CI) [†]		-0.91 [‡] (-1.09; -0.73)	-1.16 [‡] (-1.34; -0.99)
Percent of Patients Achieving HbA_{1C} < 7%	21	45 [‡]	62 [‡]
Fasting Plasma Glucose (mg/dL)			
Baseline (mean)	166	172	173
Change from baseline (adjusted mean)	8	-27	-35
Difference from placebo (adjusted mean) (95% CI) [†]		-36 [‡] (-42; -29)	-43 [‡] (-50; -37)
2-hour Postprandial Glucose (mg/dL)			
Baseline (mean)	229	250	254
Change from baseline (adjusted mean)	5	-43	-59
Difference from placebo (adjusted mean) (95% CI) [†]		-48 [‡] (-59.1; -37.0)	-64 [‡] (-75.0; -52.9)
Body Weight			
Baseline (mean) in kg	87.5	85.9	86.9
% change from baseline (adjusted mean)	-0.6	-2.8	-3.9
Difference from placebo (adjusted mean) (95% CI) [†]		-2.2 [‡] (-2.9; -1.6)	-3.3 [‡] (-4.0; -2.6)

* Intent-to-treat population using last observation in study prior to glycemic rescue therapy
 † Least squares mean adjusted for baseline value and stratification factors
 ‡ p<0.001

14.2 Combination Therapy

Add-on Combination Therapy with Metformin

A total of 1284 patients with type 2 diabetes inadequately controlled on metformin monotherapy (greater than or equal to 2,000 mg/day, or at least 1,500 mg/day if higher dose not tolerated) participated in a 26-week, double-blind, placebo- and active-controlled study to evaluate the efficacy and safety of INVOKANA in combination with metformin. The mean age was 55 years, 47% of patients were men, and the mean baseline eGFR was 89 mL/min/1.73 m². Patients already on the required metformin dose (N=1009) were randomized after completing a 2-week, single-blind, placebo run-in period. Patients taking less than the required metformin dose or patients on metformin in combination with another antihyperglycemic agent (N=275) were switched to metformin monotherapy (at doses described above) for at least 8 weeks before entering the 2-week, single-blind, placebo run-in. After the placebo run-in period, patients were randomized to INVOKANA 100 mg, INVOKANA 300 mg, sitagliptin 100 mg, or placebo, administered once daily as add-on therapy to metformin.

At the end of treatment, INVOKANA 100 mg and 300 mg once daily resulted in a statistically significant improvement in HbA_{1C} (p<0.001 for both doses) compared to placebo when added to metformin. INVOKANA 100 mg and 300 mg once daily also resulted in a greater proportion of patients achieving an HbA_{1C} less than 7%, in significant reduction in fasting plasma glucose (FPG), in improved postprandial glucose (PPG), and in percent body weight reduction compared to placebo when added to metformin (see Table 10). Statistically significant (p<0.001 for both doses) mean changes from baseline in systolic blood pressure relative to placebo were -5.4 mmHg and -6.6 mmHg with INVOKANA 100 mg and 300 mg, respectively.

Table 10: Results from 26-Week Placebo-Controlled Clinical Study of INVOKANA in Combination with Metformin*

Efficacy Parameter	Placebo + Metformin (N=183)	INVOKANA 100 mg + Metformin (N=368)	INVOKANA 300 mg + Metformin (N=367)
HbA_{1C} (%)			
Baseline (mean)	7.96	7.94	7.95
Change from baseline (adjusted mean)	-0.17	-0.79	-0.94
Difference from placebo (adjusted mean) (95% CI) [†]		-0.62 [‡] (-0.76; -0.48)	-0.77 [‡] (-0.91; -0.64)
Percent of patients achieving HbA_{1C} < 7%	30	46 [‡]	58 [‡]
Fasting Plasma Glucose (mg/dL)			
Baseline (mean)	164	169	173

Change from baseline (adjusted mean)	2	-27	-38
Difference from placebo (adjusted mean) (95% CI) [†]		-30 [‡] (-36; -24)	-40 [‡] (-46; -34)
2-hour Postprandial Glucose (mg/dL)			
Baseline (mean)	249	258	262
Change from baseline (adjusted mean)	-10	-48	-57
Difference from placebo (adjusted mean) (95% CI) [†]		-38 [‡] (-49; -27)	-47 [‡] (-58; -36)
Body Weight			
Baseline (mean) in kg	86.7	88.7	85.4
% change from baseline (adjusted mean)	-1.2	-3.7	-4.2
Difference from placebo (adjusted mean) (95% CI) [†]		-2.5 [‡] (-3.1; -1.9)	-2.9 [‡] (-3.5; -2.3)

* Intent-to-treat population using last observation in study prior to glycemic rescue therapy

[†] Least squares mean adjusted for baseline value and stratification factors

[‡] p<0.001

Initial Combination Therapy with Metformin

A total of 1186 patients with type 2 diabetes inadequately controlled with diet and exercise participated in a 26-week double-blind, active-controlled, parallel-group, 5-arm, multicenter study to evaluate the efficacy and safety of initial therapy with INVOKANA in combination with metformin XR. The median age was 56 years, 48% of patients were men, and the mean baseline eGFR was 87.6 mL/min/1.73 m². The median duration of diabetes was 1.6 years, and 72% of patients were treatment naïve. After completing a 2-week single-blind placebo run-in period, patients were randomly assigned for a double-blind treatment period of 26 weeks to 1 of 5 treatment groups (Table 11). The metformin XR dose was initiated at 500 mg/day for the first week of treatment and then increased to 1000 mg/day. Metformin XR or matching placebo was up-titrated every 2-3 weeks during the next 8 weeks of treatment to a maximum daily dose of 1500 to 2000 mg/day, as tolerated; about 90% of patients reached 2000 mg/day.

At the end of treatment, INVOKANA 100 mg and INVOKANA 300 mg in combination with metformin XR resulted in a statistically significant greater improvement in HbA_{1C} compared to their respective INVOKANA doses (100 mg and 300 mg) alone or metformin XR alone.

Table 11: Results from 26-Week Active-Controlled Clinical Study of INVOKANA Alone or INVOKANA as Initial Combination Therapy with Metformin*

Efficacy Parameter	Metformin XR (N=237)	INVOKANA 100 mg (N=237)	INVOKANA 300 mg (N=238)	INVOKANA 100 mg + Metformin XR (N=237)	INVOKANA 300 mg + Metformin XR (N=237)
HbA_{1C} (%)					
Baseline (mean)	8.81	8.78	8.77	8.83	8.90
Change from baseline (adjusted mean) [†]	-1.30	-1.37	-1.42	-1.77	-1.78
Difference from canagliflozin 100 mg				-0.40 [‡] (-0.59, -0.21)	

31

(adjusted mean) (95% CI) †					
Difference from canagliflozin 300 mg (adjusted mean) (95% CI) †					-0.36‡ (-0.56, -0.17)
Difference from metformin XR (adjusted mean) (95% CI) †		-0.06‡‡ (-0.26, 0.13)	-0.11‡‡ (-0.31, 0.08)	-0.46‡ (-0.66, -0.27)	-0.48‡ (-0.67, -0.28)
Percent of patients achieving HbA_{1C} < 7%	38	34	39	47 ^{§§}	51 ^{§§}

* Intent-to-treat population

† Least squares mean adjusted for covariates including baseline value and stratification factor

‡ Adjusted p=0.001 for superiority

‡‡ Adjusted p=0.001 for non-inferiority

§§ Adjusted p<0.05

¶ There were 121 patients without week 26 efficacy data. Analyses addressing missing data gave consistent results with the results provided in this table.

INVOKANA Compared to Glimepiride, Both as Add-on Combination With Metformin

A total of 1450 patients with type 2 diabetes inadequately controlled on metformin monotherapy (greater than or equal to 2,000 mg/day, or at least 1,500 mg/day if higher dose not tolerated) participated in a 52-week, double-blind, active-controlled study to evaluate the efficacy and safety of INVOKANA in combination with metformin.

The mean age was 56 years, 52% of patients were men, and the mean baseline eGFR was 90 mL/min/1.73 m². Patients tolerating maximally required metformin dose (N=928) were randomized after completing a 2-week, single-blind, placebo run-in period. Other patients (N=522) were switched to metformin monotherapy (at doses described above) for at least 10 weeks, then completed a 2-week single-blind run-in period. After the 2-week run-in period, patients were randomized to INVOKANA 100 mg, INVOKANA 300 mg, or glimepiride (titration allowed throughout the 52-week study to 6 or 8 mg), administered once daily as add-on therapy to metformin.

As shown in Table 12 and Figure 1, at the end of treatment, INVOKANA 100 mg provided similar reductions in HbA_{1C} from baseline compared to glimepiride when added to metformin therapy. INVOKANA 300 mg provided a greater reduction from baseline in HbA_{1C} compared to glimepiride, and the relative treatment difference was -0.12% (95% CI: -0.22; -0.02). As shown in Table 12, treatment with INVOKANA 100 mg and 300 mg daily provided greater improvements in percent body weight change, relative to glimepiride.

Table 12: Results from 52-Week Clinical Study Comparing INVOKANA to Glimepiride in Combination with Metformin*

Efficacy Parameter	INVOKANA 100 mg + Metformin (N=483)	INVOKANA 300 mg + Metformin (N=485)	Glimepiride (titrated) + Metformin (N=482)
HbA_{1c} (%)			
Baseline (mean)	7.78	7.79	7.83
Change from baseline (adjusted mean)	-0.82	-0.93	-0.81
Difference from glimepiride (adjusted mean) (95% CI) [†]	-0.01 [‡] (-0.11; 0.09)	-0.12 [‡] (-0.22; -0.02)	
Percent of patients achieving HbA_{1c} < 7%	54	60	56
Fasting Plasma Glucose (mg/dL)			
Baseline (mean)	165	164	166
Change from baseline (adjusted mean)	-24	-28	-18
Difference from glimepiride (adjusted mean) (95% CI) [†]	-6 (-10; -2)	-9 (-13; -5)	
Body Weight			
Baseline (mean) in kg	86.8	86.6	86.6
% change from baseline (adjusted mean)	-4.2	-4.7	1.0
Difference from glimepiride (adjusted mean) (95% CI) [†]	-5.2 [§] (-5.7; -4.7)	-5.7 [§] (-6.2; -5.1)	


* Intent-to-treat population using last observation in study prior to glycemic rescue therapy

[†] Least squares mean adjusted for baseline value and stratification factors

[‡] INVOKANA + metformin is considered non-inferior to glimepiride + metformin because the upper limit of this confidence interval is less than the pre-specified non-inferiority margin of < 0.3%.

[§] p<0.001

Figure 1: Mean HbA_{1c} Change at Each Time Point (Completers) and at Week 52 Using Last Observation Carried Forward (mITT Population)



Add-on Combination Therapy With Sulfonylurea

A total of 127 patients with type 2 diabetes inadequately controlled on sulfonylurea monotherapy participated in an 18-week, double-blind, placebo-controlled sub-study to evaluate the efficacy

and safety of INVOKANA in combination with sulfonylurea. The mean age was 65 years, 57% of patients were men, and the mean baseline eGFR was 69 mL/min/1.73 m². Patients treated with sulfonylurea monotherapy on a stable protocol-specified dose (greater than or equal to 50% maximal dose) for at least 10 weeks completed a 2-week, single-blind, placebo run-in period. After the run-in period, patients with inadequate glycemic control were randomized to INVOKANA 100 mg, INVOKANA 300 mg, or placebo, administered once daily as add-on to sulfonylurea.

As shown in Table 13, at the end of treatment, INVOKANA 100 mg and 300 mg daily provided statistically significant (p<0.001 for both doses) improvements in HbA_{1C} relative to placebo when added to sulfonylurea. INVOKANA 300 mg once daily compared to placebo resulted in a greater proportion of patients achieving an HbA_{1C} less than 7%, (33% vs 5%), greater reductions in fasting plasma glucose (-36 mg/dL vs +12 mg/dL), and greater percent body weight reduction (-2.0% vs -0.2%).

Table 13: Results from 18-Week Placebo–Controlled Clinical Study of INVOKANA in Combination with Sulfonylurea*

Efficacy Parameter	Placebo + Sulfonylurea (N=45)	INVOKANA 100 mg + Sulfonylurea (N=42)	INVOKANA 300 mg + Sulfonylurea (N=40)
HbA_{1C} (%)			
Baseline (mean)	8.49	8.29	8.28
Change from baseline (adjusted mean)	0.04	-0.70	-0.79
Difference from placebo (adjusted mean) (95% CI) [†]		-0.74 [‡] (-1.15; -0.33)	-0.83 [‡] (-1.24; -0.41)

* Intent-to-treat population using last observation in study prior to glycemic rescue therapy

[†] Least squares mean adjusted for baseline value

[‡] p<0.001

Add-on Combination Therapy With Metformin and Sulfonylurea

A total of 469 patients with type 2 diabetes inadequately controlled on the combination of metformin (greater than or equal to 2,000 mg/day or at least 1,500 mg/day if higher dose not tolerated) and sulfonylurea (maximal or near-maximal effective dose) participated in a 26-week, double-blind, placebo-controlled study to evaluate the efficacy and safety of INVOKANA in combination with metformin and sulfonylurea. The mean age was 57 years, 51% of patients were men, and the mean baseline eGFR was 89 mL/min/1.73 m². Patients already on the protocol-specified doses of metformin and sulfonylurea (N=372) entered a 2-week, single-blind, placebo run-in period. Other patients (N=97) were required to be on a stable protocol-specified dose of metformin and sulfonylurea for at least 8 weeks before entering the 2-week run-in period.

Following the run-in period, patients were randomized to INVOKANA 100 mg, INVOKANA 300 mg, or placebo, administered once daily as add-on to metformin and sulfonylurea.

At the end of treatment, INVOKANA 100 mg and 300 mg once daily resulted in a statistically significant improvement in HbA_{1C} (p<0.001 for both doses) compared to placebo when added to metformin and sulfonylurea. INVOKANA 100 mg and 300 mg once daily also resulted in a greater proportion of patients achieving an HbA_{1C} less than 7%, in a significant reduction in fasting plasma glucose (FPG), and in percent body weight reduction compared to placebo when added to metformin and sulfonylurea (see Table 14).

Table 14: Results from 26-Week Placebo-Controlled Clinical Study of INVOKANA in Combination with Metformin and Sulfonylurea*

Efficacy Parameter	Placebo + Metformin and Sulfonylurea (N=156)	INVOKANA 100 mg + Metformin and Sulfonylurea (N=157)	INVOKANA 300 mg + Metformin and Sulfonylurea (N=156)
HbA_{1C} (%)			
Baseline (mean)	8.12	8.13	8.13
Change from baseline (adjusted mean)	-0.13	-0.85	-1.06
Difference from placebo (adjusted mean) (95% CI) [†]		-0.71 [‡] (-0.90; -0.52)	-0.92 [‡] (-1.11; -0.73)
Percent of patients achieving A_{1C} < 7%	18	43 [‡]	57 [‡]
Fasting Plasma Glucose (mg/dL)			
Baseline (mean)	170	173	168
Change from baseline (adjusted mean)	4	-18	-31
Difference from placebo (adjusted mean) (95% CI) [†]		-22 [‡] (-31; -13)	-35 [‡] (-44; -25)
Body Weight			
Baseline (mean) in kg	90.8	93.5	93.5
% change from baseline (adjusted mean)	-0.7	-2.1	-2.6
Difference from placebo (adjusted mean) (95% CI) [†]		-1.4 [‡] (-2.1; -0.7)	-2.0 [‡] (-2.7; -1.3)

* Intent-to-treat population using last observation in study prior to glycemic rescue therapy

[†] Least squares mean adjusted for baseline value and stratification factors

[‡] p<0.001

Add-on Combination Therapy With Metformin and Sitagliptin

A total of 217 patients with type 2 diabetes inadequately controlled on the combination of metformin (greater than or equal to 1,500 mg/day) and sitagliptin 100 mg/day (or equivalent fixed-dose combination) participated in a 26-week, double-blind, placebo-controlled study to evaluate the efficacy and safety of INVOKANA in combination with metformin and sitagliptin. The mean age was 57 years, 58% of patients were men, 73% of patients were Caucasian, 15% were Asian, and 12% were Black or African-American. The mean baseline eGFR was 90 mL/min/1.73 m² and the mean baseline BMI was 32 kg/m². The mean duration of diabetes

was 10 years. Eligible patients entered a 2-week, single-blind, placebo run-in period and were subsequently randomized to INVOKANA 100 mg or placebo, administered once daily as add-on to metformin and sitagliptin. Patients with a baseline eGFR of 70 mL/min/1.73 m² or greater who were tolerating INVOKANA 100 mg and who required additional glycemic control (fasting finger stick 100 mg/dL or greater at least twice within 2 weeks) were up-titrated to INVOKANA 300 mg. While up-titration occurred as early as Week 4, most (90%) patients randomized to INVOKANA were up-titrated to INVOKANA 300 mg by 6 to 8 weeks.

At the end of 26 weeks, INVOKANA resulted in a statistically significant improvement in HbA_{1C} (p<0.001) compared to placebo when added to metformin and sitagliptin.

Table 15: Results from 26-Week Placebo-Controlled Clinical Study of INVOKANA in Combination with Metformin and Sitagliptin

Efficacy Parameter	Placebo + Metformin and Sitagliptin (N=108*)	INVOKANA + Metformin and Sitagliptin (N=109*)
HbA_{1C} (%)		
Baseline (mean)	8.40	8.50
Change from baseline (adjusted mean)	-0.03	-0.83
Difference from placebo (adjusted mean) (95% CI) ^{†§}		-0.81 [#] (-1.11; -0.51)
Percent of patients achieving HbA_{1C} < 7%[‡]	9	28
Fasting Plasma Glucose (mg/dL)[¶]		
Baseline (mean)	180	185
Change from baseline (adjusted mean)	-3	-28
Difference from placebo (adjusted mean) (95% CI)		-25 [#] (-39; -11)

* To preserve the integrity of randomization, all randomized patients were included in the analysis. The patient who was randomized once to each arm was analyzed on INVOKANA.

† Early treatment discontinuation before week 26, occurred in 11.0% and 24.1% of INVOKANA and placebo patients, respectively.

‡ Patients without week 26 efficacy data were considered as non-responders when estimating the proportion achieving HbA_{1c} < 7%.

§ Estimated using a multiple imputation method modeling a “wash-out” of the treatment effect for patients having missing data who discontinued treatment. Missing data was imputed only at week 26 and analyzed using ANCOVA.

¶ Estimated using a multiple imputation method modeling a “wash-out” of the treatment effect for patients having missing data who discontinued treatment. A mixed model for repeated measures was used to analyze the imputed data.

p<0.001

INVOKANA Compared to Sitagliptin, Both as Add-on Combination Therapy With Metformin and Sulfonylurea

A total of 755 patients with type 2 diabetes inadequately controlled on the combination of metformin (greater than or equal to 2,000 mg/day or at least 1,500 mg/day if higher dose not

tolerated) and sulfonylurea (near-maximal or maximal effective dose) participated in a 52-week, double-blind, active-controlled study to compare the efficacy and safety of INVOKANA 300 mg versus sitagliptin 100 mg in combination with metformin and sulfonylurea. The mean age was 57 years, 56% of patients were men, and the mean baseline eGFR was 88 mL/min/1.73 m². Patients already on protocol-specified doses of metformin and sulfonylurea (N=716) entered a 2-week single-blind, placebo run-in period. Other patients (N=39) were required to be on a stable protocol-specified dose of metformin and sulfonylurea for at least 8 weeks before entering the 2-week run-in period. Following the run-in period, patients were randomized to INVOKANA 300 mg or sitagliptin 100 mg as add-on to metformin and sulfonylurea.

As shown in Table 16 and Figure 2, at the end of treatment, INVOKANA 300 mg provided greater HbA_{1C} reduction compared to sitagliptin 100 mg when added to metformin and sulfonylurea (p<0.05). INVOKANA 300 mg resulted in a mean percent change in body weight from baseline of -2.5% compared to +0.3% with sitagliptin 100 mg. A mean change in systolic blood pressure from baseline of -5.06 mmHg was observed with INVOKANA 300 mg compared to +0.85 mmHg with sitagliptin 100 mg.

Table 16: Results from 52-Week Clinical Study Comparing INVOKANA to Sitagliptin in Combination with Metformin and Sulfonylurea*

Efficacy Parameter	INVOKANA 300 mg + Metformin and Sulfonylurea (N=377)	Sitagliptin 100 mg + Metformin and Sulfonylurea (N=378)
HbA_{1C} (%)		
Baseline (mean)	8.12	8.13
Change from baseline (adjusted mean)	-1.03	-0.66
Difference from sitagliptin (adjusted mean) (95% CI) [†]	-0.37 [‡] (-0.50; -0.25)	
Percent of patients achieving HbA_{1C} < 7%	48	35
Fasting Plasma Glucose (mg/dL)		
Baseline (mean)	170	164
Change from baseline (adjusted mean)	-30	-6
Difference from sitagliptin (adjusted mean) (95% CI) [†]	-24 (-30; -18)	
Body Weight		
Baseline (mean) in kg	87.6	89.6
% change from baseline (adjusted mean)	-2.5	0.3
Difference from sitagliptin (adjusted mean) (95% CI) [†]	-2.8 [§] (-3.3; -2.2)	


* Intent-to-treat population using last observation in study prior to glycemic rescue therapy

[†] Least squares mean adjusted for baseline value and stratification factors

[‡] INVOKANA + metformin + sulfonylurea is considered non-inferior to sitagliptin + metformin + sulfonylurea because the upper limit of this confidence interval is less than the pre-specified non-inferiority margin of < 0.3%.

[§] p<0.001

Figure 2: Mean HbA_{1C} Change at Each Time Point (Completers) and at Week 52 Using Last Observation Carried Forward (mITT Population)



Add-on Combination Therapy With Metformin and Pioglitazone

A total of 342 patients with type 2 diabetes inadequately controlled on the combination of metformin (greater than or equal to 2,000 mg/day or at least 1,500 mg/day if higher dose not tolerated) and pioglitazone (30 or 45 mg/day) participated in a 26-week, double-blind, placebo-controlled study to evaluate the efficacy and safety of INVOKANA in combination with metformin and pioglitazone. The mean age was 57 years, 63% of patients were men, and the mean baseline eGFR was 86 mL/min/1.73 m². Patients already on protocol-specified doses of metformin and pioglitazone (N=163) entered a 2-week, single-blind, placebo run-in period. Other patients (N=181) were required to be on stable protocol-specified doses of metformin and pioglitazone for at least 8 weeks before entering the 2-week run-in period. Following the run-in period, patients were randomized to INVOKANA 100 mg, INVOKANA 300 mg, or placebo, administered once daily as add-on to metformin and pioglitazone.

At the end of treatment, INVOKANA 100 mg and 300 mg once daily resulted in a statistically significant improvement in HbA_{1C} (p<0.001 for both doses) compared to placebo when added to metformin and pioglitazone. INVOKANA 100 mg and 300 mg once daily also resulted in a greater proportion of patients achieving an HbA_{1C} less than 7%, in significant reduction in fasting plasma glucose (FPG) and in percent body weight reduction compared to placebo when added to metformin and pioglitazone (see Table 17). Statistically significant (p<0.05 for both doses) mean changes from baseline in systolic blood pressure relative to placebo were -4.1 mmHg and -3.5 mmHg with INVOKANA 100 mg and 300 mg, respectively.

Table 17: Results from 26-Week Placebo-Controlled Clinical Study of INVOKANA in Combination with Metformin and Pioglitazone*

Efficacy Parameter	Placebo + Metformin and Pioglitazone (N=115)	INVOKANA 100 mg + Metformin and Pioglitazone (N=113)	INVOKANA 300 mg + Metformin and Pioglitazone (N=114)
HbA_{1C} (%)			
Baseline (mean)	8.00	7.99	7.84
Change from baseline (adjusted mean)	-0.26	-0.89	-1.03
Difference from placebo (adjusted mean) (95% CI) [†]		-0.62 [‡] (-0.81; -0.44)	-0.76 [‡] (-0.95; -0.58)
Percent of patients achieving HbA_{1C} < 7%	33	47 [‡]	64 [‡]
Fasting Plasma Glucose (mg/dL)			
Baseline (mean)	164	169	164
Change from baseline (adjusted mean)	3	-27	-33
Difference from placebo (adjusted mean) (95% CI) [†]		-29 [‡] (-37; -22)	-36 [‡] (-43; -28)
Body Weight			
Baseline (mean) in kg	94.0	94.2	94.4
% change from baseline (adjusted mean)	-0.1	-2.8	-3.8
Difference from placebo (adjusted mean) (95% CI) [†]		-2.7 [‡] (-3.6; -1.8)	-3.7 [‡] (-4.6; -2.8)

* Intent-to-treat population using last observation in study prior to glycemic rescue therapy

[†] Least squares mean adjusted for baseline value and stratification factors

[‡] p<0.001

Add-On Combination Therapy With Insulin (With or Without Other Antihyperglycemic Agents)

A total of 1718 patients with type 2 diabetes inadequately controlled on insulin greater than or equal to 30 units/day or insulin in combination with other antihyperglycemic agents participated in an 18-week, double-blind, placebo-controlled substudy of a cardiovascular study to evaluate the efficacy and safety of INVOKANA in combination with insulin. The mean age was 63 years, 66% of patients were men, and the mean baseline eGFR was 75 mL/min/1.73 m². Patients on basal, bolus, or basal/bolus insulin for at least 10 weeks entered a 2-week, single-blind, placebo run-in period. Approximately 70% of patients were on a background basal/bolus insulin regimen. After the run-in period, patients were randomized to INVOKANA 100 mg, INVOKANA 300 mg, or placebo, administered once daily as add-on to insulin. The mean daily insulin dose at baseline was 83 units, which was similar across treatment groups.

At the end of treatment, INVOKANA 100 mg and 300 mg once daily resulted in a statistically significant improvement in HbA_{1C} (p<0.001 for both doses) compared to placebo when added to insulin. INVOKANA 100 mg and 300 mg once daily also resulted in a greater proportion of patients achieving an HbA_{1C} less than 7%, in significant reductions in fasting plasma glucose (FPG), and in percent body weight reductions compared to placebo (see Table 18). Statistically

significant ($p < 0.001$ for both doses) mean changes from baseline in systolic blood pressure relative to placebo were -2.6 mmHg and -4.4 mmHg with INVOKANA 100 mg and 300 mg, respectively.

Table 18: Results from 18-Week Placebo-Controlled Clinical Study of INVOKANA in Combination with Insulin \geq 30 Units/Day (With or Without Other Oral Antihyperglycemic Agents)*

Efficacy Parameter	Placebo + Insulin (N=565)	INVOKANA 100 mg + Insulin (N=566)	INVOKANA 300 mg + Insulin (N=587)
HbA_{1C} (%)			
Baseline (mean)	8.20	8.33	8.27
Change from baseline (adjusted mean)	0.01	-0.63	-0.72
Difference from placebo (adjusted mean) (95% CI) [†]		-0.65 [‡] (-0.73; -0.56)	-0.73 [‡] (-0.82; -0.65)
Percent of patients achieving HbA_{1C} < 7%	8	20 [‡]	25 [‡]
Fasting Plasma Glucose (mg/dL)			
Baseline	169	170	168
Change from baseline (adjusted mean)	4	-19	-25
Difference from placebo (adjusted mean) (97.5% CI) [‡]		-23 [‡] (-29; -16)	-29 [‡] (-35; -23)
Body Weight			
Baseline (mean) in kg	97.7	96.9	96.7
% change from baseline (adjusted mean)	0.1	-1.8	-2.3
Difference from placebo (adjusted mean) (97.5% CI) [‡]		-1.9 [‡] (-2.2; -1.6)	-2.4 [‡] (-2.7; -2.1)

* Intent-to-treat population using last observation in study prior to glycemic rescue therapy

[†] Least squares mean adjusted for baseline value and stratification factors

[‡] $p < 0.001$

14.3 Studies in Special Populations

Adults 55 to 80 Years of Age

A total of 714 older patients with type 2 diabetes inadequately controlled on current diabetes therapy (either diet and exercise alone or in combination with oral or parenteral agents) participated in a 26-week, double-blind, placebo-controlled study to evaluate the efficacy and safety of INVOKANA in combination with current diabetes treatment. The mean age was 64 years, 55% of patients were men, and the mean baseline eGFR was 77 mL/min/1.73 m². Patients were randomized to the addition of INVOKANA 100 mg, INVOKANA 300 mg, or placebo, administered once daily. At the end of treatment, INVOKANA provided statistically significant improvements from baseline relative to placebo in HbA_{1C} ($p < 0.001$ for both doses) of -0.57% (95% CI: -0.71; -0.44) for INVOKANA 100 mg and -0.70% (95% CI: -0.84; -0.57) for INVOKANA 300 mg. Statistically significant ($p < 0.001$ for both doses) reductions from baseline in fasting plasma glucose (FPG) and body weight were also observed in this study relative to placebo [see *Use in Specific Populations (8.5)*].

Moderate Renal Impairment

A total of 269 patients with type 2 diabetes and a baseline eGFR of 30 mL/min/1.73 m² to less than 50 mL/min/1.73 m² inadequately controlled on current diabetes therapy participated in a 26-week, double-blind, placebo-controlled clinical study to evaluate the efficacy and safety of INVOKANA in combination with current diabetes treatment (diet or antihyperglycemic agent therapy, with 95% of patients on insulin and/or sulfonylurea). The mean age was 68 years, 61% of patients were men, and the mean baseline eGFR was 39 mL/min/1.73 m². Patients were randomized to the addition of INVOKANA 100 mg, INVOKANA 300 mg, or placebo, administered once daily.

At the end of treatment, INVOKANA 100 mg and INVOKANA 300 mg daily provided greater reductions in HbA_{1C} relative to placebo (-0.30% [95% CI: -0.53; -0.07] and -0.40%, [95% CI: -0.64; -0.17], respectively) [*see Warnings and Precautions (5.4), Adverse Reactions (6.1), and Use in Specific Populations (8.6)*].

16 HOW SUPPLIED/STORAGE AND HANDLING

INVOKANA (canagliflozin) tablets are available in the strengths and packages listed below:

100 mg tablets are yellow, capsule-shaped, film-coated tablets with “CFZ” on one side and “100” on the other side.

NDC 50458-140-30	Bottle of 30
NDC 50458-140-90	Bottle of 90
NDC 50458-140-50	Bottle of 500
NDC 50458-140-10	Blister package containing 100 tablets (10 blister cards containing 10 tablets each)

300 mg tablets are white, capsule-shaped, film-coated tablets with “CFZ” on one side and “300” on the other side.

NDC 50458-141-30	Bottle of 30
NDC 50458-141-90	Bottle of 90
NDC 50458-141-50	Bottle of 500
NDC 50458-141-10	Blister package containing 100 tablets (10 blister cards containing 10 tablets each)

Storage and Handling

Store at 25°C (77°F); excursions permitted to 15 to 30°C (59 to 86°F).

17 PATIENT COUNSELING INFORMATION

See FDA-approved patient labeling (Medication Guide).

Instructions

Instruct patients to read the Medication Guide before starting INVOKANA (canagliflozin) therapy and to reread it each time the prescription is renewed.

Inform patients of the potential risks and benefits of INVOKANA and of alternative modes of therapy. Also inform patients about the importance of adherence to dietary instructions, regular physical activity, periodic blood glucose monitoring and HbA_{1C} testing, recognition and management of hypoglycemia and hyperglycemia, and assessment for diabetes complications. Advise patients to seek medical advice promptly during periods of stress such as fever, trauma, infection, or surgery, as medication requirements may change.

Instruct patients to take INVOKANA only as prescribed. If a dose is missed, advise patients to take it as soon as it is remembered unless it is almost time for the next dose, in which case patients should skip the missed dose and take the medicine at the next regularly scheduled time. Advise patients not to take two doses of INVOKANA at the same time.

Inform patients that the most common adverse reactions associated with INVOKANA are genital mycotic infection, urinary tract infection, and increased urination.

Inform female patients of child bearing age that the use of INVOKANA during pregnancy has not been studied in humans, and that INVOKANA should only be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. Instruct patients to report pregnancies to their physicians as soon as possible.

Inform nursing mothers to discontinue INVOKANA or nursing, taking into account the importance of drug to the mother.

Laboratory Tests

Due to its mechanism of action, patients taking INVOKANA will test positive for glucose in their urine.

Lower Limb Amputation

Inform patients that INVOKANA is associated with an increased risk of amputations. Counsel patients about the importance of routine preventative foot care. Instruct patients to monitor for new pain or tenderness, sores or ulcers, or infections involving the leg or foot and to seek medical advice immediately if such signs or symptoms develop [*see Boxed Warning and Warnings and Precautions (5.1)*].

Hypotension

Inform patients that symptomatic hypotension may occur with INVOKANA and advise them to contact their doctor if they experience such symptoms [*see Warnings and Precautions (5.2)*].

Inform patients that dehydration may increase the risk for hypotension, and to have adequate fluid intake.

Ketoacidosis

Inform patients that ketoacidosis is a serious life-threatening condition. Cases of ketoacidosis have been reported during use of INVOKANA. Instruct patients to check ketones (when possible) if symptoms consistent with ketoacidosis occur even if blood glucose is not elevated. If symptoms of ketoacidosis (including nausea, vomiting, abdominal pain, tiredness, and labored breathing) occur, instruct patients to discontinue INVOKANA and seek medical advice immediately [*see Warnings and Precautions (5.3)*].

Acute Kidney Injury

Inform patients that acute kidney injury has been reported during use of INVOKANA. Advise patients to seek medical advice immediately if they have reduced oral intake (such as due to acute illness or fasting) or increased fluid losses (such as due to vomiting, diarrhea, or excessive heat exposure), as it may be appropriate to temporarily discontinue INVOKANA use in those settings [*see Warnings and Precautions (5.4)*].

Serious Urinary Tract Infections

Inform patients of the potential for urinary tract infections, which may be serious. Provide them with information on the symptoms of urinary tract infections. Advise them to seek medical advice if such symptoms occur [*see Warnings and Precautions (5.6)*].

Genital Mycotic Infections in Females (e.g., Vulvovaginitis)

Inform female patients that vaginal yeast infection may occur and provide them with information on the signs and symptoms of vaginal yeast infection. Advise them of treatment options and when to seek medical advice [*see Warnings and Precautions (5.8)*].

Genital Mycotic Infections in Males (e.g., Balanitis or Balanoposthitis)

Inform male patients that yeast infection of penis (e.g., balanitis or balanoposthitis) may occur, especially in uncircumcised males and patients with prior history. Provide them with information on the signs and symptoms of balanitis and balanoposthitis (rash or redness of the glans or foreskin of the penis). Advise them of treatment options and when to seek medical advice [*see Warnings and Precautions (5.8)*].

Hypersensitivity Reactions

Inform patients that serious hypersensitivity reactions, such as urticaria, rash, anaphylaxis, and angioedema, have been reported with INVOKANA. Advise patients to report immediately any signs or symptoms suggesting allergic reaction, and to discontinue drug until they have consulted prescribing physicians.

Bone Fracture

Inform patients that bone fractures have been reported in patients taking INVOKANA. Provide them with information on factors that may contribute to fracture risk.

Pregnancy

Advise pregnant women, and females of reproductive potential of the potential risk to a fetus with treatment with INVOKANA [*see Use in Specific Populations (8.1)*]. Instruct females of reproductive potential to report pregnancies to their physicians as soon as possible.

Lactation

Advise women that breastfeeding is not recommended during treatment with INVOKANA [*see Use in Specific Populations (8.2)*].

Active ingredient made in Belgium

Manufactured for:

Janssen Pharmaceuticals, Inc.

Titusville, NJ 08560

Finished product manufactured by:

Janssen Ortho LLC

Gurabo, PR 00778

Or

Janssen Cilag SpA

Latina, Italy

Licensed from Mitsubishi Tanabe Pharma Corporation

© 2013 Janssen Pharmaceutical Companies

Medication Guide
INVOKANA® (in-vo-KAHN-uh)
(canagliflozin)
Tablets

What is the most important information I should know about INVOKANA?

INVOKANA can cause important side effects, including:

- **Amputations. INVOKANA may increase your risk of lower limb amputations. Amputations mainly involve removal of the toe or part of the foot, however, amputations involving the leg, below and above the knee, have also occurred. Some people had more than one amputation, some on both sides of the body.**

You may be at a higher risk of lower limb amputation if you:

- have a history of amputation
- have heart disease or are at risk for heart disease
- have had blocked or narrowed blood vessels, usually in your leg
- have damage to the nerves (neuropathy) in your leg
- have had diabetic foot ulcers or sores

Call your doctor right away if you have new pain or tenderness, any sores, ulcers, or infections in your leg or foot. Your doctor may decide to stop your INVOKANA for a while if you have any of these signs or symptoms.

Talk to your doctor about proper foot care.

- **Dehydration. INVOKANA can cause some people to become dehydrated (the loss of too much body water). Dehydration may cause you to feel dizzy, faint, lightheaded, or weak, especially when you stand up (orthostatic hypotension).**

You may be at higher risk of dehydration if you:

- have low blood pressure
- take medicines to lower your blood pressure, including diuretics (water pill)
- are on a low sodium (salt) diet
- have kidney problems
- are 65 years of age or older

Talk to your doctor about what you can do to prevent dehydration including how much fluid you should drink on a daily basis.

- **Vaginal yeast infection.** Women who take INVOKANA may get vaginal yeast infections. Symptoms of a vaginal yeast infection include:

- vaginal odor
- white or yellowish vaginal discharge (discharge may be lumpy or look like cottage cheese)
- vaginal itching

- **Yeast infection of the penis (balanitis or balanoposthitis).** Men who take INVOKANA may get a yeast infection of the skin around the penis. Certain men who are not circumcised may have swelling of the penis that makes it difficult to pull back the skin around the tip of the penis. Other symptoms of yeast infection of the penis include:

- redness, itching, or swelling of the penis
- rash of the penis
- foul smelling discharge from the penis
- pain in the skin around penis

Talk to your doctor about what to do if you get symptoms of a yeast infection of the vagina or penis. Your doctor may suggest you use an over-the-counter antifungal medicine. Talk to your doctor right away if you use an over-the-counter antifungal medication and your symptoms do not go away.

What is INVOKANA?

- INVOKANA is a prescription medicine used along with diet and exercise to lower blood sugar in adults with type 2 diabetes.
- INVOKANA is not for people with type 1 diabetes.
- INVOKANA is not for people with diabetic ketoacidosis (increased ketones in blood or urine).
- It is not known if INVOKANA is safe and effective in children under 18 years of age.

Who should not take INVOKANA?

Do not take INVOKANA if you:

- are allergic to canagliflozin or any of the ingredients in INVOKANA. See the end of this Medication Guide for a list of ingredients in INVOKANA. Symptoms of allergic reaction to INVOKANA may include:
 - rash
 - raised red patches on your skin (hives)
 - swelling of the face, lips, mouth, tongue, and throat that may cause difficulty in breathing or swallowing

- have severe kidney problems or are on dialysis.

What should I tell my doctor before taking INVOKANA?

Before you take INVOKANA, tell your doctor if you:

- have a history of amputation.
- have heart disease or are at risk for heart disease.
- have had blocked or narrowed blood vessels, usually in your leg.
- have damage to the nerves (neuropathy) in your leg.
- have had diabetic foot ulcers or sores.
- have kidney problems.
- have liver problems.
- have a history of urinary tract infections or problems with urination.
- are on a low sodium (salt) diet. Your doctor may change your diet or your dose of INVOKANA.
- are going to have surgery.
- are eating less due to illness, surgery, or a change in your diet.
- have or have had problems with your pancreas, including pancreatitis or surgery on your pancreas.
- drink alcohol very often, or drink a lot of alcohol in the short-term (“binge” drinking).
- have ever had an allergic reaction to INVOKANA.
- have other medical conditions.
- are pregnant or plan to become pregnant. INVOKANA may harm your unborn baby. If you become pregnant while taking INVOKANA, tell your doctor as soon as possible. Talk with your doctor about the best way to control your blood sugar while you are pregnant.
- are breastfeeding or plan to breastfeed. INVOKANA may pass into your breast milk and may harm your baby. Talk with your doctor about the best way to feed your baby if you are taking INVOKANA. Do not breastfeed while taking INVOKANA.

Tell your doctor about all the medicines you take, including prescription and non-prescription medicines, vitamins, and herbal supplements.

INVOKANA may affect the way other medicines work, and other medicines may affect how INVOKANA works. Especially tell your doctor if you take:

- diuretics (water pills)
- phenytoin or phenobarbital (used to control seizures)
- digoxin (Lanoxin®)* (used to treat heart problems)
- rifampin (used to treat or prevent tuberculosis)
- ritonavir (Norvir®, Kaletra®)* (used to treat HIV infection)

Ask your doctor or pharmacist for a list of these medicines if you are not sure if your medicine is listed above.

Know the medicines you take. Keep a list of them and show it to your doctor and pharmacist when you get a new medicine.

How should I take INVOKANA?

- Take INVOKANA by mouth 1 time each day exactly as your doctor tells you to take it.
- Your doctor will tell you how much INVOKANA to take and when to take it. Your doctor may change your dose if needed.
- It is best to take INVOKANA before the first meal of the day.
- Your doctor may tell you to take INVOKANA along with other diabetes medicines. Low blood sugar can happen more often when INVOKANA is taken with certain other diabetes medicines. See **“What are the possible side effects of INVOKANA?”**
- If you miss a dose, take it as soon as you remember. If it is almost time for your next dose, skip the missed dose and take the medicine at the next regularly scheduled time. Do not take two doses of INVOKANA at the same time. Talk to your doctor if you have questions about a missed dose.
- If you take too much INVOKANA, call your doctor or go to the nearest hospital emergency room right away.
- When your body is under some types of stress, such as fever, trauma (such as a car accident), infection, or surgery, the amount of diabetes medicine you need may change. Tell your doctor right away if you have any of these conditions and follow your doctor’s instructions.
- Stay on your prescribed diet and exercise program while taking INVOKANA.
- Check your blood sugar as your doctor tells you to.
- INVOKANA will cause your urine to test positive for glucose.
- Your doctor may do certain blood tests before you start INVOKANA and during treatment as needed. Your doctor may change your dose of INVOKANA based on the results of your blood tests.

- Your doctor will check your diabetes with regular blood tests, including your blood sugar levels and your hemoglobin A_{1c}.

What are the possible side effects of INVOKANA?

INVOKANA may cause serious side effects including:

See “What is the most important information I should know about INVOKANA?”

- ketoacidosis (increased ketones in your blood or urine).** Ketoacidosis has happened in people who have **type 1 diabetes or type 2 diabetes**, during treatment with INVOKANA. Ketoacidosis is a serious condition, which may need to be treated in a hospital. Ketoacidosis may lead to death. **Ketoacidosis can happen with INVOKANA even if your blood sugar is less than 250 mg/dL. Stop taking INVOKANA and call your doctor right away if you get any of the following symptoms:**

- o nausea
- o vomiting
- o stomach area (abdominal) pain
- o tiredness
- o trouble breathing

If you get any of these symptoms during treatment with INVOKANA, if possible, check for ketones in your urine, even if your blood sugar is less than 250 mg/dL.

- kidney problems.** Sudden kidney injury has happened to people taking INVOKANA. Talk to your doctor right away if you:

- o reduce the amount of food or liquid you drink for example, if you are sick or cannot eat or
- o you start to lose liquids from your body for example, from vomiting, diarrhea or being in the sun too long

- a high amount of potassium in your blood (hyperkalemia)**

- serious urinary tract infections.** Serious urinary tract infections that may lead to hospitalization have happened in people who are taking INVOKANA. Tell your doctor if you have any signs or symptoms of a urinary tract infection such as a burning feeling when passing urine, a need to urinate often, the need to urinate right away, pain in the lower part of your stomach (pelvis), or blood in the urine. Sometimes people may also have a fever, back pain, nausea, or vomiting.

- low blood sugar (hypoglycemia).** If you take INVOKANA with another medicine that can cause low blood sugar, such as a sulfonylurea or insulin, your risk of getting low blood sugar is higher. The dose of your sulfonylurea medicine or insulin may need to be lowered while you take INVOKANA.

Signs and symptoms of low blood sugar may include:

- o headache
- o drowsiness
- o weakness
- o confusion
- o dizziness
- o irritability
- o hunger
- o fast heartbeat
- o sweating
- o shaking or feeling jittery

- serious allergic reaction.** If you have any symptoms of a serious allergic reaction, stop taking INVOKANA and call your doctor right away or go to the nearest hospital emergency room. See “Who should not take INVOKANA?”. Your doctor may give you a medicine for your allergic reaction and prescribe a different medicine for your diabetes.

- broken bones (fractures).** Bone fractures have been seen in patients taking INVOKANA. Talk to your doctor about factors that may increase your risk of bone fracture.

The most common side effects of INVOKANA include:

- vaginal yeast infections and yeast infections of the penis (See “What is the most important information I should know about INVOKANA?”)

- changes in urination, including urgent need to urinate more often, in larger amounts, or at night

Tell your doctor if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of INVOKANA. For more information, ask your doctor or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

You may also report side effects to Janssen Pharmaceuticals, Inc. at 1-800-526-7736.

How should I store INVOKANA?

- Store INVOKANA at room temperature between 68°F to 77°F (20°C to 25°C).
- Keep INVOKANA and all medicines out of the reach of children.**

General information about the safe and effective use of INVOKANA.

Medicines are sometimes prescribed for purposes other than those listed in the Medication Guide. Do not use INVOKANA for a condition for which it was not prescribed. Do not give INVOKANA to other people, even if they have the same symptoms you have. It may harm them.

This Medication Guide summarizes the most important information about INVOKANA. If you would like more information, talk with your doctor. You can ask your pharmacist or doctor for information about INVOKANA that is written for healthcare professionals.

For more information about INVOKANA, call 1-800-526-7736 or visit our website at www.invokana.com.

What are the ingredients of INVOKANA?

Active ingredient: canagliflozin

Inactive ingredients: croscarmellose sodium, hydroxypropyl cellulose, lactose anhydrous, magnesium stearate, and microcrystalline cellulose. In addition, the tablet coating contains iron oxide yellow E172 (100 mg tablet only), macrogol/PEG, polyvinyl alcohol, talc, and titanium dioxide.

*The brands listed are trademarks of their respective owners and are not trademarks of Janssen Pharmaceuticals, Inc.

Active ingredient made in Belgium. Manufactured for: Janssen Pharmaceuticals, Inc., Titusville, NJ 08560. Manufactured by: Janssen Ortho LLC, Gurabo, PR 00778 or Janssen Cilag SpA, Latina, Italy. Licensed from Mitsubishi Tanabe Pharma Corporation. © 2013 Janssen Pharmaceutical Companies

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Revised 07/2017