

product containing diphenhydramine, even one used on skin • if you have sleeplessness without pain **Ask doctor before use if** • stomach bleeding warning applies to you • you have problems or serious side effects from taking pain relievers or fever reducers • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke • you are taking a diuretic • you have a breathing problem such as emphysema or chronic bronchitis • you have glaucoma • you have trouble urinating due to an enlarged prostate gland **Ask a doctor or pharmacist before use if you are** • taking sedatives or tranquilizers, or any other sleep-aid • under a doctor's care for any serious condition • taking any other antihistamines • taking any other drug **When using this product** • drowsiness will occur • avoid alcoholic drinks • do not drive a motor vehicle or operate machinery • take with food or milk if stomach upset occurs **Stop use and ask a doctor if** • you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better • you have symptoms of heart problems or stroke: • chest pain • trouble breathing • weakness in one part or side of body • slurred speech • leg swelling • pain gets worse or lasts more than 10 days • sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness. • redness or swelling is present in the painful area • any new symptoms appear • you have difficulty swallowing • it feels like the pill is stuck in your throat **If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions** • **do not take more than directed** • drink a full glass of water with each dose • adults and children 12 years and over: take 2 caplets at bedtime • do not take more than 2 caplets in 24 hours • if taken with food, this product may take longer to work **Other information** • read all warnings and directions before use • each caplet contains: sodium 20 mg • store at 20-25°C (68-77°F) • avoid high humidity and excessive heat above 40°C (104°F) **Inactive ingredients** carnauba wax, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, purified water, talc, titanium dioxide **Questions or comments? 1-800-395-0689** (Mon - Fri 9AM - 5PM EST)
Dist. by: Bayer HealthCare LLC, Whippany, NJ 07981 **MADE IN GERMANY** XXXXXX

While folded on line, tear open at dot or use scissors.
Do not use if pouch is torn or open



Active ingredients (in each caplet)

Diphenhydramine hydrochloride 25 mg	Nighttime sleep-aid
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*	Pain reliever /nonsteroidal anti-inflammatory drug

Uses • for relief of occasional sleeplessness when associated with minor aches and pains • helps you fall asleep and stay asleep **Warnings Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: • hives • facial swelling • asthma (wheezing) • shock • skin reddening • rash • blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** • if you have ever had an allergic reaction to any other pain reliever/fever reducer • unless you have time for a full night's sleep • in children under 12 years of age • right before or after heart surgery • with any other



Do not use if foil seal imprinted with "SAFETY SQUEASE" on bottle opening is missing or broken.

See Carton For Full Information. **Active ingredients (in each caplet)** Diphenhydramine hydrochloride 25 mg, Naproxen sodium 220 mg (naproxen 200 mg) **Warnings**

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin redness
- rash
- hives. If an allergic reaction occurs, stop use and seek medical

LIFT HERE

Help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart

failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer. **Ask a doctor or pharmacist before use if you are taking any other drug. If pregnant or breast-feeding, ask a health professional before use. Directions Do not take more than directed.** Adults and children 12 years and over: take 2 caplets with a full glass of water at bedtime. Do not take more than 2 caplets in 24 hours. **Questions or comments? 1-800-395-0689 (Mon-Fri 9AM - 5PM EST)**
Dist. by: Bayer HealthCare LLC, Whippany, NJ 07981



Drug Facts

Active ingredients (in each caplet)
 Diphenhydramine
 hydrochloride 25 mg Nighttime sleep aid
 Naproxen sodium 220 mg
 (naproxen 220 mg) (NSAID) Pain reliever
 *nonsteroidal anti-inflammatory drug

Purposes
 • for relief of occasional sleeplessness when associated with minor aches and pains
 • helps you fall asleep and stay asleep

Uses
 • for relief of occasional sleeplessness when associated with minor aches and pains
 • helps you fall asleep and stay asleep

Warnings
Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 • hives • facial swelling • asthma (wheezing) • shock • skin reddening
 • rash • blisters
 If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
 • are age 65 or older • have had stomach ulcers or bleeding problems
 • take a blood thinner (an anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed.
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued)

Do not use • if you have ever had an allergic reaction to any other pain reliever/fever reducer • unless you have time for a full night's sleep • in children under 12 years of age
 • right before or after heart surgery • with any other product containing diphenhydramine, even one used on skin
 • if you have sleeplessness without pain

Ask a doctor before use if • stomach bleeding warning applies to you • you have problems or serious side effects from taking pain relievers or fever reducers • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver or kidney disease, asthma, or had a stroke • you are taking a diuretic • you have a breathing problem such as emphysema or chronic bronchitis • you have glaucoma • you have trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are • taking sedatives or tranquilizers, or any other sleep aid • under a doctor's care for any serious condition • taking any other antihistamines • taking any other drug

When using this product • drowsiness will occur • avoid alcoholic drinks • do not drive a motor vehicle or operate machinery • take with food or milk. If stomach upset occurs

Stop use and ask a doctor if • you experience any of the following signs of stomach bleeding • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better • you have symptoms of heart problems or stroke • chest pain • trouble breathing • weakness in one part or side of body • slurred speech • leg swelling • pain gets worse or lasts more than 10 days • sleeplessness persists continuously for more than 2 weeks
 Insomnia may be a symptom of a serious underlying medical illness. • redness or swelling is present in the painful area
 • any new symptoms appear • you have difficulty swallowing • it feels like the pill is stuck in your throat

Drug Facts (continued)

If pregnant or breast feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions • do not take more than directed • drink a full glass of water with each dose • adults and children 12 years and over: take 2 caplets at bedtime • do not take more than 2 caplets in 24 hours
 • taken with food, this product may take longer to work

Other information • read all warnings and directions before use. Keep outer carton. • each caplet contains: sodium 20 mg • store at 20-25°C (68-77°F) • avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients camauba wax, FD&C blue #2 aluminum lake, hydroxytoluene, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, purified water, talc, titanium dioxide

Questions or comments?
 1 800 395 0689 (Mon-Fri 9AM-5PM EST)


Bayer

XXXXXXXXXX
 0000
 5 90353 25866

Do not use if carton is open or if foil seal imprinted with "SAFETY SQUEASE" on bottle opening is missing or broken

Dist. by: **Bayer HealthCare LLC**, Whippany, NJ 07981
 Safety Squeeze is a registered trademark of The Procter & Gamble Company. Used with permission.

MADE IN GERMANY
 LOT/EXP





Do not use if foil seal imprinted with "SAFETY SQUEASE" on bottle opening is missing or broken.

See Carton For Full Information. **Active ingredients (in each caplet)** Diphenhydramine hydrochloride 25 mg, Naproxen sodium 220 mg (naproxen 200 mg) **Warnings**

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin redness
- rash
- hives. If an allergic reaction occurs, stop use and seek medical

LIFT HERE

Help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart

failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer. **Ask a doctor or pharmacist before use if you are taking any other drug. If pregnant or breast-feeding, ask a health professional before use. Directions Do not take more than directed.** Adults and children 12 years and over: take 2 caplets with a full glass of water at bedtime. Do not take more than 2 caplets in 24 hours. **Questions or comments? 1-800-395-0689 (Mon-Fri 9AM - 5PM EST)**
Dist. by: Bayer HealthCare LLC, Whippany, NJ 07981



ALEVE[®] pm
 naproxen sodium, 220 mg (NSAID)/
 diphenhydramine HCl, 25 mg
 Pain reliever/nighttime sleep-aid

40 CAPLETS*
 *40 TABLETS

SLEEP AID PLUS
12 HOUR PAIN RELIEVING
STRENGTH OF ALEVE

Drug Facts

Active ingredients (in each caplet)
 Diphenhydramine hydrochloride 25 mg Nighttime sleep-aid
 Naproxen sodium 220 mg (naproxen 200 mg) (NSAID) Pain reliever
 *nonsteroidal anti-inflammatory drug

Purposes
 Pain reliever/nighttime sleep-aid

Uses
 • for relief of occasional sleeplessness when associated with minor aches and pains
 • helps you fall asleep and stay asleep

Warnings
Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 • hives • facial swelling
 • asthma (wheezing) • shock
 • skin redness • rash • blisters
 If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID which may cause severe stomach bleeding. The chance is higher if you:
 • are age 60 or older
 • have had stomach ulcers or bleeding problems
 • take a blood thinning (anticoagulant) or steroid drug
 • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 • have 3 or more alcoholic drinks every day while using this product.
 • take more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued)

Do not use
 • if you have ever had an allergic reaction to any other pain reliever/fever reducer
 • unless you have time for a full night's sleep
 • in children under 12 years of age
 • right before or after heart surgery
 • with any other product containing diphenhydramine, even one used on skin
 • if you have sleeplessness without pain

Ask a doctor before use if
 • stomach bleeding warning applies to you
 • you have problems or serious side effects from taking pain relievers or fever reducers
 • you have a history of stomach problems, such as heartburn
 • you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
 • you are taking a diuretic
 • you have glaucoma
 • you have trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are
 • taking sedatives or tranquilizers, or any other sleep-aid
 • under a doctor's care for any serious condition
 • taking any other antihistamines
 • taking any other drug

When using this product
 • drowsiness will occur
 • avoid alcoholic drinks
 • do not drive a motor vehicle or operate machinery
 • take with food or milk if stomach upset occurs

Stop use and ask a doctor if
 • you experience any of the following signs of stomach bleeding:
 • feel faint • vomit blood • have bloody or black stools
 • have stomach pain that does not get better
 • you have symptoms of heart problems or stroke:
 • chest pain • trouble breathing
 • weakness in one part or side of body
 • slurred speech • leg swelling
 • pain gets worse or lasts more than 10 days
 • sleeplessness persists continuously for more than 2 weeks.
 Insomnia may be a symptom of a serious underlying medical illness.
 • redness or swelling is present in the painful area

Drug Facts (continued)

• any new symptoms appear
 • you have difficulty swallowing
 • it feels like the pill is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 • do not take more than directed
 • drink a full glass of water with each dose
 • adults and children 12 years and over: take 2 caplets at bedtime
 • do not take more than 2 caplets in 24 hours
 • if taken with food, this product may take longer to work.

Other information
 • read all warnings and directions before use. **Keep out of carton.**
 • each caplet contains: sodium 20 mg
 • store at 20-25°C (68-77°F)
 • avoid high humidity and excessive heat, above 40°C (104°F)

Inactive ingredients: carnauba wax, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polydioxane, purified water, talc, titanium dioxide.

Questions or comments?
 1-800-985-0689 (Mon-Fri 8AM-5PM EST)

XXXXXXXXXX
 0000
 12 OF 5
 CODE AREA
 REQUIRES 1/4"
 QUART ZONE
 ←9/16"→

Do not use if carton is open or if foil seal imprinted with "SAFETY SQUEASE" on bottle opening is missing or broken.

Dist. by:
 Bayer HealthCare LLC
 Whippany, NJ 07981
MADE IN GERMANY
 LOT/EXP




See a Squeeze is a registered trademark of The Process & Control Company. Use with permission.

Bayer

LOT & EXP CODE AREA

CLOSE ANGLE WAVE RULE





SOFT GRIP® CAP

043101

ALEVE® pm

naproxen sodium 220 mg (NSAID) /
diphenhydramine HCl 25 mg
Pain relief/nighttime sleep-aid

SOFT GRIP® CAP

ALEVE® pm

naproxen sodium, 220 mg (NSAID) /
diphenhydramine HCl, 25 mg
Pain relief/nighttime sleep-aid

Drug Facts

Active ingredients
(In each caplet)
Diphenhydramine hydrochloride 25 mg Nighttime sleep-aid
Naproxen sodium 220 mg
(naproxen 200 mg (NSAID) Pain reliever
hydroxyzine hydrochloride 25 mg Pain reliever

Purposes

Uses

• for relief of occasional sleeplessness when associated with minor aches and pains
• helps you fall asleep and stay asleep

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

• hives • facial swelling • asthma (wheezing)
• shock • skin redness • rash • diarrhea

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID. NSAIDs cause severe stomach bleeding. The chance is higher if you:

• are age 60 or older
• have had stomach ulcers or bleeding problems
• take other drugs containing (anticoagulant) or steroid drug NSAIDs (aspirin, ibuprofen, naproxen, or others)
• have 3 or more alcoholic drinks every day while using this product

• take more or for a longer time than directed

Heart risks and stroke warning: NSAIDs, such as aspirin, ibuprofen, and naproxen, may increase the risk of heart disease and stroke. These risks are higher if you use more than directed or for longer than directed.

Drug Facts (continued)

• pain gets worse or lasts more than 10 days
• sleeplessness persists continuously for more than 2 weeks
• hives may be a symptom of a serious underlying medical illness
• redness or swelling is present in the painful area
• any new symptoms appear
• it feels like the pill is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. If you are pregnant, using this product during the last 3 months of pregnancy unless directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Directions more than directed
• drink full glass of water with each dose
• adults and children 12 years and over: take 2 caplets at bedtime
• do not take more than 2 caplets in 24 hours
• if taken with food, this product may take longer to work

Other information

• read all warnings and directions before use. Keep outer caplet container.
• each caplet contains: sodium 20 mg
• store at 20-25°C (68-77°F)
• avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients carmelum, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyox, purified water, talc, titanium dioxide

Questions or comments? 1-800-345-0889
(Mon. - Fri. 9AM - 5PM EST)

Drug Facts (continued)

Do not use
• if you have ever had an allergic reaction to any other pain reliever/fever reducer unless you have time for a full night's sleep
• in children under 12 years of age
• with any other product containing diphenhydramine, even one used on skin
• if you have sleeplessness without pain

Ask a doctor before use if

• you have bleeding warning applies to you
• you have problems or serious side effects from taking pain relievers or fever reducers
• you have high blood pressure, heart disease, liver cirrhosis, kidney

Ask a doctor or pharmacist before use if you are

• taking sedatives or tranquilizers, or any other sleep-aid
• taking any other antacids
• taking any other anticholinergics
• you are taking a diuretic
• you are taking a diuretic
• you have glaucoma
• you have a breathing problem such as emphysema or chronic bronchitis
• you have trouble urinating due to an enlarged prostate gland

When using this product

• avoid alcoholic drinks
• do not drive a motor vehicle or operate machinery
• take with food or milk. 1 stomach upset occurs

Stop use and ask a doctor if

• you experience any of the following signs of stomach bleeding:
• have faint • vomit blood • have bloody or black stools
• have stomach pain that does not go away
• you have chest pain • trouble breathing
• weakness in one part or side of body
• slurred speech • egg swelling

XXXXXXXXXX

0000



SOFT GRIP® CAP

ALEVE® pm

naproxen sodium 220 mg (NSAID) /
diphenhydramine HCl 25 mg
Pain relief/nighttime sleep-aid

SOFT GRIP® CAP

ALEVE® pm

naproxen sodium, 220 mg (NSAID) /
diphenhydramine HCl, 25 mg
Pain relief/nighttime sleep-aid



**SLEEP AID PLUS
12 HOUR PAIN RELIEVING
STRENGTH OF ALEVE**


40 CAPLETS*
*CAPSULE-SHAPED TABLETS

Do not use this product if printed safety seal bearing "Bayer HealthCare" under cap is torn or missing.

Dist. by:
Bayer Healthcare LLC
Whippany, NJ 07981

MADE IN GERMANY
Bayer, the Bayer cross, Aleve and Soft Grip are registered trademarks of Bayer.

Bayer



3 25866 56028 5



Do not use if foil seal imprinted with "SAFETY SQUEASE" on bottle opening is missing or broken.

See Carton For Full Information. **Active ingredients (in each caplet)** Diphenhydramine hydrochloride 25 mg, Naproxen sodium 220 mg (naproxen 200 mg) **Warnings**

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin redness
- rash
- hives. If an allergic reaction occurs, stop use and seek medical

LIFT HERE


Help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart

failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer. **Ask a doctor or pharmacist before use if you are taking any other drug. If pregnant or breast-feeding, ask a health professional before use. Directions Do not take more than directed.** Adults and children 12 years and over: take 2 caplets with a full glass of water at bedtime. Do not take more than 2 caplets in 24 hours. **Questions or comments? 1-800-395-0689 (Mon-Fri 9AM - 5PM EST)**
Dist. by: Bayer HealthCare LLC, Whippany, NJ 07981



Drug Facts

Active ingredients (in each caplet)
 Diphenhydramine hydrochloride 25 mg Nighttime sleep-aid
 Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)* Pain reliever
 *nonsteroidal anti-inflammatory drug

Purposes

Uses

- for relief of occasional sleeplessness when associated with minor aches and pains
- helps you fall asleep and stay asleep

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- asthma (wheezing)
- skin reddening
- blisters
- facial swelling
- shock
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- unless you have time for a full night's sleep
- in children under 12 years of age
- right before or after heart surgery
- with any other product containing diphenhydramine, even one used on skin
- if you have sleeplessness without pain

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have a breathing problem such as emphysema or chronic bronchitis
- you have glaucoma
- you have trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers, or any other sleep-aid
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery
- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vom t blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swe ling
 - pain gets worse or lasts more than 10 days
 - sleeplessness persists continuously for more than 2 weeks.

Insomnia may be a symptom of a serious underlying medical illness.

Drug Facts (continued)

- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the pill is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions


- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets at bedtime
- do not take more than 2 caplets in 24 hours
- if taken with food, this product may take longer to work

Other information

- read all warnings and directions before use. Keep outer carton.
- each caplet contains: sodium 20 mg
- store at 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients carnauba wax, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, purified water, talc, titanium dioxide

Questions? 1-800-398-0689




Do not use if carton is open or if foil seal imprinted with "SAFETY SQUEASE" on bottle opening is missing or broken.

Dist. by:
 Bayer HealthCare LLC
 Whippany, NJ 07981
MADE IN GERMANY



LOT/EXP

Safety Squease is a registered trademark of The Procter & Gamble Company. Used with permission.

Bayer







Drug Facts

Active ingredients (in each caplet) Purposes
 Diphenhydramine hydrochloride 25 mg.....Nighttime sleep-aid
 Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*.....Pain reliever
 *nonsteroidal anti-inflammatory drug

Uses

- for relief of occasional sleeplessness when associated with minor aches and pains
- helps you fall asleep and stay asleep

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- asthma (wheezing)
- skin reddening
- blisters
- facial swelling
- shock
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- unless you have time for a full night's sleep
- in children under 12 years of age
- right before or after heart surgery
- with any other product containing diphenhydramine, even one used on skin
- if you have sleeplessness without pain

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have a breathing problem such as emphysema or chronic bronchitis
- you have glaucoma
- you have trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers, or any other sleep-aid
- under a doctor's care for any serious condition
- taking any other antihistamines
- taking any other drug

When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery
- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness.
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the pill is stuck in your throat

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets at bedtime
- do not take more than 2 caplets in 24 hours
- if taken with food, this product may take longer to work


Other information

- read all warnings and directions before use. Keep outer carton.
- each caplet contains: sodium 20 mg
- store at 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients carnauba wax, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, purified water, talc, titanium dioxide

Questions or comments?
 1-800-395-0689 (Mon - Fri 9AM - 5PM EST)

Bayer




Do not use if carton is open or if foil seal imprinted with "SAFETY SQUEASE" on bottle opening is missing or broken.

Dist. by:
 Bayer HealthCare LLC
 Whippany, NJ 07981

Safety Squeeze is a registered trademark of The Procter & Gamble Company. Used with permission.

MADE IN GERMANY

LOT/EXP



TWO 80-COUNT
BOTTLES INSIDE

ALEVE[®] pm
naproxen sodium, 220 mg (NSAID)/
diphenhydramine HCl, 25 mg
Pain reliever/nighttime sleep-aid

ALEVE[®] pm
naproxen sodium, 220 mg (NSAID)/
diphenhydramine HCl, 25 mg
Pain reliever/nighttime sleep-aid



**SLEEP AID PLUS
12 HOUR PAIN RELIEVING
STRENGTH OF ALEVE**

160 CAPLETS^{*}
*CAPSULE-SHAPED TABLETS

LOT

EXP

Do not use if carton is open or if foil seal imprinted with
"SAFETY SEAL" on bottle opening is missing or broken.
Bayer HealthCare LLC
Whispery NJ 07981
MADE IN GERMANY
Bayer, the Bayer Cross and Aleve
are registered trademarks of Bayer.

Bayer



XXXXXX

Drug Facts

Active Ingredients (in each caplet)
Diphenhydramine hydrochloride 25 mg Nighttime sleep-aid
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)* Pain reliever
*nonsteroidal anti-inflammatory drug

Uses
• for relief of occasional sleeplessness when associated with minor aches and pains
• helps you fall asleep and stay asleep

Warnings
Allergy alert: Naproxen sodium may cause a severe allergic reaction especially in people allergic to aspirin. Symptoms may include:
• hives • facial swelling
• asthma (wheezing) • shock
• skin redness • rash
• blisters
If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID which may cause severe stomach bleeding. The chance is higher if you are age 60 or older
• have had stomach ulcers or bleeding problems
• take a blood thinning (anticoagulant) or steroid drug
• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
• have 3 or more alcohol drinks every day while using this product
• take more or for a longer time than directed

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use
• if you have ever had an allergic reaction to any other pain reliever/fever reducer
• unless you have time for a full night's sleep
• in children under 12 years of age
• right before or after heart surgery
• with any other product containing diphenhydramine, even one used on skin
• if you have sleeplessness without pain

Ask a doctor before use if:
• stomach bleeding warning applies to you
• you have problems or serious side effects from taking pain relievers or fever reducers
• you have a history of stomach problems, such as heartburn
• you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
• you are taking a diuretic
• you have a breathing problem such as emphysema or chronic bronchitis
• you have glaucoma
• you have trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are:
• taking sedatives or tranquilizers or any other sleep-aid under a doctor's care for any serious condition
• taking any other drug
• taking any other drug

When using this product
• drowsiness will occur
• avoid alcoholic drinks
• do not drive a motor vehicle or operate machinery
• take with food or milk if stomach upset occurs

Stop use and ask a doctor if:
• you experience any of the following signs of stomach bleeding:
• feel faint • vomit blood • have bloody or black stools
• have stomach pain that does not get better

Drug Facts (continued)

• you have symptoms of heart problems or stroke:
• chest pain • trouble breathing
• weakness in one part or side of body
• slurred speech • leg swelling
• pain gets worse or lasts more than 10 days
• sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness.
• redness or swelling is present in the painful area
• any new symptoms appear
• you have difficulty swallowing
• it feels like the pill is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
• do not take more than directed
• drink a full glass of water with each dose
• adults and children 12 years and over: take 2 caplets at bedtime
• do not take more than 2 caplets in 24 hours
• if taken with food, this product may take longer to work

Other information
• read all warnings and directions before use. Keep outer carton.
• each caplet contains: sodium 20 mg
• store at 20-25°C (68-77°F)
• avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients: carnauba wax, FD&C blue #2, aluminum lake, hygroscopic loss, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinylpyrrolidone, purified water, talc, titanium dioxide

Questions or comments? 1-800-395-0689 (Mon - Fri 9AM - 5PM EST)





Bayer, the Bayer Cross and Aleve are registered trademarks of Bayer.



16 FREE CAPLETS WITH THIS PURCHASE



**SLEEP AID PLUS
12 HOUR PAIN RELIEVING
STRENGTH OF ALEVE**

ALEVE[®] pm

naproxen sodium, 220 mg (NSAID)/
diphenhydramine HCl, 25 mg
Pain reliever/nighttime sleep-aid

96 CAPLETS*
*CAPSULES SHOWN TABLETS

Drug Facts

Active ingredients (in each caplet)
Diphenhydramine hydrochloride 25 mg.....Nighttime sleep-aid
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID).....Pain reliever
*nonsteroidal anti-inflammatory drug

Uses

- for relief of occasional sleeplessness when associated with minor aches and pains
- helps you fall asleep and stay asleep

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- asthma (wheezing)
- skin reddening
- blisters
- facial swelling
- shock
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- unless you have time for a full night's sleep
- in children under 12 years of age
- right before or after heart surgery
- with any other product containing diphenhydramine, even one used on skin
- if you have sleeplessness without pain

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have a breathing problem such as emphysema or chronic bronchitis
- you have glaucoma
- you have trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers, or any other sleep-aid
- under a doctor's care for any serious condition
- taking any other antihistamines
- taking any other drug

When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery
- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness.
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the pill is stuck in your throat

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets at bedtime
- do not take more than 2 caplets in 24 hours
- if taken with food, this product may take longer to work

Other information

- read all warnings and directions before use. Keep out carton.
- each caplet contains: sodium 20 mg
- store at 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients carnauba wax, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystaline cellulose, polyethylene glycol, povidone, purified water, talc, titanium dioxide

Questions or comments?
1-800-395-0689 (Mon - Fri 9AM - 5PM EST)

Bayer

XXXXXXXXXX



Do not use if carton is open or if foil seal imprinted with "SAFETY SQUEASE" on bottle opening is missing or broken.

Dist. by:
Bayer HealthCare LLC
Whippany, NJ 07981
MADE IN GERMANY
LOT/EXP

Safety Squeeze is a registered trademark of The Procter & Gamble Company. Used with permission.





This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

VALERIE S PRATT
09/12/2017