

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use PAROXETINE CAPSULES safely and effectively. See full prescribing information for PAROXETINE CAPSULES.

PAROXETINE capsules, for oral use

Initial U.S. Approval: 1992

WARNING: SUICIDAL THOUGHTS AND BEHAVIORS

See full prescribing information for complete boxed warning.

- **Potential for increased risk of suicidal thinking and behavior (5.1)**
- **Monitor for worsening and emergence of suicidal thoughts and behaviors (5.1)**

-----**RECENT MAJOR CHANGES**-----

Warnings and Precautions, Serotonin Syndrome and amphetamines (5.2) 01/2017

-----**INDICATIONS AND USAGE**-----

Paroxetine capsules are a selective serotonin reuptake inhibitor (SSRI) indicated for the treatment of moderate to severe vasomotor symptoms associated with menopause (VMS) (1)
Limitation of Use: Paroxetine capsules are not indicated for the treatment of any psychiatric condition (1)

-----**DOSAGE AND ADMINISTRATION**-----

The recommended dosage of paroxetine capsules is 7.5 mg once daily, at bedtime (2.1)

-----**DOSAGE FORMS AND STRENGTHS**-----

Capsules: 7.5 mg (3)

-----**CONTRAINDICATIONS**-----

- Concurrent use with monoamine oxidase inhibitors (MAOI) or use within 14 days of MAOI use (2.2, 4.1, 5.2, 7.3)
- Use with thioridazine (4.2, 7.1)
- Use with pimozide (4.3, 7.1)
- Hypersensitivity to any ingredient in paroxetine capsules (4.4)
- Pregnancy (4.5, 8.1)

-----**WARNINGS AND PRECAUTIONS**-----

- *Suicidality:* Monitor for suicidality or unusual changes in behavior (5.1)

- *Serotonin Syndrome* Serotonin syndrome, which is potentially life-threatening, has been reported with SSRIs. Discontinue paroxetine and initiate supportive treatment (5.2, 7.3)
- *Tamoxifen* Efficacy of tamoxifen may be reduced when administered concomitantly with paroxetine (5.3, 7.1)
- *Abnormal Bleeding* Caution patients about the risk of bleeding associated with the concomitant use of paroxetine and non-steroidal anti-inflammatory drugs (NSAIDs), aspirin, or other drugs that affect coagulation (5.4, 7.1)
- *Angle-Closure Glaucoma* Angle-closure glaucoma has occurred in patients who have untreated anatomically narrow angles and who are treated with antidepressants. (5.5)
- *Hyponatremia* Can occur in association with syndrome of inappropriate antidiuretic hormone secretion (SIADH) (5.6)
- *Bone Fracture:* Epidemiological studies have reported an association between SSRI treatment and fractures (5.7)
- *Activation of Mania/Hypomania* Screen for bipolar disorder and monitor for mania/hypomania (5.8)
- *Seizures:* Use cautiously in patients with a history of seizures or with conditions that potentially lower the seizure threshold (5.9)
- *Akathisia* Can occur, most likely in the first few weeks of treatment (5.10)
- *Cognitive and Motor Impairment* May cause impairment; patients should not operate machinery or motor vehicles until certain that paroxetine does not affect them adversely (5.11)

-----**ADVERSE REACTIONS**-----

The most common adverse reactions (greater than or equal to 2%) reported in clinical trials were: headache, fatigue, and nausea/vomiting (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Actavis at 1-800-272-5525 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch

-----**DRUG INTERACTIONS**-----

- Paroxetine is a strong CYP2D6 inhibitor. Coadministration of paroxetine can alter concentrations of other drugs that are metabolized by CYP2D6. Consider potential drug interactions prior to and during therapy (5.3, 7.1, 7.3). See Full Prescribing Information for a list of clinically significant drug interactions (7.1, 7.2, 7.3)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide

Revised: 5/2017

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: SUICIDAL THOUGHTS AND BEHAVIORS

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Dosage Information
- 2.2 Use of Paroxetine Capsules Before or After a Monoamine Oxidase Inhibitor (MAOI)

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

- 4.1 Monoamine Oxidase Inhibitors
- 4.2 Thioridazine
- 4.3 Pimozide
- 4.4 Hypersensitivity to any Ingredient in Paroxetine Capsules
- 4.5 Pregnancy

5 WARNINGS AND PRECAUTIONS

- 5.1 Suicidal Thoughts and Behaviors
- 5.2 Serotonin Syndrome
- 5.3 Potential Impact on Tamoxifen Efficacy
- 5.4 Abnormal Bleeding
- 5.5 Angle-Closure Glaucoma
- 5.6 Hyponatremia
- 5.7 Bone Fracture
- 5.8 Screening Patients for Bipolar Disorder and Monitoring for Mania/Hypomania
- 5.9 Seizures
- 5.10 Akathisia
- 5.11 Potential for Cognitive and Motor Impairment

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

7 DRUG INTERACTIONS

- 7.1 Potential for Paroxetine to Affect Other Drugs
- 7.2 Potential for Other Drugs to Affect Paroxetine
- 7.3 Other Potentially Significant Drug Interactions

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment
- 8.7 Hepatic Impairment

10 OVERDOSAGE

- 10.1 Human Experience with Overdosage
- 10.2 Management of Overdosage

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from Full Prescribing Information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: SUICIDAL THOUGHTS AND BEHAVIORS

Antidepressants, including selective serotonin reuptake inhibitors (SSRIs), have been shown to increase the risk of suicidal thoughts and behavior in pediatric and young adult patients when used to treat major depressive disorder and other psychiatric disorders. Because paroxetine is an SSRI, monitor patients closely for worsening and for emergence of suicidal thoughts and behaviors. Advise families and caregivers of the need for close observation and communication with the prescriber [see Warnings and Precautions (5.1)].

1 INDICATIONS AND USAGE

Paroxetine capsules are indicated for the treatment of moderate to severe vasomotor symptoms (VMS) associated with menopause.

Limitation of Use:

Paroxetine capsules are not indicated for the treatment of any psychiatric condition. Paroxetine capsules contain a lower dose of paroxetine than that used to treat depression, obsessive compulsive disorder, panic disorder, generalized anxiety disorder, social anxiety disorder, and post-traumatic stress disorder. The safety and efficacy of this lower dose of paroxetine in paroxetine capsules have not been established for any psychiatric condition. Patients who require paroxetine for treatment of a psychiatric condition should discontinue paroxetine capsules and initiate a paroxetine-containing medication that is indicated for such use.

2 DOSAGE AND ADMINISTRATION

2.1 Dosage Information

The recommended dosage of paroxetine capsules for the treatment of moderate to severe VMS is 7.5 mg once daily, at bedtime, with or without food.

2.2 Use of Paroxetine Capsules Before or After a Monoamine Oxidase Inhibitor (MAOI)

Wait at least 14 days after discontinuation of an MAOI before initiating therapy with paroxetine capsules. Conversely, allow at least 14 days after stopping paroxetine capsules before starting an MAOI [see *Contraindications (4.1), Warnings and Precautions (5.2) and Drug Interactions (7.3)*].

3 DOSAGE FORMS AND STRENGTHS

Paroxetine capsules are available as 7.5 mg in a size “2”, hard gelatin capsule with an orange opaque cap printed with “A401” and an orange opaque body printed with “7.5 mg” in black ink, containing a white to off white blend.

4 CONTRAINDICATIONS

4.1 Monoamine Oxidase Inhibitors

Concomitant use of an MAOI with paroxetine capsules or within 14 days of stopping treatment with paroxetine capsules is contraindicated because of an increased risk of serotonin syndrome. The use of paroxetine capsules within 14 days of stopping an MAOI is also contraindicated [see *Dosage and Administration (2.2), Warnings and Precautions (5.2) and Drug Interactions (7.3)*].

Starting paroxetine capsules in a patient who is being treated with linezolid or intravenous methylene blue, both of which inhibit monoamine oxidase, is also contraindicated because of an increased risk of serotonin syndrome [see *Dosage and Administration (2.2), Warnings and Precautions (5.2) and Drug Interactions (7.3)*].

4.2 Thioridazine

Concomitant use of paroxetine capsules with thioridazine is contraindicated, because thioridazine prolongs

the QT interval, and paroxetine can increase thioridazine levels [*see Drug Interactions (7.1)*].

4.3 Pimozide

Concomitant use of paroxetine capsules with pimozide is contraindicated because pimozide prolongs the QT interval, and paroxetine increases pimozide levels [*see Drug Interactions (7.1)*].

4.4 Hypersensitivity to any Ingredient in Paroxetine Capsules

Paroxetine capsules are contraindicated in patients with a history of hypersensitivity to paroxetine or any of the other ingredients in paroxetine capsules.

4.5 Pregnancy

Menopausal VMS does not occur during pregnancy and paroxetine capsules may cause fetal harm [*see Use in Specific Populations (8.1)*].

5 WARNINGS AND PRECAUTIONS

5.1 Suicidal Thoughts and Behaviors

Paroxetine is not approved for any psychiatric condition.

Antidepressants, including those that contain an SSRI, increase the risk of suicidal thinking and behavior (suicidality) in pediatric and young adult patients when used to treat major depressive disorder (MDD) and other psychiatric disorders. There is limited information regarding suicidality in women who use paroxetine for treatment of VMS. The paroxetine trials excluded women with a presence or history of previous psychiatric disorders.

Consider discontinuing paroxetine in patients with worsening depression or those who experience emergent suicidality or symptoms that might be precursors to worsening depression or suicidality, especially if these symptoms are severe, abrupt in onset, or were not part of the patient's presenting symptoms.

All patients being treated with paroxetine should be observed closely for clinical worsening, suicidality, and unusual changes in behavior, especially during the initial few months of treatment.

Anxiety, agitation, panic attacks, insomnia, irritability, hostility, aggressiveness, impulsivity, akathisia (psychomotor restlessness), hypomania, and mania have been reported in patients being treated with antidepressants for MDD as well as for other psychiatric and nonpsychiatric indications. Although a causal link between the emergence of such symptoms and either the worsening of depression and/or the emergence of suicidal impulses has not been established, there is concern that such symptoms may represent precursors to emerging suicidality.

Families and caregivers of patients being treated with paroxetine should be alerted about the need to monitor patients for the emergence of agitation, irritability, unusual changes in behavior, and the other symptoms described above, as well as the emergence of suicidality, and to report such symptoms immediately to healthcare providers.

5.2 Serotonin Syndrome

The development of a potentially life-threatening serotonin syndrome has been reported with SSRIs, including paroxetine, alone but particularly with concomitant use of serotonergic drugs (including triptans, tricyclic antidepressants, fentanyl, lithium, tramadol, tryptophan, buspirone, amphetamines, and St. John's Wort), and with drugs that impair metabolism of serotonin (in particular, MAOIs, both those intended to treat depression and others such as linezolid and intravenous methylene blue).

Serotonin syndrome symptoms may include mental status changes (e.g., agitation, hallucinations, delirium,

and coma), autonomic instability (e.g., tachycardia, labile blood pressure, dizziness, diaphoresis, flushing, hyperthermia), neuromuscular symptoms (e.g., tremor, rigidity, myoclonus, hyperreflexia, incoordination), and/or gastrointestinal symptoms (e.g., nausea, vomiting, diarrhea). Monitor patients for the emergence of serotonin syndrome.

The concomitant use of paroxetine with MAOIs is contraindicated. Do not start paroxetine in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue. All reports with methylene blue that provided information on the route of administration involved intravenous administration in the dose range of 1 mg/kg to 8 mg/kg. No reports involved the administration of methylene blue by other routes (such as oral tablets or local tissue injection) or at lower doses. There may be circumstances when it is necessary to initiate treatment with an MAOI such as linezolid or intravenous methylene blue in a patient taking paroxetine. Paroxetine should be discontinued before initiating treatment with the MAOI [*see Contraindications (4.1) and Dosage and Administration (2.2)*].

If concomitant use of paroxetine with other serotonergic drugs (e.g., triptans, tricyclic antidepressants, fentanyl, lithium, tramadol, tryptophan, buspirone, amphetamines, and St. John's Wort) is clinically warranted, consider the increased risk of serotonin syndrome and carefully observe the patient, particularly during treatment initiation [*see Contraindications (4.1) Drug Interactions (7.3)*].

Discontinue paroxetine and any concomitant serotonergic agents immediately if the above events occur and initiate supportive symptomatic treatment.

5.3 Potential Impact on Tamoxifen Efficacy

It is uncertain whether the coadministration of paroxetine and tamoxifen has a significant adverse effect on the efficacy of tamoxifen. Some studies have shown that the efficacy of tamoxifen, as measured by the risk of breast cancer relapse/mortality, may be reduced when co-prescribed with paroxetine as a result of paroxetine's irreversible inhibition of CYP2D6 [*see Drug Interactions (7.1)*]. However, other studies have failed to demonstrate such a risk. When tamoxifen is used for the treatment or prevention of breast cancer, weigh the likely benefit of paroxetine for treating VMS vs. the risk of possible decreased tamoxifen effectiveness, and consider avoiding the concomitant use of paroxetine for VMS treatment.

5.4 Abnormal Bleeding

SSRIs, including paroxetine, may increase the risk of bleeding events. Concomitant use of aspirin, nonsteroidal anti-inflammatory drugs (NSAIDs), warfarin, and other anticoagulants may add to this risk. Case reports and epidemiological studies (case-control and cohort design) have demonstrated an association between use of drugs that interfere with serotonin reuptake and the occurrence of gastrointestinal bleeding. Bleeding events related to SSRIs have ranged from ecchymosis, hematoma, epistaxis, and petechiae to life-threatening hemorrhages. Caution patients about the risk of bleeding associated with the concomitant use of paroxetine and NSAIDs, aspirin, or other drugs that affect coagulation [*see Drug Interactions (7.1)*].

5.5 Angle-Closure Glaucoma

The pupillary dilation that occurs following use of many antidepressants and paroxetine may trigger an angle-closure attack in a patient with anatomically narrow angles who does not have a patent iridectomy.

5.6 Hyponatremia

Hyponatremia may occur as a result of treatment with SSRIs, including paroxetine. Elderly patients may be at greater risk. In many cases, the hyponatremia appears to be the result of the syndrome of inappropriate antidiuretic hormone secretion (SIADH). Cases with serum sodium lower than 110 mmol/L have been reported in patients using SSRIs. Also, patients taking diuretics or who are volume-depleted can be at greater risk. Consider discontinuation of paroxetine in patients with symptomatic hyponatremia and institute appropriate medical intervention.

Signs and symptoms of hyponatremia include headache, difficulty concentrating, memory impairment, confusion, weakness, and unsteadiness, which can lead to falls. Signs and symptoms associated with more severe and/or acute cases have included hallucination, syncope, seizure, coma, respiratory arrest, and death.

5.7 Bone Fracture

Epidemiological studies on bone fracture risk following exposure to SSRIs have reported an association between SSRI treatment and fractures. It is unknown to what extent fracture risk is directly attributable to SSRI treatment. If a paroxetine-treated patient presents with unexplained bone pain, point tenderness, swelling, or bruising, consider the possibility of a fragility fracture.

5.8 Screening Patients for Bipolar Disorder and Monitoring for Mania/Hypomania

Paroxetine is only indicated for the treatment of moderate to severe VMS and is not approved for use in treating either depression or bipolar depression. However, prior to initiating treatment with paroxetine, all patients should be adequately screened to determine if they are at risk for bipolar disorder; such screening should include a detailed psychiatric history, including a family history of suicide, bipolar disorder, and depression. It is generally believed (though not established in controlled trials) that use of an antidepressant alone may increase the likelihood of precipitation of a mixed/manic episode in patients at risk for bipolar disorder.

5.9 Seizures

In premarketing testing of paroxetine, seizures occurred in 0.1% of paroxetine-treated patients. Use paroxetine cautiously in patients with a history of seizures or with conditions that potentially lower the seizure threshold. Evaluate and consider discontinuing use in any patient who develops seizures.

5.10 Akathisia

The use of paroxetine or other SSRIs has been associated with the development of akathisia, which is characterized by an inner sense of restlessness and psychomotor agitation such as an inability to sit or stand still usually associated with subjective distress. This is most likely to occur within the first few weeks of treatment. Discontinue treatment with paroxetine if akathisia occurs.

5.11 Potential for Cognitive and Motor Impairment

Paroxetine has the potential to impair judgment, thinking, or motor skills. Patients should be cautioned about operating hazardous machinery, including motor vehicles, until they are reasonably certain that the drug treatment does not affect them adversely.

6 ADVERSE REACTIONS

The following serious adverse reactions are discussed elsewhere in labeling:

- Suicidality [*see Warnings and Precautions (5.1)*]
- Serotonin syndrome [*see Warnings and Precautions (5.2)*]
- Abnormal bleeding [*see Warnings and Precautions (5.4)*]
- Angle-Closure Glaucoma [*see Warnings and Precautions (5.5)*]
- Hyponatremia [*see Warnings and Precautions (5.6)*]
- Bone Fracture [*see Warnings and Precautions (5.7)*]
- Mania/Hypomania [*see Warnings and Precautions (5.8)*]
- Seizure [*see Warnings and Precautions (5.9)*]
- Akathisia [*see Warnings and Precautions (5.10)*]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the

clinical trials of a drug cannot directly be compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The data described below reflect exposure to paroxetine in the one 8-week Phase 2 randomized, placebo-controlled trial and the two Phase 3 randomized, placebo-controlled, 12-week and 24-week trials for the treatment of moderate to severe VMS [see *Clinical Studies (14)*]. In these trials, a total of 635 women were exposed to paroxetine 7.5 mg administered orally once daily and 641 women received placebo. The majority of paroxetine-treated patients were Caucasian (68%) and African American (30%), with a mean age of 55 years (range 40 to 73 years). Women with a history of suicidal ideation or suicidal behavior were excluded from these studies.

Adverse Reactions Leading to Study Discontinuation: A total of 4.7% of women taking paroxetine discontinued from the clinical trials due to an adverse reaction, compared to 3.7% of women on placebo; the most frequent adverse reactions leading to discontinuation among paroxetine-treated women were: abdominal pain (0.3%), attention disturbances (0.3%), headache (0.3%), and suicidal ideation (0.3%).

Common Adverse Reactions: Overall, based on investigators' determinations about what events were likely to be drug-related, about 20% of women treated with paroxetine reported at least 1 adverse reaction in the three controlled studies. The most common adverse reactions (greater than or equal to 2% and more common among paroxetine-treated women) reported in these studies were headache, fatigue/malaise/lethargy, and nausea/vomiting. Of these commonly reported adverse reactions, nausea occurred primarily within the first 4 weeks of treatment and fatigue occurred primarily within the first week of treatment, and decreased in frequency with continued therapy.

The adverse reactions that occurred in at least 2% of patients in the paroxetine group and at a higher incidence than placebo are shown in Table 1 for the pooled Phase 2 and Phase 3 trials.

Table 1 Frequency of Adverse Reactions in the Phase 2 and Phase 3 Trials ($\geq 2\%$ and at a higher incidence than placebo)

	Frequency n (%)	
	Paroxetine (n = 635)	Placebo (n = 641)
Nervous system disorders		
Headache	40 (6.3)	31 (4.8)
General disorders and administration site conditions		
Fatigue, malaise, lethargy	31 (4.9)	18 (2.8)
Gastrointestinal disorders		
Nausea, vomiting	27 (4.3)	15 (2.3)

Certain symptoms were seen more frequently in women at the time of discontinuation of paroxetine compared to women discontinuing placebo, and have also been reported upon discontinuation of other formulations of paroxetine, particularly when abrupt. These include increased dreaming/nightmares, muscle cramps/spasms/twitching, headache, nervousness/anxiety, fatigue/tiredness, restless feeling in legs, and trouble sleeping/insomnia. While these events are generally self-limiting, there have been reports of serious discontinuation symptoms with other formulations of paroxetine.

Serious Adverse Reactions: In the pooled Phase 2 and Phase 3 trials, three paroxetine-treated patients reported a serious adverse reaction of suicidal ideation and one paroxetine-treated patient reported a serious adverse reaction of suicide attempt. There were no serious adverse reactions of suicidal ideation or suicide attempt reported among the placebo-treated patients.

6.2 Postmarketing Experience

The following adverse reactions have been identified from clinical studies of paroxetine and during post-approval use of other formulations of paroxetine. Because some of these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Blood and Lymphatic System Disorders: Idiopathic thrombocytopenic purpura, Events related to impaired hematopoiesis (including aplastic anemia, pancytopenia, bone marrow aplasia, agranulocytosis).

Cardiac Disorders: Atrial fibrillation, Pulmonary edema, Ventricular fibrillation, Ventricular tachycardia (including torsades de pointes).

Gastrointestinal Disorders: Pancreatitis, Pancreatitis hemorrhagic, Vomiting.

General Disorders and Administration Site Conditions: Death, Drug withdrawal syndrome, Malaise.

Hepatobiliary Disorders: Drug-induced liver injury, Hepatic failure, Jaundice.

Immune System Disorders: Anaphylactoid reaction, Angioedema, Toxic epidermal necrolysis.

Investigations: Elevated liver tests (the most severe cases were deaths due to liver necrosis, and grossly elevated transaminases associated with severe liver dysfunction).

Metabolism and Nutrition Disorders: Diabetes mellitus inadequate control, Type 2 diabetes mellitus.

Nervous System Disorders: Neuroleptic malignant syndrome, Paresthesia, Somnolence, Tremor.

Psychiatric Disorders: Aggression, Agitation, Anxiety, Confusional state, Depression, Disorientation, Homicidal ideation, Insomnia, Restlessness.

Respiratory, Thoracic and Mediastinal Disorders: Pulmonary hypertension.

Skin and Subcutaneous Tissue Disorders: Hyperhidrosis, Stevens-Johnson syndrome.

7 DRUG INTERACTIONS

No drug-drug interaction studies have been conducted with paroxetine.

7.1 Potential for Paroxetine to Affect Other Drugs

Paroxetine is a strong CYP2D6 inhibitor. Clinical drug interaction studies have been performed with substrates of CYP2D6 and show that paroxetine can inhibit the metabolism of drugs metabolized by CYP2D6 [see *Clinical Pharmacology (12.3)*]. Table 2 contains examples of drugs with a metabolism that may be affected by coadministration with paroxetine.

Table 2 Effects of Paroxetine on Other Drugs

Concomitant Drug Name	Effect of Paroxetine on Other Drugs	Clinical Recommendations
Thioridazine	Increased plasma concentrations of thioridazine Potential QTc prolongation	Concomitant use of thioridazine and paroxetine is contraindicated.
Pimozide	Increased plasma concentrations of pimozide. Potential QTc prolongation	Concomitant use of pimozide and paroxetine is contraindicated.
Tamoxifen	Reduced plasma concentrations of active tamoxifen metabolite	Consider avoiding concomitant use of tamoxifen and paroxetine.
Tricyclic Antidepressant (TCA) (e.g., Desipramine)	Increased plasma concentrations and elimination half-life	Plasma TCA concentrations may need to be monitored and the dose of TCA may need to be reduced if a TCA is coadministered with paroxetine. Monitor tolerability.
Risperidone	Increased plasma concentrations of risperidone	A lower dosage of risperidone may be necessary (see the Full Prescribing Information for risperidone). Monitor tolerability.
Atomoxetine	Increased exposure of atomoxetine	A lower dosage of atomoxetine may be necessary (see Full Prescribing Information for atomoxetine). Monitor tolerability.
Drugs Highly Bound to Plasma Protein (e.g., Warfarin)	Increased free plasma concentrations	The dosage of warfarin may need to be reduced. Monitor tolerability and the International Normalized Ratio.
Digoxin	Decreased plasma concentrations of digoxin	Dosage of digoxin may need to be increased. Monitor digoxin concentrations and clinical effect.
Theophylline	Increased plasma concentrations of theophylline	Dosage of theophylline may need to be decreased. Monitor theophylline concentrations and tolerability.

Use caution if coadministering paroxetine with other drugs that are metabolized by CYP2D6, including nortriptyline, amitriptyline, imipramine, desipramine, fluoxetine, phenothiazines, risperidone, and Type 1C antiarrhythmics (e.g., propafenone, flecainide, and encainide).

7.2 Potential for Other Drugs to Affect Paroxetine

The metabolism and pharmacokinetics of paroxetine may be affected by the induction and inhibition of drug metabolizing enzymes such as CYP2D6. Table 3 contains a list of drugs that may affect the pharmacokinetics

of paroxetine when administered concomitantly [see *Clinical Pharmacology (12.3)*].

Table 3 Effects of Other Drugs on Paroxetine

Concomitant Drug Name	Effect of Concomitant Drug on Paroxetine	Clinical Recommendations
Phenobarbital	Decreased paroxetine exposure	No dose adjustment for paroxetine. Monitor clinical effect of paroxetine.
Phenytoin	Decreased paroxetine exposure	
Fosamprenavir/ Ritonavir	Decreased plasma concentration of paroxetine	
Cimetidine	Increased plasma concentration of paroxetine	

Use caution if coadministering paroxetine with other drugs that inhibit CYP2D6 (e.g., quinidine).

7.3 Other Potentially Significant Drug Interactions

Monoamine Oxidase Inhibitors (MAOIs)

Serious adverse reactions such as serotonin syndrome have been reported in patients receiving a concomitant SSRI and MAOI, in patients started on an SSRI who recently received an MAOI and in patients started on an MAOI who recently received an SSRI. Therefore, concomitant use of MAOIs with paroxetine or use of paroxetine and an MAOI within 14 days of each other is contraindicated [see *Dosage and Administration (2.2)*, *Contraindications (4.1)* and *Warnings and Precautions (5.2)*].

Serotonergic Drugs

If concomitant use of paroxetine with other serotonergic drugs (e.g., triptans, tricyclic antidepressants, fentanyl, lithium, tramadol, tryptophan, buspirone, amphetamines, and St. John's Wort) is clinically warranted, consider the increased risk of serotonin syndrome and carefully observe the patient, particularly during treatment initiation [see *Warnings and Precautions (5.2)*].

An interaction between paroxetine and tryptophan may occur when they are coadministered. Adverse experiences, consisting primarily of headache, nausea, sweating, and dizziness, have been reported when tryptophan was administered to patients taking paroxetine. Consequently, concomitant use of paroxetine with tryptophan is not recommended.

If concomitant use of paroxetine with a serotonergic drug is warranted, carefully observe the patient, particularly during treatment initiation. There have been postmarketing reports of serotonin syndrome with the use of an SSRI and a triptan.

Paroxetine capsules contain paroxetine, which is also the active ingredient in other drugs. The concomitant use of paroxetine with other paroxetine products is not recommended [see *Indications and Usage (1)*].

Drugs that Interfere with Hemostasis (e.g., NSAIDs, Aspirin, and Warfarin)

Altered anticoagulant effects, including increased bleeding, have been reported when SSRIs are coadministered with NSAIDs, aspirin, and warfarin or other drugs that affect coagulation. There may be a pharmacodynamic interaction between paroxetine and warfarin that causes an increased bleeding diathesis despite unaltered prothrombin time. Carefully monitor patients receiving warfarin therapy when paroxetine is initiated or discontinued [see *Warnings and Precautions (5.4)*].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category X

Risk Summary

Paroxetine is contraindicated in pregnant women because menopausal VMS does not occur during pregnancy and paroxetine can cause fetal harm. Epidemiological studies have shown that infants exposed to paroxetine in the first trimester of pregnancy may have an increased risk of cardiovascular malformations. Cardiac malformations are a common congenital abnormality. These data would suggest that the risk of a cardiac abnormality following paroxetine exposure in the first trimester may increase the risk from 1% to 2%. Exposure to SSRIs in late pregnancy may lead to an increased risk for neonatal complications requiring prolonged hospitalization, respiratory support, and tube feeding, and/or persistent pulmonary hypertension of the newborn (PPHN). No teratogenicity was seen in reproductive development studies conducted in rats and rabbits. However, an increase in rat pup deaths was seen during the first 4 days of lactation when dosing occurred during the last trimester of gestation and continued throughout lactation, at a dose approximately equal to the maximum recommended human dose (MRHD) for VMS (7.5 mg) on an mg/m² basis. If this drug is used during pregnancy, or if the patient becomes pregnant while taking this drug, the patient should be apprised of the potential hazard to a fetus.

Human Data

First-Trimester Pregnancy Exposure

- Epidemiologic studies which include data from the Swedish National Registry, a retrospective cohort study using United Healthcare data and a meta-analysis of studies (1992 to 2008) have shown a less than 2-fold increased risk of cardiac malformations, primarily ventricular septal and atrial septal defects, with first-trimester paroxetine exposure. Two case-control studies using separate databases with greater than 9000 birth defect cases and greater than 4000 controls showed 7 and 6 paroxetine-exposed infants respectively, with right ventricular outflow tract obstructions, a 2- to 3-fold increased risk. An increase in overall congenital malformations with first-trimester paroxetine use was not observed in all studies.

Third-Trimester Pregnancy Exposure

- Neonates exposed to SSRIs late in the third trimester have developed complications requiring prolonged hospitalization, respiratory support, and tube feeding. Such complications can arise immediately upon delivery. Reported clinical findings have included respiratory distress, cyanosis, apnea, seizures, temperature instability, feeding difficulty, vomiting, hypoglycemia, hypotonia, hypertonia, hyperreflexia, tremor, jitteriness, irritability, and constant crying. These features are consistent with either a direct toxic effect of SSRIs or, possibly, a drug discontinuation syndrome. It should be noted that in some cases, the clinical picture is consistent with serotonin syndrome [*see Warnings and Precautions (5.2)*].
- Infants exposed to SSRIs in late pregnancy may have an increased risk for persistent pulmonary hypertension of the newborn (PPHN). PPHN occurs in 1 to 2 per 1000 live births in the general population and is associated with substantial neonatal morbidity and mortality. In a retrospective case-control study of 377 women whose infants were born with PPHN and 836 women whose infants were born healthy, the risk for developing PPHN was approximately 6-fold higher for infants exposed to SSRIs after the 20th week of gestation compared to infants who had not been exposed to antidepressants during pregnancy. There is currently no corroborative evidence regarding the risk for PPHN following exposure to SSRIs in pregnancy; this is the first study that has investigated the potential risk. The study did not include enough cases with exposure to individual SSRIs to determine if all SSRIs posed similar levels of PPHN risk.

Animal Data

Reproduction studies were performed at doses up to 50 mg/kg/day in rats and 6 mg/kg/day in rabbits administered during organogenesis. These doses are approximately 65 (rat) and 16 (rabbit) times the maximum recommended human dose (MRHD) for VMS on an mg/m² basis. There were no teratogenic effects. However,

in rats, there was an increase in pup deaths during the first 4 days of lactation when dosing occurred during the last trimester of gestation and continued throughout lactation. This effect occurred at a dose of 1 mg/kg/day or approximately equal to the MRHD for VMS on an mg/m² basis. The no-effect dose for rat pup mortality was not determined. The cause of these deaths is unknown.

8.3 Nursing Mothers

Paroxetine is excreted in human milk. Because of the potential for serious adverse reactions in nursing infants from paroxetine, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

8.4 Pediatric Use

Safety and effectiveness in pediatric patients have not been established; paroxetine is not indicated in the pediatric population.

8.5 Geriatric Use

Clinical studies of paroxetine did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Elderly patients may have elevated paroxetine plasma concentrations compared to younger patients. However, no paroxetine dose adjustment is considered necessary in elderly patients [*see Clinical Pharmacology (12.3)*].

SSRIs have been associated with cases of clinically significant hyponatremia in elderly patients, who may be at greater risk for this adverse event [*see Warnings and Precautions (5.6)*].

8.6 Renal Impairment

No paroxetine dose adjustment is considered necessary in patients with renal impairment [*see Clinical Pharmacology (12.3)*].

8.7 Hepatic Impairment

No paroxetine dose adjustment is considered necessary in patients with liver impairment [*see Clinical Pharmacology (12.3)*].

10 OVERDOSAGE

10.1 Human Experience with Overdosage

There is limited clinical experience with paroxetine overdosage in humans, as there were no overdoses reported in the clinical studies.

Spontaneous cases of deliberate or accidental overdosage during paroxetine treatment have been reported; some of these cases were fatal and some of the fatalities appeared to involve paroxetine alone. Of nonfatal cases with known outcome, most recovered without sequelae. The largest known ingestion involved 2000 mg of paroxetine (267 times the maximum recommended daily dose) in a patient who recovered.

Commonly reported adverse reactions associated with paroxetine overdosage include somnolence, coma, nausea, tremor, tachycardia, confusion, vomiting, and dizziness. Other notable signs and symptoms observed with overdoses involving paroxetine (alone or with other substances) include mydriasis, convulsions (including status epilepticus), ventricular dysrhythmias (including torsades de pointes), hypertension, aggressive reactions, syncope, hypotension, stupor, bradycardia, dystonia, rhabdomyolysis, symptoms of hepatic dysfunction (including hepatic failure, hepatic necrosis, jaundice, hepatitis, and hepatic steatosis), serotonin syndrome, manic reactions, myoclonus, acute renal failure, and urinary retention.

10.2 Management of Overdosage


Treatment should consist of those general measures employed in the management of overdosage with any

SSRI. Consult with a certified poison control center for up-to-date guidance and advice on treatment of overdose.

Ensure an adequate airway, oxygenation, and ventilation. Monitor cardiac rhythm and vital signs. General supportive and symptomatic measures are also recommended. Induction of emesis is not recommended. In managing overdose, consider the possibility of multiple drug involvement.

11 DESCRIPTION

Paroxetine is an orally administered selective serotonin reuptake inhibitor (SSRI) for the treatment of moderate to severe VMS associated with menopause. It is identified chemically as (-)-*trans*-4R-(4'-fluorophenyl)-3S-[(3',4'-methylenedioxyphenoxy)methyl]piperidine mesylate and has the molecular formula of $C_{19}H_{20}FNO_3 \cdot CH_3SO_3H$. The molecular weight is 425.5 (329.4 as free base). The structural formula is:



The mesylate salt of paroxetine is an odorless, off-white powder, having a melting point range of 147° to 150°C and a solubility of more than 1 g/mL in water.

Each orange capsule contains 9.69 mg paroxetine mesylate equivalent to 7.5 mg paroxetine base.

Inactive ingredients consist of: anhydrous citric acid, colloidal silicon dioxide, dibasic calcium phosphate anhydrous, FD&C Yellow #6, gelatin, iron oxide black, magnesium stearate, microcrystalline cellulose, potassium hydroxide, propylene glycol, shellac, sodium lauryl sulfate, sodium starch glycolate and titanium dioxide.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Nonclinical studies have shown that paroxetine is an SSRI. Paroxetine is not an estrogen, and its mechanism of action for the treatment of VMS is unknown.

12.2 Pharmacodynamics

Studies at clinically relevant doses in humans have demonstrated that paroxetine blocks the uptake of serotonin into human platelets. *In vitro* studies in animals also suggest that paroxetine is a selective inhibitor of neuronal serotonin reuptake and has weak effects on norepinephrine and dopamine neuronal reuptake. *In vitro* radioligand binding studies indicate that paroxetine has little affinity for muscarinic α_1 -, α_2 -, beta-adrenergic-, dopamine (D_2)-, 5-HT₁-, 5-HT₂-, and histamine (H_1)-receptors.

12.3 Pharmacokinetics

Absorption, Distribution, Metabolism and Excretion

Absorption

Paroxetine is completely absorbed after oral dosing of the mesylate salt. In a study in which healthy postmenopausal women (n=24) received paroxetine 7.5 mg capsules as a daily dose for 14 days, steady-state paroxetine concentrations were achieved by approximately 12 days of dosing for most subjects, although it may take substantially longer in an occasional patient. Peak concentrations were reached at a median of 6 hours (3 to 8 hours range). Steady-state mean values of C_{max} , C_{min} , and AUC_{0-last} were 13.10 ng/mL (CV 91%), 7.17 ng/mL (CV 99%), and 237 hr*ng/mL (CV 94%), respectively.

Steady-state AUC_{0-24} values were about 3 times those of AUC_{0-inf} following a single dose, indicating non-linear pharmacokinetics. Steady-state C_{max} values were approximately 5 times greater than those attained after a single dose and steady-state exposure based on AUC_{0-24} was about 10 times greater than AUC_{0-24} after a single dose.

The nonlinear kinetics and excess accumulation are due to the fact that CYP2D6, an enzyme that is in part responsible for paroxetine metabolism, is readily saturable.

The effects of food on the bioavailability of paroxetine were studied with paroxetine tablets at higher strength. AUC was only slightly increased (6%) when drug was administered with food but the C_{max} was 29% greater, while the time to reach peak plasma concentration decreased from 6.4 hours post-dosing to 4.9 hours. Paroxetine can be taken with or without food.

Distribution

Paroxetine distributes throughout the body, including the central nervous system, with only 1% remaining in the plasma.

Approximately 95% and 93% of paroxetine is bound to plasma protein at 100 ng/mL and 400 ng/mL, respectively. Under clinical conditions, paroxetine concentrations would normally be less than 100 ng/mL. Paroxetine does not alter the *in vitro* protein binding of phenytoin or warfarin.

Metabolism

Paroxetine is extensively metabolized after oral administration. The principal metabolites are polar and conjugated products of oxidation and methylation, which are readily cleared. Conjugates with glucuronic acid and sulfate predominate, and major metabolites have been isolated and identified. Data indicate that the metabolites have no more than 1/50 the potency of the parent compound at inhibiting serotonin uptake. The metabolism of paroxetine is accomplished in part by cytochrome CYP2D6. Saturation of this enzyme at clinical doses appears to account for the nonlinearity of paroxetine kinetics with increasing dose and increasing duration of treatment. The role of this enzyme in paroxetine metabolism also suggests potential drug-drug interactions [see *Drug Interactions (7)*]. At steady state, when the CYP2D6 pathway is essentially saturated, paroxetine clearance is governed by alternative P450 isozymes, which, unlike CYP2D6, show no evidence of saturation.

Excretion

Approximately 64% of a 30 mg oral solution of paroxetine was excreted in the urine with 2% as the parent compound and 62% as metabolites over a 10-day post-dosing period. About 36% of the dose was excreted in the feces (probably via the bile), mostly as metabolites and less than 1% as the parent compound over the 10-day post-dosing period.

Specific Populations

Renal and Liver Impairment

Increased plasma concentrations of paroxetine occur in subjects with renal and hepatic impairment. The mean plasma concentration in patients with creatinine clearance below 30 mL/min was approximately 4 times greater than seen in normal volunteers. Patients with creatinine clearance of 30 to 60 mL/min and patients with hepatic impairment had about a 2-fold increase in plasma concentrations (AUC , C_{max}). No paroxetine dose

adjustment is considered necessary in patients with renal or hepatic impairment.

Elderly Patients

In a multiple-dose study in the elderly at daily paroxetine doses of 20, 30, and 40 mg, C_{\min} concentrations were about 70% to 80% greater than the respective C_{\min} concentrations in nonelderly subjects. No paroxetine dose adjustment is considered necessary in elderly patients.

Drug Interaction Studies

Potential Effect of Paroxetine on Other Drugs

Drugs Metabolized by CYP3A4

An *in vivo* drug interaction study involving the coadministration under steady-state conditions of paroxetine and terfenadine, a substrate for cytochrome CYP3A4, revealed no effect of paroxetine on terfenadine pharmacokinetics. *In vitro* studies have shown ketoconazole, a potent inhibitor of CYP3A4 activity, to be at least 100 times more potent than paroxetine as an inhibitor of the metabolism of several substrates for CYP3A4, including astemizole, triazolam, and cyclosporine. Based on the assumption that the relationship between paroxetine's *in vitro* K_i and its lack of effect on terfenadine's *in vivo* clearance predicts its effect on other CYP3A4 substrates, paroxetine's extent of inhibition of CYP3A4 activity is not likely to be of clinical significance.

Drugs Metabolized by CYP2D6

Many drugs are metabolized by the cytochrome P450 isozyme CYP2D6. Like other agents that are metabolized by CYP2D6, paroxetine may significantly inhibit the activity of this isozyme. In most patients (greater than 90%), this CYP2D6 isozyme is saturated early during paroxetine dosing.

Specific studies investigating the effect of paroxetine on drugs metabolized by CYP2D6 are listed below:

Pimozide: Higher doses of paroxetine have been shown to elevate plasma levels of pimozide. In a controlled study of healthy volunteers, after paroxetine was titrated to 60 mg daily, coadministration of a single dose of 2 mg pimozide was associated with mean increases in pimozide AUC of 151% and C_{\max} of 62%, compared to pimozide administered alone [*see Drug Interactions (7.1)*].

Tamoxifen: It is uncertain whether the coadministration of paroxetine and tamoxifen has a significant adverse effect on the efficacy of tamoxifen. Some studies have shown that the efficacy of tamoxifen, as measured by the risk of breast cancer relapse/mortality, may be reduced when co-prescribed with paroxetine as a result of paroxetine's irreversible inhibition of CYP2D6. However, other studies have failed to demonstrate such a risk [*see Warnings and Precautions (5.3) and Drug Interactions (7.1)*].

Desipramine: In one study, daily dosing of paroxetine (20 mg once daily) under steady-state conditions increased single dose desipramine (100 mg) C_{\max} , AUC, and $T_{1/2}$ by an average of approximately 2-, 5-, and 3-fold, respectively [*see Drug Interactions (7.1)*].

Risperidone: Daily dosing of paroxetine 20 mg in patients stabilized on risperidone (4 to 8 mg/day), a CYP2D6 substrate, increased mean plasma concentrations of risperidone approximately 4-fold, decreased 9-hydroxyrisperidone concentrations approximately 10%, and increased concentrations of the active moiety (the sum of risperidone plus 9-hydroxyrisperidone) approximately 1.4-fold [*see Drug Interactions (7.1)*].

Atomoxetine: The effect of paroxetine on the pharmacokinetics of atomoxetine has been evaluated when both drugs were at steady state. In healthy volunteers who were extensive metabolizers of CYP2D6, paroxetine 20 mg daily was given in combination with 20 mg atomoxetine every 12 hours. This resulted in increases in steady-state atomoxetine AUC values that were 6- to 8-fold greater and in atomoxetine C_{\max} values that were

3- to 4-fold greater than when atomoxetine was given alone [see *Drug Interactions (7.1)*].

Digoxin: Mean digoxin AUC at steady state decreased by 15% in the presence of paroxetine [see *Drug Interactions (7.1)*].

Beta Blockers: In a study in which propranolol (80 mg twice daily) was dosed orally for 18 days, the steady-state plasma concentrations of propranolol were unaltered during coadministration with paroxetine (30 mg once daily) for the final 10 days. The effects of propranolol on paroxetine have not been evaluated.

Potential Effect of Other Drugs on Paroxetine

Concomitant use of paroxetine with other drugs that alter CYP enzymes activities including CYP2D6 may affect the plasma concentrations of paroxetine. Specific studies investigating the effect of other drugs on paroxetine are listed below:

Cimetidine: Cimetidine inhibits many cytochrome P450 enzymes. In a study in which paroxetine (30 mg once daily) was dosed orally for 4 weeks, steady-state plasma concentrations of paroxetine were increased by approximately 50% during coadministration with oral cimetidine (300 mg three times daily) for the final week [see *Drug Interactions (7.2)*].

Phenobarbital: Phenobarbital induces many cytochrome P450 enzymes. When a single oral 30 mg dose of paroxetine was administered at phenobarbital steady state (100 mg once daily for 14 days), paroxetine AUC and $T_{1/2}$ were reduced (by an average of 25% and 38%, respectively) compared to paroxetine administered alone. The effect of paroxetine on phenobarbital pharmacokinetics was not studied. Because paroxetine exhibits nonlinear pharmacokinetics, the results of this study may not address the case where the 2 drugs are both being chronically dosed [see *Drug Interactions (7.2)*].

Phenytoin: When a single oral 30 mg dose of paroxetine was administered at phenytoin steady state (300 mg once daily for 14 days), paroxetine AUC and $T_{1/2}$ were reduced (by an average of 50% and 35%, respectively) compared to paroxetine administered alone. In a separate study, when a single oral 300 mg dose of phenytoin was administered at paroxetine steady state (30 mg once daily for 14 days), phenytoin AUC was slightly reduced (12% on average) compared to phenytoin administered alone. Because both drugs exhibit nonlinear pharmacokinetics, the above studies may not address the case where the 2 drugs are both being chronically dosed [see *Drug Interactions (7.2)*].

Digoxin: A clinical drug interaction study showed that concurrent use of digoxin did not affect paroxetine exposure.

Diazepam: A clinical drug interaction study showed that concurrent use of diazepam did not affect paroxetine exposure.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Two-year carcinogenicity studies were conducted in rodents given paroxetine in the diet at 1, 5, and 25 mg/kg/day (mice) and 1, 5, and 20 mg/kg/day (rats). The doses used in these carcinogenicity studies were approximately 16 (mouse) and 26 (rat) times the MHRD for VMS. There was a significantly greater number of male rats in the high-dose group with reticulum cell sarcomas (1/100, 0/50, 0/50, and 4/50 for control, low-, middle-, and high-dose groups, respectively) and a significantly increased linear trend across groups for the occurrence of lymphoreticular tumors in male rats. Female rats were not affected. Although there was a dose-related increase in the number of tumors in mice, there was no drug-related increase in the number of mice with tumors. The relevance of these findings to humans is unknown.

Mutagenesis

Paroxetine produced no genotoxic effect in a battery of 5 *in vitro* and 2 *in vivo* assays that included the following: bacterial mutation assay, mouse lymphoma mutation assay, unscheduled DNA synthesis assay, and tests for cytogenetic aberrations *in vivo* in mouse bone marrow and *in vitro* in human lymphocytes and in a dominant lethal test in rats.

Impairment of Fertility

A reduced pregnancy rate was found in reproduction studies in rats at a paroxetine dose of 15 mg/kg/day, which is 19 times the MRHD for VMS on an mg/m² basis. Irreversible lesions occurred in the reproductive tract of male rats after dosing in toxicity studies for 2 to 52 weeks. These lesions consisted of vacuolation of epididymal tubular epithelium at 50 mg/kg/day and atrophic changes in the seminiferous tubules of the testes with arrested spermatogenesis at 25 mg/kg/day (65 times and 32 times the MHRD for VMS on an mg/m² basis, respectively).

14 CLINICAL STUDIES

The efficacy of paroxetine as a treatment for moderate to severe VMS associated with menopause was established in two Phase 3 studies (at a dose of 7.5 mg once daily at bedtime) in 1174 postmenopausal women with a minimum of 7 to 8 moderate to severe vasomotor symptoms per day at baseline (greater than or equal to 50 per week) for 30 days prior to receiving study drug.

Study 1 was a 12-week, randomized, double-blind, placebo-controlled clinical trial with a total of 606 postmenopausal women (average age 55 years, 65% Caucasian and 33% African American, 18% surgically menopausal and 82% naturally menopausal).

Study 2 was a 24-week, randomized, double-blind, placebo-controlled clinical trial with a total of 568 postmenopausal women (average age 54 years, 76% Caucasian and 22% African American, 20% surgically menopausal and 81% naturally menopausal).

The co-primary efficacy endpoints for both studies were the reduction from baseline in VMS frequency and severity at Weeks 4 and 12. Data from Study 1 showed a statistically significant reduction from baseline in the frequency of moderate to severe vasomotor symptoms at Week 4 and Week 12 and a statistically significant reduction in the severity of moderate to severe VMS at Week 4 for paroxetine compared to placebo (Table 4). Data from Study 2 showed a statistically significant reduction from baseline in the frequency and severity of moderate to severe vasomotor symptoms at Week 4 and Week 12 for paroxetine compared to placebo (Table 5).

Table 4 Study 1: Changes in the Daily Frequency and Daily Severity of Moderate to Severe VMS at Weeks 4 and 12 (MITT Population)

	Frequency		Severity	
	Paroxetine Capsules	Placebo	Paroxetine Capsules	Placebo
Baseline				
n	301	305	301	305
Median	10.4	10.4	2.5	2.5
Change from baseline at Week 4				
n	289	293	281	289
Median	-4.3	-3.1	-0.05	0.00
Treatment Difference*	-1.2		-0.05	
P-value [#]	<0.01		<0.01	
Change from baseline at Week 12				
n	264	274	236	253
Median	-5.9	-5.0	-0.06	-0.02
Treatment Difference*	-0.9		-0.04	
P-value [#]	<0.01		0.17	

MITT population: all consented and randomized subjects with valid baseline daily hot flash diary data who had taken at least 1 dose of study medication and had at least 1 day of on-treatment daily hot flash diary data.

* Treatment Difference: the difference between the median changes from baseline.

P-value is obtained from rank-ANCOVA model.

Table 5 Study 2: Changes in the Daily Frequency and Daily Severity of Moderate to Severe VMS at Weeks 4 and 12 (MITT Population)

	Frequency		Severity	
	Paroxetine Capsules	Placebo	Paroxetine Capsules	Placebo
Baseline				
n	284	284	284	284
Median	9.9	9.6	2.5	2.5
Change from baseline at Week 4				
n	276	274	268	271
Median	-3.8	-2.5	-0.04	-0.01
Treatment Difference*	-1.3		-0.03	
P-value [#]	<0.01		0.04	
Change from baseline at Week 12				
n	257	244	245	236
Median	-5.6	-3.9	-0.05	0.00
Treatment Difference*	-1.7		-0.05	
P-value [#]	<0.01		<0.01	

MITT population: all consented and randomized subjects with valid baseline daily hot flash diary data who had taken at least 1 dose of study medication and had at least 1 day of on-treatment daily hot flash diary data.

* Treatment Difference: the difference between the median changes from baseline.

P-value is obtained from rank-ANCOVA model.

Persistence of benefit at 24 weeks in Study 2 was evaluated with a responder analysis where responders were defined as those patients who achieved greater than or equal to 50% reduction from baseline in the frequency of moderate to severe VMS at Week 24. The proportion of patients achieving a greater than or equal to 50% reduction in the frequency of moderate to severe VMS from baseline to Week 24 was 48% in the paroxetine group and 36% in the placebo group at Week 24.

16 HOW SUPPLIED/STORAGE AND HANDLING

Paroxetine capsules are available as 7.5 mg, size “2”, hard gelatin capsule with an orange opaque cap printed with “A401” and an orange opaque body printed with “7.5 mg” in black ink, containing a white to off white blend.

NDC 0591-4100-30, blister packs of 30

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature]. Protect from light and humidity.

17 PATIENT COUNSELING INFORMATION

See FDA-approved patient labeling (*Medication Guide*).

Instruct patients to read the Medication Guide before starting therapy with paroxetine capsules and to reread it each time the prescription is renewed.

- Advise patients, their families, and their caregivers to look for the emergence of suicidality, especially early during treatment [see *Boxed Warning and Warnings and Precautions (5.1)*].
- Instruct patients not to take paroxetine capsules with an MAOI or within 14 days of stopping an MAOI and

allow 14 days after stopping paroxetine capsules before starting an MAOI [*see Dosage and Administration (2.2) and Contraindications (4.1)*].

- Advise patients not to take paroxetine capsules with thioridazine or pimozide [*see Contraindications (4.2 and 4.3)*].
- Caution patients about the risk of serotonin syndrome, particularly with the concomitant use of paroxetine capsules with triptans, tricyclic antidepressants, linezolid, tramadol, amphetamines, St. John's Wort, lithium, tryptophan supplements, other serotonergic agents, or antipsychotic drugs [*see Warnings and Precautions (5.2) and Drug Interactions (7.3)*].
- Caution patients that efficacy of tamoxifen may be reduced when administered concomitantly and counsel them about the likely benefit of paroxetine for treating VMS vs. the risk of possible decreased tamoxifen effectiveness [*see Warnings and Precautions (5.3)*].
- Caution patients about the concomitant use of paroxetine capsules and NSAIDs, aspirin, warfarin, and other anticoagulants because combined use of drugs that interfere with serotonin reuptake has been associated with an increased risk of bleeding [*see Warnings and Precautions (5.4)*].
- Advise patients that taking paroxetine can cause mild pupillary dilation, which in susceptible individuals, can lead to an episode of angle-closure glaucoma. Pre-existing glaucoma is almost always open-angle glaucoma because angle-closure glaucoma, when diagnosed, can be treated definitively with iridectomy. Open-angle glaucoma is not a risk factor for angle-closure glaucoma. Patients may wish to be examined to determine whether they are susceptible to angle-closure, and have a prophylactic procedure (e.g., iridectomy), if they are susceptible [*see Warnings and Precautions (5.5)*].
- Caution patients about the risk of hyponatremia, particularly elderly patients and those who are taking diuretics or are volume-depleted [*see Warnings and Precautions (5.6)*].
- Inform patients that there is the possibility for an increased risk of fracture [*see Warnings and Precautions (5.7)*].
- Advise patients, their families, and their caregivers to observe for signs of activation of mania/hypomania [*see Warnings and Precautions (5.8)*].
- Advise patients to notify their physician if they become pregnant during therapy [*see Contraindications (4.5) and Use in Specific Populations (8.1)*].
- Caution patients about operating hazardous machinery, including motor vehicles, until they are reasonably certain that paroxetine therapy does not affect their ability to engage in such activities [*see Warnings and Precautions (5.11)*].
- Advise patients to inform their healthcare provider if they are taking, or plan to take, any prescription or over-the-counter drugs, including herbal supplements, because there is a potential for interaction with paroxetine [*see Drug Interactions (7.3)*].
- Advise patients that paroxetine, the active ingredient in paroxetine capsules, is also the active ingredient in certain other drugs and these medications should not be taken concomitantly [*see Indications and Usage (1) and Drug Interactions (7.3)*].

Manufactured by:
Actavis Laboratories FL, Inc.
Fort Lauderdale, FL 33314 USA

Distributed by:
Actavis Pharma, Inc.
Parsippany, NJ 07054 USA

Revised: May 2017



MEDICATION GUIDE

Paroxetine (pa-ROX-e-teen) Capsules

Read the Medication Guide that comes with paroxetine capsules before you start taking it and each time you get a refill. There may be new information. This Medication Guide does not take the place of talking to your healthcare provider about your medical condition or treatment. Talk with your healthcare provider if there is something you do not understand or want to learn more about. Paroxetine capsules contain a lower dose of paroxetine, a medicine also used to treat a number of psychiatric disorders. The lower dose of paroxetine in paroxetine capsules has not been studied in any psychiatric conditions and paroxetine capsules are not approved for any psychiatric uses.

What is the most important information I should know about paroxetine capsules?

Paroxetine capsules may cause serious side effects.

Call your healthcare provider right away if you have any of the following symptoms, or go to the nearest emergency room:

1. Suicidal thoughts or actions:

- **Paroxetine capsules, and related antidepressant medicines, may increase suicidal thoughts or actions within the first few months of treatment.**
- Depression or other serious mental illnesses are the most important causes of suicidal thoughts or actions.
- Watch for these changes and call your healthcare provider right away if you notice:
 - New or sudden changes in mood, behavior, actions, thoughts, or feelings, especially if severe.
 - Pay particular attention to such changes when paroxetine capsules are started.

Keep all follow-up visits with your healthcare provider and call between visits if you are worried about symptoms.

Call your healthcare provider right away or go to the nearest emergency room if you have any of the following symptoms, especially if they are new, worse, or worry you:

- attempts to commit suicide
- acting on dangerous impulses
- acting aggressive or violent
- thoughts about suicide or dying
- new or worse depression
- new or worse anxiety or panic attacks
- feeling agitated, restless, angry or irritable
- trouble sleeping
- an increase in activity or talking more than what is normal for you
- other unusual changes in behavior or mood

2. Serotonin Syndrome. This condition can be life-threatening and may include:

- agitation (nervousness), hallucinations, coma or other changes in mental status
- coordination problems or muscle twitching (small movements of the muscles that you cannot control)
- racing heartbeat, high or low blood pressure
- sweating or fever
- nausea, vomiting, or diarrhea

- muscle rigidity
- dizziness
- flushing
- tremors
- seizures

3. Reduced effectiveness of tamoxifen. Tamoxifen (a medicine used to treat breast cancer) may not work as well if it is taken while you take paroxetine capsules. If you are taking tamoxifen, tell your healthcare provider before starting paroxetine capsules.

4. Abnormal bleeding. Paroxetine capsules may increase your risk of bleeding or bruising, especially if you take the blood thinner warfarin or non-steroidal anti-inflammatory drugs (NSAIDs) like ibuprofen, naproxen, or aspirin.

5. Visual problems.

- Eye pain
- Changes in vision
- Swelling or redness in or around the eye

Only some people are at risk for these problems. You may want to undergo an eye examination to see if you are at risk and receive preventative treatment if you are.

6. Low salt (sodium) levels in the blood. Elderly people may be at greater risk for this condition. Symptoms may include:

- headache
- weakness or feeling unsteady
- confusion, problems concentrating or thinking or memory problems

7. Bone Fractures. Women who take paroxetine capsules may have a higher risk of bone fractures. Contact your healthcare provider if you have pain in a bone.

8. Manic episodes:

- greatly increased energy
- severe trouble sleeping
- racing thoughts
- reckless behavior
- unusually grand ideas
- excessive happiness or irritability
- talking more or faster than usual

9. Seizures or convulsions.

10. Restlessness. Women who take paroxetine capsules may feel an inner restlessness, agitation (nervousness), or be unable to sit still or stand still especially when they start taking paroxetine capsules. Call your healthcare provider if this happens to you.

11. Driving. Paroxetine capsules may affect your ability to make decisions, think clearly, or react quickly. Do not drive, operate heavy machinery, or do other potentially dangerous activities until you know how paroxetine capsules affect you.

What are paroxetine capsules?

Paroxetine capsules are a prescription medicine used to reduce moderate to severe hot flashes associated with menopause. Paroxetine capsules are a selective serotonin reuptake inhibitor (SSRI). It is not a hormone. The way paroxetine treats hot flashes associated with menopause is not known. Paroxetine capsules do not prevent or treat osteoporosis or dryness, itching or burning in and around the vagina.

Paroxetine capsules are not for psychiatric problems such as depression, obsessive compulsive disorder, panic disorder, generalized anxiety disorder, social anxiety disorder, and post-traumatic stress disorder.

Paroxetine capsules are not for use in children.

Talk to your healthcare provider if you do not think that your hot flashes are getting better while taking paroxetine capsules.

Who should not take paroxetine capsules?

Do not take paroxetine capsules if you:

- **take a Monoamine Oxidase Inhibitor (MAOI).** Ask your healthcare provider or pharmacist if you are not sure if you take an MAOI, including the antibiotic linezolid.
 - Do not take an MAOI within 14 days of stopping paroxetine capsules unless directed to do so by your healthcare provider.
 - Do not start paroxetine capsules if you stopped taking an MAOI in the last 14 days unless directed to do so by your healthcare provider.
 - **People who take paroxetine capsules close in time to an MAOI may have serious or life-threatening side effects. Get medical help right away if you have any of these symptoms:**
 - high fever
 - uncontrolled muscle spasms
 - stiff muscles
 - rapid changes in heart rate or blood pressure
 - confusion
 - loss of consciousness (pass out)
- **take thioridazine.** Do not take thioridazine together with paroxetine capsules because this can cause serious heart rhythm problems or sudden death.
- **take the antipsychotic medicine pimozide.** Do not take pimozide together with paroxetine capsules because this can cause serious heart problems.
- **are allergic to paroxetine or any of the ingredients in paroxetine capsules. See the end of this Medication Guide for a complete list of ingredients in paroxetine capsules.**
- **are pregnant.** Paroxetine capsules are not for pregnant women. Paroxetine, the active ingredient in paroxetine capsules, can harm your unborn baby. Risks to your unborn baby include an increased risk of birth defects, particularly heart defects. Your baby may also have certain other serious symptoms shortly after birth.

What should I tell my healthcare provider before taking paroxetine capsules?

Before starting paroxetine capsules, tell your healthcare provider if you:

- have liver problems
- have kidney problems
- have or had seizures or convulsions
- have bipolar disorder or mania
- have low sodium levels in your blood

- have or had bleeding problems
- have glaucoma (high pressure in the eye)
- have any other medical conditions
- **are breastfeeding or plan to breastfeed.** Paroxetine passes into breast milk. Talk to your healthcare provider before taking paroxetine capsules if you are breastfeeding.

Tell your healthcare provider about all the medicines that you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Paroxetine capsules and some medicines may interact with each other, may not work as well, or may cause serious side effects when taken together.

If you take paroxetine capsules, you should not take any other medicines that contain paroxetine, including Paxil, Paxil CR and Pexeva.

Especially tell your healthcare provider if you take:

- triptans used to treat migraine headache
- medicines used to treat mood, anxiety, psychotic or thought disorders, including MAOIs, SSRIs, tricyclics, lithium, buspirone, or antipsychotics
- tramadol, fentanyl or over-the-counter supplements such as tryptophan or St. John’s Wort
- amphetamines
- thioridazine
- pimozide
- tamoxifen
- atomoxetine
- cimetidine
- digoxin
- theophylline
- medicines to treat irregular heart rate (like propafenone, flecainide, and encainide)
- medicines used to treat schizophrenia
- certain medicines used to treat HIV infection
- the blood thinner warfarin
- nonsteroidal anti-inflammatory drugs (NSAIDs) (like ibuprofen, naproxen, or aspirin)
- certain medicines used to treat seizures (like phenobarbital and phenytoin)
- other drugs containing paroxetine, the medicine in paroxetine capsules.

Ask your healthcare provider if you are not sure if you are taking any of these medications.

Your healthcare provider or pharmacist can tell you if it is safe to take paroxetine capsules with your other medicines. Do not start or stop any medicine while taking paroxetine capsules without talking to your healthcare provider first.

How should I take paroxetine capsules?

- Take paroxetine capsules exactly as your healthcare provider tells you to take it.
- Take paroxetine capsules 1 time each day at bedtime.
- Paroxetine capsules may be taken with or without food.
- If you miss a dose of paroxetine capsules, take the missed dose as soon as you remember. If it is almost time for the next dose, skip the missed dose and take your next dose at the regular time. Do not take two doses of paroxetine capsules at the same time.
- If you take too many paroxetine capsules, call your healthcare provider or poison control center right away, or go to the nearest emergency room right away.

What should I avoid while taking paroxetine capsules?

- Paroxetine capsules can cause sleepiness or may affect your ability to make decisions, think clearly, or react quickly. You should not drive, operate heavy machinery, or do other dangerous activities until you know how paroxetine capsules affect you.

What are the possible side effects of paroxetine capsules?

Paroxetine capsules may cause serious side effects, including:

- See “What is the most important information I should know about paroxetine capsules?”

The most common side effects of paroxetine capsules include:

- headache
- tiredness
- nausea and vomiting

Tell your healthcare provider if you have any side effect that bothers you or does not go away. These are not all the possible side effects of paroxetine capsules. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store paroxetine capsules?

- Store paroxetine capsules at room temperature between 68° to 77°F (20° to 25°C).
- Keep paroxetine capsules out of the light.
- Keep paroxetine capsules dry.
- **Keep paroxetine capsules and all medicines out of the reach of children.**

General information about the safe and effective use of paroxetine capsules.

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use paroxetine capsules for a condition for which they were not prescribed. Do not give paroxetine capsules to other people, even if they have the same condition. It may harm them.

This Medication Guide summarizes the most important information about paroxetine capsules. If you would like more information, talk with your healthcare provider. You may ask your healthcare provider or pharmacist for information about paroxetine capsules that is written for healthcare professionals.

For more information about paroxetine capsules call 1-800-272-5525 or go to www.actavis.com.

What are the ingredients in paroxetine capsules?

Active ingredient: paroxetine

Inactive ingredients: anhydrous citric acid, colloidal silicon dioxide, dibasic calcium phosphate anhydrous, FD&C Yellow #6, gelatin, iron oxide black, magnesium stearate, microcrystalline cellulose, potassium hydroxide, propylene glycol, shellac, sodium lauryl sulfate, sodium starch glycolate and titanium dioxide.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Brands listed are the trademarks of their respective owners.

Manufactured by:

Actavis Laboratories FL, Inc.
Fort Lauderdale, FL 33314 USA

Distributed by:
Actavis Pharma, Inc.
Parsippany, NJ 07054 USA

Revised: May 2017

MEDICATION GUIDE

Paroxetine (pa-ROX-e-teen) Capsules

Read the Medication Guide that comes with paroxetine capsules before you start taking it and each time you get a refill. There may be new information. This Medication Guide does not take the place of talking to your healthcare provider about your medical condition or treatment. Talk with your healthcare provider if there is something you do not understand or want to learn more about. Paroxetine capsules contain a lower dose of paroxetine, a medicine also used to treat a number of psychiatric disorders. The lower dose of paroxetine in paroxetine capsules has not been studied in any psychiatric conditions and paroxetine capsules are not approved for any psychiatric uses.

What is the most important information I should know about paroxetine capsules?

Paroxetine capsules may cause serious side effects.

Call your healthcare provider right away if you have any of the following symptoms, or go to the nearest emergency room:

1. Suicidal thoughts or actions:

- **Paroxetine capsules, and related antidepressant medicines, may increase suicidal thoughts or actions within the first few months of treatment.**
- Depression or other serious mental illnesses are the most important causes of suicidal thoughts or actions.
- Watch for these changes and call your healthcare provider right away if you notice:
 - New or sudden changes in mood, behavior, actions, thoughts, or feelings, especially if severe.
 - Pay particular attention to such changes when paroxetine capsules are started.

Keep all follow-up visits with your healthcare provider and call between visits if you are worried about symptoms.

Call your healthcare provider right away or go to the nearest emergency room if you have any of the following symptoms, especially if they are new, worse, or worry you:

- attempts to commit suicide
- acting on dangerous impulses
- acting aggressive or violent
- thoughts about suicide or dying
- new or worse depression
- new or worse anxiety or panic attacks
- feeling agitated, restless, angry or irritable
- trouble sleeping
- an increase in activity or talking more than what is normal for you
- other unusual changes in behavior or mood

2. Serotonin Syndrome. This condition can be life-threatening and may include:

- agitation (nervousness), hallucinations, coma or other changes in mental status
- coordination problems or muscle twitching (small movements of the muscles that you cannot control)
- racing heartbeat, high or low blood pressure
- sweating or fever
- nausea, vomiting, or diarrhea
- muscle rigidity

- dizziness
- flushing
- tremors
- seizures

3. Reduced effectiveness of tamoxifen. Tamoxifen (a medicine used to treat breast cancer) may not work as well if it is taken while you take paroxetine capsules. If you are taking tamoxifen, tell your healthcare provider before starting paroxetine capsules.

4. Abnormal bleeding. Paroxetine capsules may increase your risk of bleeding or bruising, especially if you take the blood thinner warfarin or non-steroidal anti-inflammatory drugs (NSAIDs) like ibuprofen, naproxen, or aspirin.

5. Visual problems.

- Eye pain
- Changes in vision
- Swelling or redness in or around the eye

Only some people are at risk for these problems. You may want to undergo an eye examination to see if you are at risk and receive preventative treatment if you are.

6. Low salt (sodium) levels in the blood. Elderly people may be at greater risk for this condition. Symptoms may include:

- headache
- weakness or feeling unsteady
- confusion, problems concentrating or thinking or memory problems

7. Bone Fractures. Women who take paroxetine capsules may have a higher risk of bone fractures. Contact your healthcare provider if you have pain in a bone.

8. Manic episodes:

- greatly increased energy
- severe trouble sleeping
- racing thoughts
- reckless behavior
- unusually grand ideas
- excessive happiness or irritability
- talking more or faster than usual

9. Seizures or convulsions.

10. Restlessness. Women who take paroxetine capsules may feel an inner restlessness, agitation (nervousness), or be unable to sit still or stand still especially when they start taking paroxetine capsules. Call your healthcare provider if this happens to you.

11. Driving. Paroxetine capsules may affect your ability to make decisions, think clearly, or react quickly. Do not drive, operate heavy machinery, or do other potentially dangerous activities until you know how paroxetine capsules affect you.

What are paroxetine capsules?

Paroxetine capsules are a prescription medicine used to reduce moderate to severe hot flashes associated with menopause. Paroxetine capsules are a selective serotonin reuptake inhibitor (SSRI). It is not a hormone. The way paroxetine treats hot flashes associated with menopause is not known. Paroxetine capsules do not prevent or treat osteoporosis or dryness, itching or burning in and around the vagina.

Paroxetine capsules are not for psychiatric problems such as depression, obsessive compulsive disorder, panic disorder, generalized anxiety disorder, social anxiety disorder, and post-traumatic stress disorder.

Paroxetine capsules are not for use in children.

Talk to your healthcare provider if you do not think that your hot flashes are getting better while taking paroxetine capsules.

Who should not take paroxetine capsules?

Do not take paroxetine capsules if you:

- **take a Monoamine Oxidase Inhibitor (MAOI).** Ask your healthcare provider or pharmacist if you are not sure if you take an MAOI, including the antibiotic linezolid.
 - Do not take an MAOI within 14 days of stopping paroxetine capsules unless directed to do so by your healthcare provider.
 - Do not start paroxetine capsules if you stopped taking an MAOI in the last 14 days unless directed to do so by your healthcare provider.
 - **People who take paroxetine capsules close in time to an MAOI may have serious or life-threatening side effects. Get medical help right away if you have any of these symptoms:**
 - high fever
 - uncontrolled muscle spasms
 - stiff muscles
 - rapid changes in heart rate or blood pressure
 - confusion
 - loss of consciousness (pass out)
- **take thioridazine.** Do not take thioridazine together with paroxetine capsules because this can cause serious heart rhythm problems or sudden death.
- **take the antipsychotic medicine pimozide.** Do not take pimozide together with paroxetine capsules because this can cause serious heart problems.
- **are allergic to paroxetine or any of the ingredients in paroxetine capsules. See the end of this Medication Guide for a complete list of ingredients in paroxetine capsules.**
- **are pregnant.** Paroxetine capsules are not for pregnant women. Paroxetine, the active ingredient in paroxetine capsules, can harm your unborn baby. Risks to your unborn baby include an increased risk of birth defects, particularly heart defects. Your baby may also have certain other serious symptoms shortly after birth.

What should I tell my healthcare provider before taking paroxetine capsules?

Before starting paroxetine capsules, tell your healthcare provider if you:

- have liver problems
- have kidney problems
- have or had seizures or convulsions
- have bipolar disorder or mania
- have low sodium levels in your blood
- have or had bleeding problems

- have glaucoma (high pressure in the eye)
- have any other medical conditions
- **are breastfeeding or plan to breastfeed.** Paroxetine passes into breast milk. Talk to your healthcare provider before taking paroxetine capsules if you are breastfeeding.

Tell your healthcare provider about all the medicines that you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Paroxetine capsules and some medicines may interact with each other, may not work as well, or may cause serious side effects when taken together.

If you take paroxetine capsules, you should not take any other medicines that contain paroxetine, including Paxil, Paxil CR and Pexeva.

Especially tell your healthcare provider if you take:

- triptans used to treat migraine headache
- medicines used to treat mood, anxiety, psychotic or thought disorders, including MAOIs, SSRIs, tricyclics, lithium, buspirone, or antipsychotics
- tramadol, fentanyl or over-the-counter supplements such as tryptophan or St. John’s Wort
- amphetamines
- thioridazine
- pimozide
- tamoxifen
- atomoxetine
- cimetidine
- digoxin
- theophylline
- medicines to treat irregular heart rate (like propafenone, flecainide, and encainide)
- medicines used to treat schizophrenia
- certain medicines used to treat HIV infection
- the blood thinner warfarin
- nonsteroidal anti-inflammatory drugs (NSAIDs) (like ibuprofen, naproxen, or aspirin)
- certain medicines used to treat seizures (like phenobarbital and phenytoin)
- other drugs containing paroxetine, the medicine in paroxetine capsules.

Ask your healthcare provider if you are not sure if you are taking any of these medications.

Your healthcare provider or pharmacist can tell you if it is safe to take paroxetine capsules with your other medicines. Do not start or stop any medicine while taking paroxetine capsules without talking to your healthcare provider first.

How should I take paroxetine capsules?

- Take paroxetine capsules exactly as your healthcare provider tells you to take it.
- Take paroxetine capsules 1 time each day at bedtime.
- Paroxetine capsules may be taken with or without food.
- If you miss a dose of paroxetine capsules, take the missed dose as soon as you remember. If it is almost time for the next dose, skip the missed dose and take your next dose at the regular time. Do not take two doses of paroxetine capsules at the same time.
- If you take too many paroxetine capsules, call your healthcare provider or poison control center right away, or go to the nearest emergency room right away.

What should I avoid while taking paroxetine capsules?

- Paroxetine capsules can cause sleepiness or may affect your ability to make decisions, think clearly, or react quickly. You should not drive, operate heavy machinery, or do other dangerous activities until you know how paroxetine capsules affect you.

What are the possible side effects of paroxetine capsules?

Paroxetine capsules may cause serious side effects, including:

- See “What is the most important information I should know about paroxetine capsules?”

The most common side effects of paroxetine capsules include:

- headache
- tiredness
- nausea and vomiting

Tell your healthcare provider if you have any side effect that bothers you or does not go away. These are not all the possible side effects of paroxetine capsules. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store paroxetine capsules?

- Store paroxetine capsules at room temperature between 68° to 77°F (20° to 25°C).
- Keep paroxetine capsules out of the light.
- Keep paroxetine capsules dry.
- **Keep paroxetine capsules and all medicines out of the reach of children.**

General information about the safe and effective use of paroxetine capsules.

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use paroxetine capsules for a condition for which they were not prescribed. Do not give paroxetine capsules to other people, even if they have the same condition. It may harm them.

This Medication Guide summarizes the most important information about paroxetine capsules. If you would like more information, talk with your healthcare provider. You may ask your healthcare provider or pharmacist for information about paroxetine capsules that is written for healthcare professionals.

For more information about paroxetine capsules call 1-800-272-5525 or go to www.actavis.com.

What are the ingredients in paroxetine capsules?

Active ingredient: paroxetine

Inactive ingredients: anhydrous citric acid, colloidal silicon dioxide, dibasic calcium phosphate anhydrous, FD&C Yellow #6, gelatin, iron oxide black, magnesium stearate, microcrystalline cellulose, potassium hydroxide, propylene glycol, shellac, sodium lauryl sulfate, sodium starch glycolate and titanium dioxide.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Brands listed are the trademarks of their respective owners.

Manufactured by:
Actavis Laboratories FL, Inc.

Fort Lauderdale, FL 33314 USA

Distributed by:
Actavis Pharma, Inc.
Parsippany, NJ 07054 USA

Revised: May 2017