




DRUG FACTS TEXT DEFINED	TYPE SIZE
DRUG FACTS TITLE	9 pt
DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
SPACE BEFORE BULLET	2 ems
BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces













o p.
6 pt
6.5 pt
<39
5 pt
2 ems
1.5 pt, .5 pt
2 spaces



PRODUCTION (NOT COLOR ACCURATE) (*)



DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	5.5 pt
• SUBHEADINGS/BODY TEXT	5.5 pt
• LEADING	5.5 pt
• BULLETS	4.5 pt
SPACE BEFORE BULLET	2 ems
• WARNING BOX LINE, HAIRLINES	.5 pt



PRODUCTION


(NOT COLOR ACCURATE)


DRUG FACTS TEXT DEFINED	TYPE SIZE
DRUG FACTS TEXT DEFINED	TTPE SIZE
DRUG FACTS TITLE	9.0 pt
• DRUG FACTS CONTINUED	8.0 pt
• HEADINGS	8.0 pt
• SUBHEADINGS/BODY TEXT	6.0 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5.0 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

(b) (4 Full size file



Cut
Crease
1/8 1/8 perf
1/2 x 3/16 Cut Land






PRODUCTION (NO


(NOT COLOR ACCURATE)


(b) (4)



DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	5.5 pt
SUBHEADINGS/BODY TEXT	5.5 pt
• LEADING	5.5 pt
• BULLETS	4.5 pt
• HAIRLINES	.5 pt







(b) (4 Full size file

(NOT COLOR ACCURATE)

128 CODE = 090870




DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9.0 pt
• DRUG FACTS CONTINUED	8.0 pt
• HEADINGS	8.0 pt
• SUBHEADINGS/BODY TEXT	6.0 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5.0 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces





PRODUCTION	(NOT COLOR ACCURATE)		
		(b) (4)	
			• DRUG
			• DRUG
			HEADIN
			• SUBHE
			LEADIN
			• # OF CI
			• BULLE
			SPACE
			BARLIN
			SPACE

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	10 pt
• DRUG FACTS CONTINUED	9 pt
• HEADINGS	9 pt
• SUBHEADINGS/BODY TEXT	7 pt
• LEADING	7.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	6 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



Full size file

(b) (4)

(b) (4

	Directions a duties 18 years of age and older this product is to be used once a day (every 24 hours), every day for 14 days may take 1 to 4 days for full effect may take 1 to 4 days for full effect	Uses I teats frequent heartburn (occurs <u>2 or more</u> days a week) I to 4 days for full effect 1 to 4 days for full effect
	If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children . In case of overdose, get medical help or contact a Poison Control Center right away.	PrincipalPittingEsomegrazole 20 mg
	Drug Facts (continued)	Drug Facts
эш гри о	 Tips for Managing Heartburk Arold foods or drinks that are more likely for visual such as rich, spicy, fatty a case heartburk, such as rich, spicy, spi	esomeprazole magnesium delayed-release tablets 20 mg/acid reducer

prescription antiretrovirals (medicines for HIV infection)	Questions or comments? call toll-free weekdays 9 AM to
Ask a doctor or pharmacist before use if you are taking wardarin, clopidogrel or cilostazol (blood-thinning medicines) digoxin (heart medicine) diazepam (anxiety medicine) tacrolimus or mycophenolate modetil (immune system medicines) tacrolimus or mycophenolate modetil (immune system medicines)	Other Information ead the directions and varnings before use ead the acetions and varnings before use store at 20-25°C (68-77°F) active ingredients con starch, crospovidone, D&C red no. 27 aluminum lake, FD&C blue monostearate, hydroxypropyl cellulose, hypromellose, magnesum monostearate, hydroxypropyl cellulose, hypromellose, magnesum paraffin, polyethylene glycol, polysorbate 80, sodium stearyl tumarate, sucrose, talc, fitanium dixide, friethyl citrate sucrose, talc, fitanium dixide, friethyl citrate
■ nausea or vomiting ■ stomach pain	
 had heartburn over 3 months. This may be a sign of a more serious contition. frequent wheezing, particularly with heartburn Inexplained weight loss 	
Ask a doctor before use if you have	children may sometimes be caused by a serious condition.
Warnings Allergy alert: Do not use if you are allergic to esomeprazole Do not use if you have: I trouble or pain swallowing food, vomiting with blood, or bloody or black stools I chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness frequent chest pain I these may be signs of a serious condition. See your doctor.	 14-Day Course of Treatment 14-Day Course of Treatment availow 1 fablet with a glass of water before eating in the morning atke every day for 14 days do not take more than 1 tablet a day do not take more than 1 tablet a day you may repeat a 14-Day Course (it needed) you may repeat a 14-day source (it needed) you may repeat a 14-day source often than every may not take more than 14 days unless directed by your doctor do not use for more than 14 days or more often than every may not use for more than 14 days or more often than every
Uses ■ treats frequent heartburn (occurs <u>2 or more</u> days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect	Directions adults 18 years of age and older it its product is to be used once a day (every 24 hours), every day for 14 days may take 1 to 4 days for full effect
Purpose ingredient (in each tablet) Purpose Esomeprazole 20 mg. Acid reducer Each delayed-release tablet corresponds to 22.3 mg (Each delayed-release tablet corresponds to 22.3 mg esometric)	If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

VALERIE S PRATT 12/18/2017