

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use KINEVAC safely and effectively. See full prescribing information for KINEVAC

KINEVAC (sincalide for injection), for intravenous use

Initial U.S. Approval: 1976

RECENT MAJOR CHANGES

Contraindications (4)	03/2018
Warnings and Precautions (5.1)	03/2018

INDICATIONS AND USAGE

Kinevac is a cholecystokin (CCK) analog indicated in adults to:

- stimulate gallbladder contraction, as may be assessed by various methods of diagnostic imaging, or to obtain by duodenal aspiration a sample of concentrated bile for analysis of cholesterol, bile salts, phospholipids, and crystals. (1)
- stimulate pancreatic secretion in combination with secretin prior to obtaining a duodenal aspirate for analysis of enzyme activity, composition, and cytology. (1)
- accelerate the transit of a barium meal through the small bowel, thereby decreasing the time and extent of radiation associated with fluoroscopy and x-ray examination of the intestinal tract. (1)

DOSAGE AND ADMINISTRATION

Recommended Adult Dosage and Administration by Indication:

To Stimulate Contraction of the Gallbladder

- 0.02 mcg/kg as a single dose over 30 to 60 seconds via intravenous injection. If satisfactory contraction does not occur in 15 minutes, administer a dose of 0.04 mcg/kg over 30 to 60 seconds. (2.1)
- Alternatively consider an intravenous infusion to reduce gastrointestinal adverse reactions: 0.12 mcg/kg diluted in 100 mL of 0.9% Sodium Chloride Injection USP and infused over 50 minutes at a rate of 2 mL per minute. (2.1, 2.2, 5.3)

To Stimulate Pancreatic Secretion in Combination with Secretin

- 30 minutes after initiation of secretin for injection, administer 0.02 mcg/kg diluted in 30 mL of 0.9% Sodium Chloride Injection USP and infused over 30 minutes at a rate of 1 mL per minute. (2.1, 2.2)

To Accelerate Transit of a Barium Meal Through the Small Intestine

- After the barium meal is beyond the proximal jejunum, administer 0.04 mcg/kg over 30 to 60 seconds via intravenous injection. (2.1)
- If satisfactory transit of the barium meal has not occurred in 30 minutes, administer a second dose of 0.04 mcg/kg over 30 to 60 seconds. (2.1)
- Alternatively consider an intravenous infusion to reduce gastrointestinal adverse reactions: 0.12 mcg/kg diluted in 100 mL 0.9% Sodium Chloride Injection USP and infused over 30 minutes. (2.1, 2.2, 5.3)

DOSAGE FORMS AND STRENGTHS

For injection: 5 mcg of sincalide as a lyophilized powder in a single-dose vial for reconstitution (3)

CONTRAINDICATIONS

- History of hypersensitivity to sincalide. (4, 5.1)
- Intestinal obstruction. (4)

WARNINGS AND PRECAUTIONS

- Anaphylaxis, Anaphylactic Shock and Other Hypersensitivity Reactions:** May occur during or soon after administration. If symptoms occur, discontinue the drug. (4, 5.1)
- Evacuation of Gallstones:** Stimulation of gallbladder contraction in patients with small gallbladder stones could lead to the evacuation of the stones from the gallbladder, resulting in their lodging in the cystic duct or in the common bile duct. (5.2)
- Gastrointestinal Adverse Reactions with Intravenous Injection:** Administration as an intravenous injection may cause transient nausea, vomiting, abdominal pain or cramping, dizziness or flushing. To reduce the risk of adverse reactions when used to simulate contraction of the gallbladder or accelerate transit of a barium meal through the small intestine, administer as an intravenous infusion over 50 or 30 minutes, respectively. (2.1, 5.3)
- Preterm Labor or Spontaneous Abortion:** Advise pregnant women of the potential risk for preterm labor and spontaneous abortion. (5.4, 8.1)

ADVERSE REACTIONS

Most common adverse reactions ($\geq 20\%$) are: abdominal discomfort or pain, and nausea. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Bracco Diagnostics Inc. at 1-800-257-5181 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

Drugs that Affect Gallbladder Motility or Contractile Response: May interfere with response to sincalide. Consider discontinuing these drugs prior to administration of Kinevac, when used to stimulate contraction of the gallbladder. (7.1)

See 17 for PATIENT COUNSELING INFORMATION

Revised: 12/2018

FULL PRESCRIBING INFORMATION:CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- Recommended Dosage and Administration Instructions
- Preparation Instructions

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- Anaphylaxis, Anaphylactic Shock and Other Hypersensitivity Reactions
- Evacuation of Gallstones
- Gastrointestinal Adverse Reactions with Intravenous Injection
- Preterm Labor or Spontaneous Abortion

6 ADVERSE REACTIONS

7 DRUG INTERACTIONS

- Drugs that Affect Gallbladder Motility or Contractile Response

8 USE IN SPECIFIC POPULATIONS

- Pregnancy
- Lactation
- Pediatric Use
- Geriatric Use

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- Mechanism of Action
- Pharmacodynamics

13 NONCLINICAL TOXICOLOGY

- Carcinogenesis, Mutagenesis, Impairment of Fertility

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Kinevac is indicated in adults to:

- to stimulate gallbladder contraction, as may be assessed by various methods of diagnostic imaging, or to obtain by duodenal aspiration a sample of concentrated bile for analysis of cholesterol, bile salts, phospholipids, and crystals;
- to stimulate pancreatic secretion in combination with secretin prior to obtaining a duodenal aspirate for analysis of enzyme activity, composition, and cytology;
- to accelerate the transit of a barium meal through the small bowel, thereby decreasing the time and extent of radiation associated with fluoroscopy and x-ray examination of the intestinal tract.

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosage and Administration Instructions

The recommended dosage and administration of Kinevac by indication is shown in Table 1. For preparation instructions see *Dosage and Administration* (2.2).

Table 1: Recommended Adult Dosage and Administration of Kinevac by Treatment Indication

Indication	Recommended Adult Dosage and Administration of KINEVAC
To stimulate contraction of the gallbladder	<p>Kinevac 0.02 mcg/kg as a single dose over 30 to 60 seconds via intravenous injection. If satisfactory contraction does not occur in 15 minutes, administer a dose of 0.04 mcg/kg over 30 to 60 seconds.</p> <p><u>Alternatively, Consider an Intravenous Infusion to Reduce Gastrointestinal Adverse Reactions [see Warnings and Precautions (5.3)]:</u> 0.12 mcg/kg diluted in 100 mL of 0.9% Sodium Chloride Injection USP and infused over 50 minutes at a rate of 2 mL per minute.</p>
To stimulate pancreatic secretion in combination with secretin for injection	<p>Secretin for Injection: 0.25 units/kg as intravenous infusion over 60 minutes</p> <p>Kinevac: 30 minutes after initiation of secretin infusion, administer Kinevac 0.02 mcg/kg diluted in 30 mL of 0.9% Sodium Chloride Injection USP and infused over 30 minutes at a rate of 1 mL per minute.</p>
To accelerate the transit of a barium meal through the small intestine	<p>After the barium meal is beyond the proximal jejunum, administer Kinevac 0.04 mcg/kg over 30 to 60 seconds via intravenous injection.</p> <p>If satisfactory transit of the barium meal has not occurred in 30 minutes, administer a second dose of 0.04 mcg/kg over 30 to 60 seconds.</p> <p><u>Alternatively, Consider an Intravenous Infusion to Reduce Gastrointestinal Adverse Reactions [see Warnings and Precautions (5.3)]:</u> 0.12 mcg/kg diluted in 100 mL 0.9% Sodium Chloride Injection USP and infused over 30 minutes.</p>

2.2 Preparation Instructions

For Intravenous Injection

- Reconstitute Kinevac aseptically by adding 5 mL of Sterile Water for Injection USP to the vial.
- Inspect the reconstituted solution visually for particulate matter and discoloration after reconstitution and prior to administration.
- Withdraw the prescribed dose of the reconstituted solution from the vial and administer as an intravenous injection over 30 to 60 seconds, as shown in Table 1. Discard the unused portion.
- Store the reconstituted solution at room temperature. *Discard after 8 hours.*
- For single use only; discard unused portion.

For Intravenous Infusion

- Reconstitute Kinevac aseptically by adding 5 mL of Sterile Water for Injection USP to the vial.
- After reconstitution, withdraw the prescribed dose of the solution from the vial. Discard unused portion.
- Dilute the reconstituted solution in 30 mL or 100 mL of 0.9% Sodium Chloride Injection USP, depending on the indication, as described in Table 1.
- Inspect the Kinevac solutions visually for particulate matter and discoloration after reconstitution, dilution and prior to administration.
- Store the diluted solution at room temperature. *Discard after 1 hour.*

3 DOSAGE FORMS AND STRENGTHS

For injection: 5 mcg of sincalide as a lyophilized white powder for reconstitution in a single-dose vial.

4 CONTRAINDICATIONS

KINEVAC is contraindicated in patients with:

- a history of hypersensitivity to sincalide. Serious hypersensitivity reactions have included anaphylaxis and anaphylactic shock [*see Warnings and Precautions (5.1), Adverse Reactions (6)*].
- intestinal obstruction.

5 WARNINGS AND PRECAUTIONS

5.1 Anaphylaxis, Anaphylactic Shock and Other Hypersensitivity Reactions

In postmarketing experience, anaphylaxis, anaphylactic shock and other serious hypersensitivity reactions have been reported during and within one hour following administration of Kinevac [see *Adverse Reactions (6)*].

Due to the potential for anaphylaxis, appropriate medical support should be readily available when Kinevac is administered. If anaphylaxis or other hypersensitivity reactions occur, immediately discontinue the infusion and initiate appropriate medical treatment. Observe patients closely during and after the infusion. Do not reinitiate Kinevac in patients who have experienced symptoms of hypersensitivity [see *Contraindications (4)*].

5.2 Evacuation of Gallstones

Stimulation of gallbladder contraction in patients with small gallbladder stones could lead to the evacuation of the stones from the gallbladder, resulting in their lodging in the cystic duct or in the common bile duct.

5.3 Gastrointestinal Adverse Reactions with Intravenous Injection

Administration of Kinevac as an intravenous injection may cause adverse reactions such as nausea, vomiting, abdominal pain or cramping, dizziness, and flushing [see *Adverse Reactions (6)*]. These reactions are generally transient. To reduce the risk of adverse reactions with intravenous injection when used to simulate contraction of the gallbladder or accelerate transit of a barium meal through the small intestine, administer Kinevac as an intravenous infusion over 50 or 30 minutes, respectively [see *Dosage and Administration (2.1)*].

5.4 Preterm Labor or Spontaneous Abortion

Because of Kinevac's effect on smooth muscle, pregnant patients should be advised that spontaneous abortion or premature induction of labor may occur [see *Use in Specific Populations (8.1)*].

6 ADVERSE REACTIONS

The following adverse reactions associated with the use of Kinevac were identified in clinical trials or postmarketing reports. Because these reactions were reported voluntarily from a population of uncertain size, it is not always possible to estimate their frequency, reliably, or to establish a causal relationship to drug exposure.

The most frequent adverse reactions (20% or greater) are gastrointestinal: abdominal discomfort or pain, and nausea; these may not necessarily indicate an abnormality of the biliary tract unless there is other clinical or radiologic evidence of disease.

Less common adverse reactions include:

Hypersensitivity reactions: anaphylaxis and anaphylactic shock, hypotension, throat tightness, bradycardia, shortness of breath, nausea, abdominal cramping, diaphoresis, hives, rash, itching; and numbness of face, lips and eyes [see *Contraindications (4)*, *Warnings and Precautions (5.1)*].

Neurological reactions: seizures, headache.

Vasovagal reactions: dizziness, loss of consciousness, nausea, diaphoresis, syncope and hypotension (generally self-limiting).

Other: nausea, vomiting, flushing, hypertension, urge to defecate, diarrhea, sneezing.

7 DRUG INTERACTIONS

7.1 Drugs that Affect Gallbladder Motility or Contractile Response

Drugs that may stimulate or inhibit gallbladder motility or contractile response may interfere with the response to sincalide. Consider discontinuing these drugs prior to administration of Kinevac, when used to simulate contraction of the gallbladder.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Based on limited human data and mechanism of action, sincalide may cause preterm labor or spontaneous abortion [see *Warnings and Precautions (5.4)*]. Available data with sincalide for injection are insufficient to establish a drug-associated risk of major birth defects, miscarriage or adverse maternal or fetal outcomes. In animal embryo-fetal development studies in which sincalide was administered to hamsters and rats during the period of organogenesis, no effects were seen at doses comparable to the maximum recommended clinical dose on a mg/kg basis. However, in a prenatal development study in which rats were administered sincalide during organogenesis through parturition, decreased weight gain and developmental delays were observed at a dose 122 times higher than the maximum recommended human dose based on body surface area.

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

Data

Animal Data

There were no effects on embryo-fetal development in hamsters when sincalide was administered subcutaneously at 250 or 750 ng/kg during organogenesis (Gestation Days 7 to 13) at doses up to 0.8 times the maximum recommended dose of 120 ng/kg on a body surface area basis. No effects on embryo-fetal development were observed in Sprague-Dawley rats at subcutaneous doses of 250, 450, or 750 ng/kg from Gestation Days 6 to 16, representing 1.0 time the maximum recommended human dose on a body surface area basis. In a separate study at a higher dose of 90 mcg/kg administered subcutaneously to CFY rats from Gestation Day 10 through parturition (representing 122 times the maximum recommended human dose on a body surface area basis), offspring showed decreased growth, behavioral changes, and developmental delays.

8.2 Lactation

Risk Summary

There are no data regarding the presence of sincalide in human or animal milk, the effects on the breastfed infant, or the effects on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for Kinevac and any potential adverse effect on the breastfed infant from Kinevac or from the underlying condition.

8.4 Pediatric Use

The safety and effectiveness in pediatric patients have not been established.

8.5 Geriatric Use

Clinical studies of Kinevac did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects.

10 OVERDOSAGE

In the event of an overdose, symptoms related to vagal stimulation, such as gastrointestinal symptoms

possible impairment of fertility in males or females.

16 HOW SUPPLIED/STORAGE AND HANDLING

Kinevac (sincalide for injection) is supplied as 5 mcg of sincalide as a lyophilized white powder for reconstitution in a single-dose vial; in packages of 10 vials (NDC 0270-0556-15).

Store at 25° C (77° F); excursions permitted to 15-30° C (59-86° F) [See USP Controlled Room Temperature].

17 PATIENT COUNSELING INFORMATION

Anaphylaxis, Anaphylactic Shock and Other Hypersensitivity Reactions

Inform patients that hypersensitivity reactions, including anaphylaxis and anaphylactic shock have been reported during or following administration of Kinevac. Advise patients to report immediately to a healthcare provider if they experience symptoms of a hypersensitivity reaction [*see Warnings and Precautions (5.1)*].

Gastrointestinal Adverse Reactions

Advise patients that Kinevac may cause transient gastrointestinal symptoms [*see Warnings and Precautions (5.3)*].

Pregnancy

Advise pregnant women of the potential risk for preterm labor and spontaneous abortion [*see Warnings and Precautions (5.4), Use in Specific Populations (8.1)*].

U.S. Patent 6,803,046

Rx only

Manufactured for
Bracco Diagnostics Inc.
Monroe Township, NJ 08831
By Jubilant HollisterStier LLC
Spokane, WA 99207