


TOP (BACK)

Warnings

- **Ask your doctor before use** if you are pregnant, under a doctor's care for a serious condition, age 60 or over, taking any other drug or have stomach problems.
- This product may cause a **severe allergic reaction**, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Do not use** this product if you have ever had an allergic reaction to any pain reliever/fever reducer.
- **Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. [The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed.

COMPOSITE



BASE

■ **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Questions or comments? call toll free 1-800-88-ADVIL

DO NOT USE
unless vial is sealed in
unbroken plastic blister on
cardboard card.

For most recent product information,
visit www.Advii.com
Pfizer, Madison, NJ 07940 USA
© 2018 Pfizer Inc.
PAA099923.FDA01

TOP (FACE)

Advil®

Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)

10 Tablets **Tablets**

READ AND KEEP CARD FOR COMPLETE
WARNINGS AND INFORMATION

Uses

- temporarily relieves minor aches and pains due to: headache, muscular aches, menstrual cramps, the common cold, backache, toothache, minor pain of arthritis.
- temporarily reduces fever

OPEN HERE TO VIEW MORE
PRODUCT INFORMATION

PAA099923.FDA01

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	6 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• WARNING BOX LINE	.5 pt

TOP LAYER (BACK)

Drug Facts (continued)

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

 If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug


When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better

COMPOSITE



TOP LAYER (FACE)

Drug Facts (continued)

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients acetylated monoglycerides, colloidal silicon dioxide, corn starch, croscarmellose sodium, methylparaben, microcrystalline cellulose, pharmaceutical glaze, pharmaceutical ink, povidone, pregelatinized starch, propylparaben, sodium benzoate, sodium lauryl sulfate, stearic acid, sucrose, synthetic iron oxide, titanium dioxide, white wax

Questions or comments? call toll free 1-800-88-ADVIL

For most recent product information, visit www.Advil.com
Pfizer, Madison, NJ 07940 USA
© 2018 Pfizer Inc.

PAA099924.FDA01

81% FPO

3 0573 0151 12 2

DO NOT USE IF SEAL AROUND CAP IS BROKEN OR MISSING. ADVIL SAFETY SEALED.

Advil®
Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)
10 Coated Tablets **Advil Tablets**

Drug Facts

Active ingredient (in each tablet) Purpose
Ibuprofen 200 mg (NSAID)* ... Pain reliever/Fever reducer
*nonsteroidal anti-inflammatory drug

LIFT HERE for More Drug Facts

BASE LAYER

ADVIL® SAFETY SEALED ADVIL® SAFETY SEALED ADVIL® SAFETY SEALED

TEAR HERE TO OPEN

Drug Facts (continued)

- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
 - any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

PAA099924.FDA01

DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.25 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Advil®

Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)

- Headache
- Backache
- Muscle Ache
- Menstrual Pain
- Minor Arthritis Pain



Advil®

Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)



Advil®

Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)

- Headache
- Backache
- Muscle Ache
- Menstrual Pain
- Minor Arthritis Pain



Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)

Advil®

Home • Office • Travel

Contains: 1 Dozen, 10 Count Pocket Packs

Pfizer, Madison, NJ 07940 USA
© 2016 Pfizer Inc.

For most recent product information, visit www.Advil.com PAA070670

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

KAREN M MAHONEY
10/02/2018