


Carrier 1.00" x .90" (.125" corner radius)



Face .56" x .90"

Flat 3.00" x .56"





Actual Size 100%



Actual Size 100%

(b) (4)

CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	-
Drug Facts	N/A	8 pt	-
(continued):	N/A	8 pt	-
Headings:	N/A	8 pt	4.5 pt
Subheadings:	N/A	6 pt	4.5 pt
Text:	N/A	6 pt	4.5 pt
Bullets:	N/A	5 pt (no larger)	4 pt
Leading:	N/A	+0.5 pt	4.75 pt
Characters Per Inch:	N/A	No more than 39	>39



Bayer
Made in

Dist. by:
Bayer HealthCare LLC
Whippany, NJ 07881

Lot
Exp.

Bayer, the Bayer Cross, Aleve and All Day Strong are registered trademarks of Bayer.

Drug Facts
Active ingredient **Purposes**
(in each tablet)
Naproxen sodium 220 mg
(naproxen 200 mg) (NSAID) Pain reliever/
fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - headache
 - the common cold
- temporarily reduces fever

Warnings

- Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
- asthma (wheezing)
 - facial swelling
 - hives
 - skin redness
 - rash
 - blisters
- If an allergic reaction occurs, stop use and seek medical help right away.
- Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
- are age 60 or older
 - have had stomach ulcers or bleeding problems
 - take a blood thinning (anticoagulant) or steroid drug
 - take other drugs containing prescription or over-the-counter NSAIDs (aspirin, ibuprofen, naproxen, or others)



Drug Facts (continued)

- have 3 or more alcoholic drinks every day while using this product
 - take more or for a longer time than directed
- Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug


When using this product

- take with food or milk if stomach upset occurs
- you experience any of the following signs of stomach bleed ing:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- stomach pain gets worse or lasts more than 3 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

Drug Facts (continued)

Questions or comments? 1-800-395-0899
(Mon-Fri 9AM - 5PM EST)

Do not use if carton is open or if foil seal imprinted with "SAFETY SQUELSE" on bottle opening is missing or broken.



Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy, unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
 - the smallest effective dose should be used
 - drink a full glass of water with each dose
- Adults and children 12 years and older
- take 1 tablet every 8 to 12 hours while symptoms last
 - take 2 tablets within the first hour
 - do not exceed 2 tablets in any 8- to 12-hour period
 - do not exceed 3 tablets in a 24-hour period

Children under 12 years

- ask a doctor

Other Information

- each tablet contains sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients: FD&C blue #2 lake, hydroxypropylcellulose, magnesium stearate, microcrystalline cellulose, polyvinylpyrrolidone, povidone, talc, titanium dioxide

ALL DAY STRONG®
ALEVE®



naproxen sodium tablets, 220 mg (NSAID)
Pain reliever/fever reducer



STRENGTH TO LAST 12 HOURS

50 TABLETS



Actual Size 100%

(b) (4)

CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	8.1 pt
Drug Facts	N/A	8 pt	8 pt
(continued):	N/A	8 pt	8 pt
Headings:	N/A	8 pt	8 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6-6.5 pt
Characters Per Inch:	N/A	No more than 39	≤ 39




Actual Size 100%



Actual Size 100%

(b) (4)

CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	-
Drug Facts	N/A	8 pt	-
(continued):	N/A	8 pt	-
Headings:	N/A	8 pt	4.5 pt
Subheadings:	N/A	6 pt	4.5 pt
Text:	N/A	6 pt	4.5 pt
Bullets:	N/A	5 pt (no larger)	4 pt
Leading:	N/A	+0.5 pt	4.75 pt
Characters Per Inch:	N/A	No more than 39	>39



Bayer
Made in
Lot
Exp

SafetyNet is a registered trademark of The Procter & Gamble Company. Used with permission.
Bayer, the Bayer Cross, Aleve and All Day Strong are registered trademarks of Bayer.

Dist. by:
Bayer HealthCare LLC
Whispery, NJ 07981

Drug Facts
Active Ingredient
(in each caplet)
Naproxen sodium 220 mg (naproxen 220 mg) (NSAID) *
Purposes
Pain reliever/fever reducer

*nonsteroidal antiinflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - muscular aches
 - menstrual cramps
 - backache
 - headache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, asthma (wheezing), facial swelling, skin redness, shock, or rash.

If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older; have had stomach ulcers or bleeding problems; have alcohol drinking (ethanol) or steroid drug use; take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others); have 3 or more alcoholic drinks every day while using this product; take more or for a longer time than directed.

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug


When using this product:

- take with food or milk if stomach upset occurs
- stop use and ask a doctor if:
 - you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
 - pain gets worse or lasts more than 10 days
 - you have difficulty swallowing
 - it feels like the pill is stuck in your throat
 - redness or swelling is present in the painful area
 - any new symptoms appear

Drug Facts (continued)

Questions or comments? 1-800-395-0688
(Mon-Fri 9AM-5PM EST)

Do not use if carton is open or if foil seal imprinted with "SAFETY SEAL" on bottle opening is missing or broken.



Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
 - the smallest effective dose should be used
 - drink a full glass of water with each dose
- Adults and older than 12 years and older**
- take 1 caplet every 8 to 12 hours while symptoms last
 - for the first dose you may take 2 caplets within the first hour
 - do not exceed 2 caplets in any 8- to 12-hour period
 - do not exceed 3 caplets in a 24-hour period

Other Information

- each caplet contains: sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients: FD&C Blue #2 lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Chloramphenicol: ask a doctor

12 years

ask a doctor


ALL DAY STRONG®
ALEVE®
naproxen sodium tablets, 220 mg (NSAID)
Pain reliever/fever reducer

STRENGTH TO LAST 12 HOURS **50 CAPLETS**
CAPSULE-SHAPED TABLETS




Actual Size 100%

(b) (4)

CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	8.1 pt
Drug Facts	N/A	8 pt	8 pt
(continued):	N/A	8 pt	8 pt
Headings:	N/A	8 pt	8 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6-6.5 pt
Characters Per Inch:	N/A	No more than 39	≤ 39



Actual Size 100%



Actual Size 100%

(b) (4)

Drug Facts Title:	N/A	>8.0 pt	-
Drug Facts	N/A	8 pt	-
(continued):	N/A	8 pt	-
Headings:	N/A	8 pt	-
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	-
Leading:	N/A	+0.5 pt	6.5 pt
Characters Per Inch:	N/A	No more than 39	≤ 39



Actual Size 100%

(b) (4)

CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	8.1 pt
Drug Facts	N/A	8 pt	8 pt
(continued):	N/A	8 pt	8 pt
Headings:	N/A	8 pt	8 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6.5 pt
Characters Per Inch:	N/A	No more than 39	≤ 39