

CONSUMER CARTON
Nasacort Allergy 24 Hour 30 Spray - OUTSIDE CARTON (inside carton/coupon on next page)
 06.28.17
 NAS_Allergy_24hr_30sp_CON_CARTN_FNL4



Sanoofi Regulatory Specifications		Verified Date: 06/28/17
Drug Facts Info		
Drug Facts Title	Font name: Helvetica Neue 77 Bold Condensed Oblique	14 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique / Helvetica Neue 57 Condensed	8.00 point type
Warnings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8.00 point type
Directions	Font name: Helvetica Neue 77 Bold Condensed	6.00 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6.00 point type
Subtitle	Font name: Universal News with Commercial Pi 97	6.00 point type
Buttons on same line, end of statement separated from bulleted statement by two sets		Yes
Spacing of the hairlines from edge of box - i.e. Minimum of 1 space either side of Drug Facts Box		Yes
Tracking	0 to 16 pt.	94% 100%
Leading	Minimum space in body of Drug Facts:	6.25 pt.
	Maximum Characters/line:	32
	Hairlines:	2.5 pt.
		50 pt
Principal Display Panel Info		
Font name of text located in bottom 1/3 of the box (not the height in inches)		125 pt.
PDP dimensions in square inches		12 sq. in.
Font size of Statement of Identity (if not line text, to be measured in Helvetica capital "M")	V 11 pt. H 8.00 pt.	
Font size of Largest Legend Copy or PDP (if not line text, to be measured in Helvetica capital "M")	V 4.5 pt. H 0.00 pt.	
Ratio of Statement of Identity to Largest Legend Copy on PDP		V 25% H 0%

PROFESSIONAL CARTON
Nasacort Allergy 24 Hour 30 Spray - OUTSIDE CARTON (inside carton/coupon on next page)
 06.28.17
 NAS_Allergy_24hr_30sp_PRO_CARTN_FNL4



Sanoofi Regulatory Specifications		Verified Date: 06/28/17
Drug Facts Info		
Drug Facts Title	Font name: Helvetica Neue 77 Bold Condensed Oblique	14 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique / Helvetica Neue 57 Condensed	8.00 point type
Warnings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8.00 point type
Cautioning	Font name: Helvetica Neue 77 Bold Condensed	6.00 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6.00 point type
Substitutes	Font name: Universal News with Commercial Pi 97	6.00 point type
Substitutes on same line, end of statement separated from bulleted statement by two ems		Yes
Spacing of the hairlines from edge of box - i.e. Minimum of 1 space either side of Drug Facts box		Yes
Tracking	0 to 16 pt.	94% 100%
Leading: Minimum space in body of Drug Facts:	6.25 pt.	Maximum Characters/Inch: 32
Trailers:	2.5 pt.	Hairlines: 50 pt
Principal Display Panel Info		
Font name of text located in bottom 6 inches of the outer height in inches		125 in.
PDP dimensions in square inches		12 sq. in.
Font size of Statement of Identity (if not true text, to be measured in Helvetica capital "M")	V 11 pt. H 8.00 pt.	
Font size of Largest Legend Copy or PDP (if not true text, to be measured in Helvetica capital "M")	V 4.5 pt. H 0.20 pt.	
Ratio of Statement of Identity to Largest Legend Copy on PDP		
Statement of Identity gtc (Divided by) Largest Legend Copy gtc		V 25% H 0%

Nasacort Allergy 24 Hour 60 Spray Childrens

06.28.17

NAS_Allergy_24hr_60sp_KIDS_LID_FNL3



Sanofi Regulatory Specifications		Verified Date: 06/28/17
Drug Facts Info		
Drug Facts (Title)	Font name:	N/A
Drug Facts (continued)	Font name:	N/A
Headings	Font name:	N/A
Subheadings	Font name:	N/A
Body Text	Font name:	N/A
Bullets	Font name:	N/A
Bullets on same lines: end of statement separated from bulleted statement by two ems		N/A
Spacing of the hairlines from edge of box – i.e. Minimum of 2 spaces either side of Drug Facts Box		N/A
Tracking:	N/A	Horizontal Scale: N/A
Leading (Minimum space in body of Drug Facts):	N/A	Maximum Characters/Inch: N/A
Barlines:	N/A	Hairlines: N/A
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		.125 in.
PDP dimensions (in square inches)		19 sq. in.
Font size of Statement of Identity (If not live text, to be measured in Helvetica capital "M")		V 14.5 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (If not live text, to be measured in Helvetica capital "M")		V 58 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		
Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		V 25% H 0%

Nasacort Allergy 24 Hour 60 Spray

06.28.17

NAS_Allergy_24hr_60sp_LID_FNL3



Sanofi Regulatory Specifications		Verified Date: 06/28/17
Drug Facts Info		
Drug Facts (Title)	Font name:	N/A
Drug Facts (continued)	Font name:	N/A
Headings	Font name:	N/A
Subheadings	Font name:	N/A
Body Text	Font name:	N/A
Bullets	Font name:	N/A
Bullets on same lines: end of statement separated from bulleted statement by two ems		N/A
Spacing of the hairlines from edge of box – i.e. Minimum of 2 spaces either side of Drug Facts Box		N/A
Tracking:	N/A	Horizontal Scale: N/A
Leading (Minimum space in body of Drug Facts):	N/A	Maximum Characters/Inch: N/A
Barlines:	N/A	Hairlines: N/A
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		.125 in.
PDP dimensions (in square inches)		19 sq. in.
Font size of Statement of Identity (If not live text, to be measured in Helvetica capital "M")		V 14.5 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (If not live text, to be measured in Helvetica capital "M")		V 58 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		
Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		V 25% H 0%

Nasacort Allergy 24 Hour 120 Spray

06.28.17

NAS_Allergy_24hr_120sp_LID_FNL3



Sanofi Regulatory Specifications		Verified Date: 06/28/17
Drug Facts Info		
<i>Drug Facts</i> (Title)	Font name:	N/A
<i>Drug Facts</i> (continued)	Font name:	N/A
Headings	Font name:	N/A
Subheadings	Font name:	N/A
Body Text	Font name:	N/A
Bullets	Font name:	N/A
Bullets on same lines: end of statement separated from bulleted statement by two ems		N/A
Spacing of the hairlines from edge of box – i.e. Minimum of 2 spaces either side of Drug Facts Box		N/A
Tracking:	N/A	Horizontal Scale: N/A
Leading (Minimum space in body of Drug Facts):	N/A	Maximum Characters/Inch: N/A
Barlines:	N/A	Hairlines: N/A
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		.125 in.
PDP dimensions (in square inches)		19 sq. in.
Font size of Statement of Identity (If not live text, to be measured in Helvetica capital "M")		V 14.5 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (If not live text, to be measured in Helvetica capital "M")		V 58 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		
Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		V 25% H 0%

Nasacort Allergy 24 Hour 120+30 Spray Bonus Pack

06.28.17

NAS_Allergy_24hr_120+30sp_Bonus-Pk_LID_FNL4



Sanofi Regulatory Specifications		Verified Date: 06/28/17
Drug Facts Info		
Drug Facts (Title)	Font name:	N/A
Drug Facts (continued)	Font name:	N/A
Headings	Font name:	N/A
Subheadings	Font name:	N/A
Body Text	Font name:	N/A
Bullets	Font name:	N/A
Bullets on same lines: end of statement separated from bulleted statement by two ems		N/A
Spacing of the hairlines from edge of box – i.e. Minimum of 2 spaces either side of Drug Facts Box		N/A
Tracking:	N/A	Horizontal Scale: N/A
Leading (Minimum space in body of Drug Facts):	N/A	Maximum Characters/Inch: N/A
Barlines:	N/A	Hairlines: N/A
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		.125 in.
PDP dimensions (in square inches)		19 sq. in.
Font size of Statement of Identity (If not live text, to be measured in Helvetica capital "M")		V 14.5 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (If not live text, to be measured in Helvetica capital "M")		V 58 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		
Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		V 25% H 0%

Nasacort Allergy 24 Hour 120+30 Spray Multi-Pack

06.28.17

NAS_Allergy_24hr_120+30sp_Multi-Pk_LID_FNL3



Sanofi Regulatory Specifications		Verified Date: 06/28/17
Drug Facts Info		
Drug Facts (Title)	Font name:	N/A
Drug Facts (continued)	Font name:	N/A
Headings	Font name:	N/A
Subheadings	Font name:	N/A
Body Text	Font name:	N/A
Bullets	Font name:	N/A
Bullets on same lines: end of statement separated from bulleted statement by two ems		N/A
Spacing of the hairlines from edge of box – i.e. Minimum of 2 spaces either side of Drug Facts Box		N/A
Tracking:	N/A	Horizontal Scale: N/A
Leading (Minimum space in body of Drug Facts):	N/A	Maximum Characters/Inch: N/A
Barlines:	N/A	Hairlines: N/A
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		.125 in.
PDP dimensions (in square inches)		19 sq. in.
Font size of Statement of Identity (If not live text, to be measured in Helvetica capital "M")		V 14.5 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (If not live text, to be measured in Helvetica capital "M")		V 58 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		
Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		V 25% H 0%

Nasacort Allergy 24 Hour 240 Spray Multi-Pack

06.28.17

NAS_Allergy_24hr_240sp_Multi-Pk_LID_FNL3



Sanofi Regulatory Specifications		Verified Date: 06/28/17
Drug Facts Info		
Drug Facts (Title)	Font name:	N/A
Drug Facts (continued)	Font name:	N/A
Headings	Font name:	N/A
Subheadings	Font name:	N/A
Body Text	Font name:	N/A
Bullets	Font name:	N/A
Bullets on same lines: end of statement separated from bulleted statement by two ems		N/A
Spacing of the hairlines from edge of box – i.e. Minimum of 2 spaces either side of Drug Facts Box		N/A
Tracking:	N/A	Horizontal Scale: N/A
Leading (Minimum space in body of Drug Facts):	N/A	Maximum Characters/Inch: N/A
Barlines:	N/A	Hairlines: N/A
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		.125 in.
PDP dimensions (in square inches)		19 sq. in.
Font size of Statement of Identity (If not live text, to be measured in Helvetica capital "M")		V 14.5 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (If not live text, to be measured in Helvetica capital "M")		V 58 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		
Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		V 25% H 0%

Nasacort Allergy 24 Hour 240 Spray Multi-Pack - ALTERNATE

06.28.17


NAS_Allergy_24hr_240sp_Multi-Pk_LID_ALT_FNL3



Sanofi Regulatory Specifications		Verified Date: 06/28/17
Drug Facts Info		
<i>Drug Facts</i> (Title)	Font name:	N/A
<i>Drug Facts</i> (continued)	Font name:	N/A
Headings	Font name:	N/A
Subheadings	Font name:	N/A
Body Text	Font name:	N/A
Bullets	Font name:	N/A
Bullets on same lines: end of statement separated from bulleted statement by two ems		N/A
Spacing of the hairlines from edge of box – i.e. Minimum of 2 spaces either side of Drug Facts Box		N/A
Tracking:	N/A	Horizontal Scale: N/A
Leading (Minimum space in body of Drug Facts):	N/A	Maximum Characters/Inch: N/A
Barlines:	N/A	Hairlines: N/A
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		.125 in.
PDP dimensions (in square inches)		19 sq. in.
Font size of Statement of Identity (If not live text, to be measured in Helvetica capital "M")		V 14.5 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (If not live text, to be measured in Helvetica capital "M")		V 58 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		
Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		V 25% H 0%

06.28.17

NAS_Allergy_24hr_360sp_Club-Pk_CARD_FNL4



Sanofi Regulatory Specifications		Verified Date: 06/28/17
Drug Facts Info		
Drug Facts (Title)	Font name:	N/A
Drug Facts (continued)	Font name:	N/A
Headings	Font name:	N/A
Subheadings	Font name:	N/A
Body Text	Font name:	N/A
Bullets	Font name:	N/A
Bullets on same lines: end of statement separated from bulleted statement by two ems		N/A
Spacing of the hairlines from edge of box – i.e. Minimum of 2 spaces either side of Drug Facts Box		N/A
Tracking:	N/A	Horizontal Scale:
Leading (Minimum space in body of Drug Facts):	N/A	Maximum Characters/Inch:
Barlines:	N/A	Hairlines:
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		.138 in
PDP dimensions (in square inches)		48 sq. in.
Font size of Statement of Identity (If not live text, to be measured in Helvetica capital "M")		V 16 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (If not live text, to be measured in Helvetica capital "M")		V 61 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		
Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		V 26% H 0%

06.28.17

NAS_Allergy_24hr_360sp_Club-Pk_CARD_FNL4

LOT _____
 EXP _____

<p>Drug Facts</p> <p>Active ingredient (in each spray) Purpose Triamcinolone acetonide Allergy symptom reliever 55 mcg (glucocorticoid)</p> <p>Uses temporarily relieves these symptoms of hay fever or other upper respiratory allergies: ■ nasal congestion ■ runny nose ■ sneezing ■ itchy nose</p> <p>Warnings Do not use ■ in children under 2 years of age ■ if you have ever had an allergic reaction to any of the ingredients</p> <p>Ask a doctor before use if you ■ have had recent nose ulcers or nose surgery ■ have had a nose injury that has not healed ■ are using a steroid medicine for asthma, allergies or skin rash ■ have an eye infection ■ have or had glaucoma or cataracts</p> <p>When using this product ■ the growth rate of some children may be slower ■ some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief. ■ do not share this bottle with anyone else as this may spread germs ■ remember to tell your doctor about all the medicines you take, including this one</p> <p>Stop use and ask a doctor if ■ you have, or come into contact with someone who has, chickenpox, measles or tuberculosis ■ you have or develop symptoms of an infection such as a persistent fever ■ you have any change in vision ■ you have severe or frequent nosebleeds</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>	<p>Drug Facts (continued)</p> <p>Directions Read insert (inside package) on how to: ■ get a new bottle ready (primed) before first use ■ prime bottle again if not used for more than 2 weeks ■ use the spray ■ clean the spray nozzle</p> <p style="text-align: center;">ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER</p> <p>adults and children 12 years of age and older ■ once daily, spray 2 times into each nostril while sniffing gently ■ once your allergy symptoms improve, reduce to 1 spray in each nostril per day</p> <p style="text-align: center;">CHILDREN 2 TO UNDER 12 YEARS OF AGE</p> <p>■ the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.</p> <p>children 6 to under 12 years of age ■ an adult should supervise use ■ once daily, spray 1 time into each nostril while sniffing gently ■ if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.</p> <p>children 2 to under 6 years of age ■ an adult should supervise use ■ once daily, spray 1 time into each nostril while sniffing gently</p> <p>children under 2 years of age ■ do not use</p> <p>■ do not use more than directed ■ if you forget a dose, do not double the next dose ■ do not spray into eyes or mouth ■ if allergy symptoms do not improve after one week, stop using and talk to a doctor ■ do not use for the common cold ■ shake well before each use</p> <p>Other information ■ do not use if sealed package is torn or opened ■ keep package and insert. They contain important information. ■ store between 20°-25°C (68°-77°F)</p> <p>Inactive ingredients benzalkonium chloride, carbomethylcellulose sodium, dextrose, edetate disodium, hydrochloric acid or sodium hydroxide (for pH adjustment), microcrystalline cellulose, polysorbate 80, purified water</p> <p>Questions or comments? call toll-free 1-800-633-1610</p>
---	---

Dist. by: Chatterm, Inc., a Sanofi Company
 Chattanooga, TN 37409-0219 © 2017
 Origin France 0000000-01

Sanofi Regulatory Specifications		Verified Date: 06/28/17
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	14 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique / Helvetica Neue 57 Condensed	8.00 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8.00 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6.00 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6.00 point type
Bullets	Font name: Universal News with Commercial Pi 97	6.00 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		Yes
Spacing of the hairlines from edge of box – i.e. Minimum of 2 spaces either side of Drug Facts Box		Yes
Tracking:	0 to 16 pt.	Horizontal Scale: 94%-100%
Leading (Minimum space in body of Drug Facts):	6.5 pt.	Maximum Characters/Inch: 29
Barlines:	2.5 pt.	Hairlines: .50 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		.138 in
PDP dimensions (in square inches)		48 sq. in.
Font size of Statement of Identity (If not live text, to be measured in Helvetica capital "M")		V 16 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (If not live text, to be measured in Helvetica capital "M")		V 61 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		
Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		V 26% H 0%

Nasacort Allergy 24 Hour 1"x1" IRC

06.27.17

NAS_1x1_IRC_FNL3



MANUFACTURER'S COUPON EXPIRATION DATE: 00/00/00

Consumer: There is a limit of one coupon per purchase, and the coupon can't be combined with other offers. Good only in the U.S. and overseas commissaries on the purchase of the specified product/size. Void where prohibited or if altered, copied, transferred, auctioned, or sold. You must pay sales tax. Cash value: 1/100 of 1¢. **Retailer:** We will reimburse you for the face value of this coupon plus 8¢ if redeemed in compliance with our Coupon Reimbursement Policy (available at address below). Invoices proving purchase of sufficient stock to cover coupons presented for redemption must be provided upon request. Failure to follow the policy may constitute fraud and may void all coupons submitted. Mail to: Chatterm, Inc. 1050, NCH Marketing Services, P.O. Box 880001, El Paso, TX 88588-0001. ©2017 Chatterm, Inc. 0000000-00

Children's Nasacort Allergy 24 Hour Consumer Information Leaflet

06.27.17

NAS_Kids_Consumer_Info_Leaflet_FNL4




IMPORTANT INFORMATION

- Read both sides of this insert for complete instructions on how to get a bottle ready (primed), how to use the spray bottle and how to clean the spray nozzle.
- Keep this insert as it contains important information.

STEPS TO GET A NEW BOTTLE READY FOR USE (PRIMED)

1. Before first use, a new bottle must be primed.

- Remove cap (Figure A).
- Shake bottle.
- Press and release spray nozzle until a fine mist is produced (as shown in Figure B). This may take several times. Take care not to spray in face.



PRIME BOTTLE AGAIN IF NOT USED FOR MORE THAN 2 WEEKS


- Repeat steps for priming a bottle (see above).

USE INSTRUCTIONS

- Blow nose gently to clear nostrils.
- Remove cap, then shake bottle.
- Hold bottle with thumb under bottle and spray nozzle between fingers (as shown in Figure B).
- Press against the outside of your nose with your finger to close off one nostril (as shown in Figure C). The spray nozzle should not reach far into the nose. Aim nozzle toward back of nose (as shown in Figure C).



DO NOT spray toward nasal septum (the wall between the 2 nostrils) (as shown in Figure D).



- While sniffing gently, spray into nostril.

IMPORTANT: For complete dosing instructions, See "Directions for use" on next side.

- Repeat steps "4" through "6" for the other nostril.
- After using the nasal spray, wipe nozzle with a tissue and replace cap by pressing it down over the spray nozzle.

NOTE: Avoid blowing nose for 15 minutes after use.

If nozzle does not spray properly, see cleaning instructions on next side.

(continued on side 2)

SIDE 1

DIRECTIONS FOR USE

- BEFORE first use, you must get a new bottle ready (primed).**
- If bottle is not used for more than 2 weeks, prime bottle again.**

Dosing

ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER

- | | |
|---|--|
| adults and children 12 years of age and older | <ul style="list-style-type: none"> once daily, spray 2 times into each nostril while sniffing gently once your allergy symptoms improve, reduce to 1 spray in each nostril per day |
|---|--|

CHILDREN 2 TO UNDER 12 YEARS OF AGE

- the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.

- | | |
|-------------------------------------|--|
| children 6 to under 12 years of age | <ul style="list-style-type: none"> an adult should supervise use once daily, spray 1 time into each nostril while sniffing gently if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day. |
|-------------------------------------|--|

- | | |
|------------------------------------|---|
| children 2 to under 6 years of age | <ul style="list-style-type: none"> an adult should supervise use once daily, spray 1 time into each nostril while sniffing gently |
|------------------------------------|---|

- | | |
|-------------------------------|--|
| children under 2 years of age | <ul style="list-style-type: none"> do not use |
|-------------------------------|--|

Other important information for use

- Do not use more than directed.
- If you forget a dose, do not double the next dose.
- Do not spray into eyes or mouth.
- If you get the spray in your eyes, rinse well with water.
- If allergy symptoms do not improve after one week, stop using and talk to a doctor.
- Do not share this bottle with anyone else as this may spread germs.


IF PUMP DOES NOT SPRAY PROPERLY, THE NOZZLE MAY BE BLOCKED

- Never try to unblock nozzle with a pin or any other object (as shown in Figure E).
- Clean the nozzle as shown below.



CLEANING INSTRUCTIONS

- Gently pull spray nozzle away from bottle (as shown in Figure F).
- Rinse **SPRAY NOZZLE ONLY** under warm water (as shown in Figure G).
- Shake or tap to remove excess water.
- Re-attach spray nozzle to bottle.
- Press and release spray nozzle until a fine spray is produced, taking care not to spray in face.



Children's Nasacort® Allergy 24HR pump is now ready to use.

Where can I get more information?

1-800-633-1610

Store between 20°-25°C (68°-77°F)

Distributed by: Chattem, Inc., a Sanofi Company
Chattanooga, TN 37409-0219 XXXXXXX-XX © 2017 Chattem, Inc.

SIDE 2

Nasacort Allergy 24 Hour Consumer Information Leaflet

06.27.17

NAS_Consumer_Info_Leaflet_FNL4




IMPORTANT INFORMATION

- Read both sides of this insert for complete instructions on how to get a bottle ready (primed), how to use the spray bottle and how to clean the spray nozzle.
- Keep this insert as it contains important information.

STEPS TO GET A NEW BOTTLE READY FOR USE (PRIMED)

1. Before first use, a new bottle must be primed.

- Remove cap (Figure A).
- Shake bottle.
- Press and release spray nozzle until a fine mist is produced (as shown in Figure B). This may take several times. Take care not to spray in face.



PRIME BOTTLE AGAIN IF NOT USED FOR MORE THAN 2 WEEKS

- Repeat steps for priming a bottle (see above).

USE INSTRUCTIONS

- Blow nose gently to clear nostrils.
- Remove cap, then shake bottle.
- Hold bottle with thumb under bottle and spray nozzle between fingers (as shown in Figure B).
- Press against the outside of your nose with your finger to close off one nostril (as shown in Figure C).
- Place the tip of the spray nozzle into the other nostril. The spray nozzle should not reach far into the nose. Aim nozzle toward back of nose (as shown in Figure C).



DO NOT spray toward nasal septum (the wall between the 2 nostrils) (as shown in Figure D).



- While sniffing gently, spray into nostril.

IMPORTANT: For complete dosing instructions, See "Directions for use" on next side.

- Repeat steps "4" through "6" for the other nostril.
- After using the nasal spray, wipe nozzle with a tissue and replace cap by pressing it down over the spray nozzle.

NOTE: Avoid blowing nose for 15 minutes after use.

If nozzle does not spray properly, see cleaning instructions on next side.

(continued on side 2)

SIDE 1

DIRECTIONS FOR USE

- BEFORE first use, you must get a new bottle ready (primed).**
- If bottle is not used for more than 2 weeks, prime bottle again.**

Dosing

ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER

- | | |
|---|--|
| adults and children 12 years of age and older | <ul style="list-style-type: none"> once daily, spray 2 times into each nostril while sniffing gently once your allergy symptoms improve, reduce to 1 spray in each nostril per day |
|---|--|

CHILDREN 2 TO UNDER 12 YEARS OF AGE

- the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.

- | | |
|-------------------------------------|--|
| children 6 to under 12 years of age | <ul style="list-style-type: none"> an adult should supervise use once daily, spray 1 time into each nostril while sniffing gently if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day. |
|-------------------------------------|--|

- | | |
|------------------------------------|---|
| children 2 to under 6 years of age | <ul style="list-style-type: none"> an adult should supervise use once daily, spray 1 time into each nostril while sniffing gently |
|------------------------------------|---|

- | | |
|-------------------------------|--|
| children under 2 years of age | <ul style="list-style-type: none"> do not use |
|-------------------------------|--|

Other important information for use

- Do not use more than directed.
- If you forget a dose, do not double the next dose.
- Do not spray into eyes or mouth.
- If you get the spray in your eyes, rinse well with water.
- If allergy symptoms do not improve after one week, stop using and talk to a doctor.
- Do not share this bottle with anyone else as this may spread germs.

IF PUMP DOES NOT SPRAY PROPERLY, THE NOZZLE MAY BE BLOCKED

- Never try to unblock nozzle with a pin or any other object (as shown in Figure E).
- Clean the nozzle as shown below.



CLEANING INSTRUCTIONS

- Gently pull spray nozzle away from bottle (as shown in Figure F).
- Rinse **SPRAY NOZZLE ONLY** under warm water (as shown in Figure G).
- Shake or tap to remove excess water.
- Re-attach spray nozzle to bottle.
- Press and release spray nozzle until a fine spray is produced, taking care not to spray in face.



Nasacort® Allergy 24HR pump is now ready to use.

Where can I get more information?

1-800-633-1610

Store between 20°-25°C (68°-77°F)

Distributed by: Chattem, Inc., a Sanofi Company
Chattanooga, TN 37409-0219 XXXXXXX-XX © 2017 Chattem, Inc.

SIDE 2

Nasacort Allergy 24 Hour Consumer Information Leaflet

07.06.17

NAS_Consumer_Info_Leaflet_with CPN_FNL5



Nasacort®
ALLERGY 24HR

COUPON
BELOW

IMPORTANT INFORMATION


- Read both sides of this insert for complete instructions on how to get a bottle ready (primed), how to use the spray bottle and how to clean the spray nozzle.
- Keep this insert as it contains important information.

STEPS TO GET A NEW BOTTLE READY FOR USE (PRIMED)

1. Before first use, a new bottle must be primed.

- Remove cap (Figure A).
- Shake bottle.
- Press and release spray nozzle until a fine mist is produced (as shown in Figure B). This may take several times. Take care not to spray in face.

Figure A



Cap

Spray Nozzle

Bottle

PRIME BOTTLE AGAIN IF NOT USED FOR MORE THAN 2 WEEKS

- Repeat steps for priming a bottle (see above).

USE INSTRUCTIONS

- Blow nose gently to clear nostrils.
- Remove cap, then shake bottle.
- Hold bottle with thumb under bottle and spray nozzle between fingers (as shown in Figure B).
- Press against the outside of your nose with your finger to close off one nostril (as shown in Figure C).
- Place the tip of the spray nozzle into the other nostril. The spray nozzle should not reach far into the nose. Aim nozzle toward back of nose (as shown in Figure C).

Figure B



Figure C



Figure D



DO NOT spray toward nasal septum (the wall between the 2 nostrils) (as shown in Figure D).

Correct Position **Wrong Position**

- While sniffing gently, spray into nostril.

IMPORTANT: For complete dosing instructions, See "Directions for use" on next side.

- Repeat steps "4" through "6" for the other nostril.
- After using the nasal spray, wipe nozzle with a tissue and replace cap by pressing it down over the spray nozzle.

NOTE: Avoid blowing nose for 15 minutes after use.

If nozzle does not spray properly, see cleaning instructions on next side.

(continued on side 2)

SIDE 1

SAVE \$X

on any one
Nasacort® Allergy 24HR
120 spray or larger

Use as directed. ©2017 Chatterm, Inc.

DIRECTIONS FOR USE

- BEFORE first use, you must get a new bottle ready (primed).**
- If bottle is not used for more than 2 weeks, prime bottle again.**

Dosing	
ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER	
adults and children 12 years of age and older	<ul style="list-style-type: none"> once daily, spray 2 times into each nostril while sniffing gently once your allergy symptoms improve, reduce to 1 spray in each nostril per day
CHILDREN 2 TO UNDER 12 YEARS OF AGE	
<ul style="list-style-type: none"> the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year. 	
children 6 to under 12 years of age	<ul style="list-style-type: none"> an adult should supervise use once daily, spray 1 time into each nostril while sniffing gently if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.
children 2 to under 6 years of age	<ul style="list-style-type: none"> an adult should supervise use once daily, spray 1 time into each nostril while sniffing gently
children under 2 years of age	<ul style="list-style-type: none"> do not use

Other important information for use

- Do not use more than directed.
- If you forget a dose, do **not** double the next dose.
- Do not spray into eyes or mouth.
- If you get the spray in your eyes, rinse well with water.
- If allergy symptoms do not improve after one week, stop using and talk to a doctor.
- Do not share this bottle with anyone else as this may spread germs.

IF PUMP DOES NOT SPRAY PROPERLY, THE NOZZLE MAY BE BLOCKED

- Never try to unblock nozzle with a pin or any other object (as shown in Figure E).
- Clean the nozzle as shown below.

Figure E



CLEANING INSTRUCTIONS

- Gently pull spray nozzle away from bottle (as shown in Figure F).
- Rinse **SPRAY NOZZLE ONLY** under warm water (as shown in Figure G).
- Shake or tap to remove excess water.
- Re-attach spray nozzle to bottle.
- Press and release spray nozzle until a fine spray is produced, taking care not to spray in face.

Figure F



Figure G



Nasacort® Allergy 24HR pump is now ready to use.

Where can I get more information?
1-800-633-1610

Store between 20°-25°C (68°-77°F)
Distributed by: Chatterm, Inc., a Sanofi Company
Chattanooga, TN 37409-0219 XXXXXXX-XX © 2017 Chatterm, Inc.

SIDE 2

MANUFACTURER'S COUPON | EXPIRES: XX/XX/XX

Consumer: There is a limit of one coupon per purchase, and the coupon can't be combined with other offers. Good only in the U.S. and overseas commissaries on the purchase of the specified product/size. Void where prohibited or if altered, copied, transferred, auctioned, or sold. You must pay sales tax. Cash value: 1/100 of 1¢. Retailer: We will reimburse you for the face value of this coupon plus 8¢ if redeemed in compliance with our Coupon Reimbursement Policy (available at address below). Invoices proving purchase of sufficient stock to cover coupons presented for redemption must be provided upon request. Failure to follow the policy may constitute fraud and may void all coupons submitted. Mail to: Chatterm, Inc. 1050, NCH Marketing Services, P.O. Box 880001, El Paso, TX 88568-0001



Nasacort Allergy 24 Hour Drug Facts Label

11.13.17
NAS_60sp_KIDS_PeelAway_DFL_FNL2

Drug Facts

Active ingredient (in each spray) *Purpose*
Triamcinolone acetate 50 mcg (1.0 mg/0.1 mL) Allergy symptoms reliever

Uses Temporarily relieves these symptoms of hay fever or other upper respiratory allergies: **nasal congestion** **runny nose** **sneezing** **itchy nose**

Warnings
Do not use

- in children under 2 years of age
- if you have ever had an allergic reaction to any of the ingredients

Ask a doctor before use if you

- have had recent nose ulcers or nose surgery
- have had a nose injury that has not healed
- are using a steroid medicine for asthma, allergies or skin rash
- have an eye infection
- have or had glaucoma or cataracts

When using this product

- the growth rate of some children may be slower
- some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief.
- do not share this bottle with anyone else as this may spread germs
- remember to tell your doctor about all the medicines you take, including this one

Dist. by: Chatter, Inc., a Sanofi Company, Chattanooga, TN 37409-0219 ©2017 Origin France

00000-01

PEEL HERE

Drug Facts (continued)

Stop use and ask a doctor if

- you have, or come into contact with someone who has, chickenpox, measles or tuberculosis
- you have or develop symptoms of an infection such as a persistent fever
- you have any change in vision
- you have severe or frequent nosebleeds

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
Read insert (inside package) on how to use the spray

- once daily, spray 2 times into each nostril while sniffing gently
- get a new bottle ready (primed) before first use
- prime bottle again if not used for more than 2 weeks
- clean the spray nozzle

ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER

- once daily, spray 2 times into each nostril while sniffing gently
- once your allergy symptoms improve, reduce to 1 spray in each nostril per day

CHILDREN 2 TO UNDER 12 YEARS OF AGE

- once daily, spray 1 time into each nostril while sniffing gently

Children 6 to under 12 years of age

- an adult should supervise use

Drug Facts (continued)

CHILDREN 2 TO UNDER 12 YEARS OF AGE (continued)

children 6 to under 12 years of age (continued)

- if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.

children 2 to under 6 years of age (continued)

- an adult should supervise use
- once daily, spray 1 time into each nostril while sniffing gently

children under 2 years of age (continued)

- do not use

Do not use more than directed

- if you forget a dose, do not double the next dose
- do not spray into eyes or mouth
- if allergy symptoms do not improve after one week, stop using and talk to a doctor
- do not use for the common cold
- shake well before each use

Other information

- do not use if sealed package is torn or opened
- avoid package heat (heat). They contain important information.
- store between 20°-25°C (68°-77°F)

Inactive ingredients
benzalkonium chloride, carbonylmethylcellulose sodium, dextrase, edetate disodium, hydrochloric acid or sodium hydroxide (for pH adjustment), microcrystalline cellulose, polyborate 80, purified water

Questions or comments? call toll-free 1-800-633-1610

Sanofi Regulatory Specifications		Verified Date: 11/13/17
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	14.00 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique / Helvetica Neue 57 Condensed	8.00 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8.00 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6.00 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6.00 point type
Bullets	Font name: Universal News with Commercial PI 97	6.00 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		Yes
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		Yes
Tracking:	0 to 16 Horizontal Scale:	94%-100%
Leading (Minimum space in body of Drug Facts):	6.3 pt. Maximum Characters/Inch (Characters plus spaces):	32
Barlines:	2.0 pt. Hairlines:	0.50 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		NA
PDP dimensions (in square inches)		NA
Font size of Statement of Identity (if not live text, to be measured in Helvetica capital "M")		NA
Font size of Logo/Largest Copy on PDP (if not live text, to be measured in Helvetica capital "M")		NA
Ratio of Statement of Identity to Logo/Largest Copy on PDP		NA
Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		NA

Nasacort Allergy 24 Hour Drug Facts Label

11.13.17

NAS_60sp_PeelAway_DFL_FNL6

Drug Facts

Active ingredient (in each spray) **Purpose**
Triamcinolone acetate 50 mcg (1.0 mg) per dose Allergy symptom reliever

Uses Temporarily relieves these symptoms of hay fever or other upper respiratory allergies:
 ■ nasal congestion ■ runny nose ■ sneezing ■ itchy nose

Warnings
 Do not use
 ■ in children under 2 years of age
 ■ if you have ever had an allergic reaction to any of the ingredients

Ask a doctor before use if you
 ■ have had recent nose ulcers or nose surgery
 ■ have had a nose injury that has not healed
 ■ are using a steroid medicine for asthma, allergies or skin rash
 ■ have an eye infection ■ have or had glaucoma or cataracts

When using this product
 ■ the growth rate of some children may be slower
 ■ some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief.
 ■ do not share this bottle with anyone else as this may spread germs
 ■ remember to tell your doctor about all the medicines you take, including this one

Dist. by: Chatter, Inc., a Sanofi Company, Chattanooga, TN 37409-0219 ©2017 Origin France

00000-01

PEEL HERE

Drug Facts (continued)

Stop use and ask a doctor if
 ■ you have, or come into contact with someone who has, chickenpox, measles or tuberculosis
 ■ you have or develop symptoms of an infection such as a persistent fever
 ■ you have any change in vision
 ■ you have severe or frequent nosebleeds

Directions
 Read insert (inside package) on how to use the spray
 ■ if pregnant or breast-feeding, ask a health professional before use
 ■ keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER
 ■ once daily, spray 2 times into each nostril while sniffing gently
 ■ once your allergy symptoms improve, reduce to 1 spray in each nostril per day

CHILDREN 2 TO UNDER 12 YEARS OF AGE
 ■ the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.
 ■ once daily, spray 1 time into each nostril while sniffing gently

Drug Facts (continued)

CHILDREN 2 TO UNDER 12 YEARS OF AGE (continued)
 ■ if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.
 ■ an adult should supervise use
 ■ once daily, spray 1 time into each nostril while sniffing gently

CHILDREN UNDER 2 YEARS OF AGE
 ■ do not use
 ■ do not use more than directed
 ■ if you forget a dose, do not double the next dose
 ■ do not spray into eyes or mouth
 ■ if allergy symptoms do not improve after one week, stop using and talk to a doctor
 ■ do not use for the common cold
 ■ shake well before each use

Other information
 ■ do not use if sealed package is torn or opened
 ■ store between 20°-25°C (68°-77°F)
 ■ store between 20°-25°C (68°-77°F)

Inactive ingredients
 benzalkonium chloride, carbonylmethylcellulose sodium, dextrase, edetate disodium, hydrochloric acid or sodium hydroxide (for pH adjustment), microcrystalline cellulose, polyborate 80, purified water

Questions or comments? call toll-free 1-800-633-1610

Sanofi Regulatory Specifications		Verified Date: 11/13/17
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	14.00 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique / Helvetica Neue 57 Condensed	8.00 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8.00 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6.00 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6.00 point type
Bullets	Font name: Universal News with Commercial PI 97	6.00 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		Yes
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		Yes
Tracking:	0 to 16 Horizontal Scale:	94%-100%
Leading (Minimum space in body of Drug Facts):	6.3 pt. Maximum Characters/Inch (Characters plus spaces):	32
Barlines:	2.0 pt. Hairlines:	0.50 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		NA
PDP dimensions (in square inches)		NA
Font size of Statement of Identity (if not live text, to be measured in Helvetica capital "M")		NA
Font size of Logo/Largest Copy on PDP (if not live text, to be measured in Helvetica capital "M")		NA
Ratio of Statement of Identity to Logo/Largest Copy on PDP		NA
Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		NA

Nasacort Allergy 24 Hour Drug Facts Label

11.13.17
NAS_120sp_PeelAway DFL_FNL2

Drug Facts

Active ingredient (in each spray) Fluticasone propionate 50 mcg (1.0 mg) per spray
Purpose Allergy symptom reliever

Uses Temporarily relieves these symptoms of hay fever or other upper respiratory allergies: **nasal congestion**, **runny nose**, **sneezing**, **itchy nose**

Warnings
Do not use

- in children under 2 years of age
- if you have ever had an allergic reaction to any of the ingredients

Ask a doctor before use if you

- have had recent nose ulcers or nose surgery
- have had a nose injury that has not healed
- are using a steroid medicine for asthma, allergies or skin rash
- have an eye infection
- have or had glaucoma or cataracts

When using this product

- the growth rate of some children may be slower
- some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief.
- do not share this bottle with anyone else as this may spread germs
- remember to tell your doctor about all the medicines you take, including this one

Dist. by: Chatter, Inc., a Sanofi Company, Chattanooga, TN 37409-0219 ©2017 Origin France

000000-01

PEEL HERE

Drug Facts (continued)

Stop use and ask a doctor if

- you have, or come into contact with someone who has, chickenpox, measles or tuberculosis
- you have or develop symptoms of an infection such as a persistent fever
- you have any change in vision
- you have severe or frequent nosebleeds

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
Read insert (inside package) on how to use the spray

- if pregnant or breast-feeding, ask a health professional before use
- if you are pregnant, get a health professional's advice before using this medicine
- if you are breastfeeding, ask a health professional before use
- if you are pregnant, get a health professional's advice before using this medicine
- if you are breastfeeding, ask a health professional before use
- if you are pregnant, get a health professional's advice before using this medicine
- if you are breastfeeding, ask a health professional before use

ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER

- once daily, spray 2 times into each nostril while sniffing gently

CHILDREN 2 TO UNDER 12 YEARS OF AGE

- once daily, spray 1 time into each nostril while sniffing gently

Drug Facts (continued)

CHILDREN 2 TO UNDER 12 YEARS OF AGE (continued)

children 6 to under 12 years of age (continued)

- if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.

children 2 to under 6 years of age (continued)

- an adult should supervise use
- once daily, spray 1 time into each nostril while sniffing gently

children under 2 years of age (continued)

- do not use

Other information

- do not use more than directed
- if you forget a dose, do not double the next dose
- do not spray into eyes or mouth
- if allergy symptoms do not improve after one week, stop using and talk to a doctor
- do not use for the common cold
- shake well before each use

Inactive ingredients
benzalkonium chloride, carbonylmethylcellulose sodium, dextrose, edetate disodium, hydrochloric acid or sodium hydroxide (for pH adjustment), microcrystalline cellulose, polyborate 80, purified water

Questions or comments? call toll-free 1-800-633-1610

Sanofi Regulatory Specifications		Verified Date: 11/13/17
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	14.00 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique / Helvetica Neue 57 Condensed	8.00 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8.00 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6.00 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6.00 point type
Bullets	Font name: Universal News with Commercial PI 97	6.00 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		Yes
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		Yes
Tracking:	0 to 16 Horizontal Scale:	94%-100%
Leading (Minimum space in body of Drug Facts):	6.3 pt. Maximum Characters/Inch (Characters plus spaces):	32
Barlines:	2.0 pt. Hairlines:	0.50 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		NA
PDP dimensions (in square inches)		NA
Font size of Statement of Identity (if not live text, to be measured in Helvetica capital "M")		NA
Font size of Logo/Largest Copy on PDP (if not live text, to be measured in Helvetica capital "M")		NA
Ratio of Statement of Identity to Logo/Largest Copy on PDP		NA
Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		NA

Nasacort Allergy 24 Hour Drug Facts Label

11.13.17

NAS_120+30sp_Bonus-Pk_PeelAway_DFL_FNL2

Drug Facts

Active ingredient (in each spray) *Fluticasone propionate* **Purpose** Allergy symptom reliever

Uses Temporarily relieves these symptoms of hay fever or other upper respiratory allergies: **nasal congestion** **runny nose** **sneezing** **itchy nose**

Warnings

Do not use

- in children under 2 years of age
- if you have ever had an allergic reaction to any of the ingredients

Ask a doctor before use if you:

- have had recent nose ulcers or nose surgery
- have had a nose injury that has not healed
- are using a steroid medicine for asthma, allergies or skin rash
- have an eye infection
- have or had glaucoma or cataracts

When using this product:

- the growth rate of some children may be slower
- some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief.
- do not share this bottle with anyone else as this may spread germs
- remember to tell your doctor about all the medicines you take, including this one

Dist. by: Chatter, Inc., a Sanofi Company, Chattanooga, TN 37409-0219 ©2017 Origin France

000000-01



PEEL HERE

Drug Facts (continued)

Stop use and ask a doctor if:

- you have, or come into contact with someone who has, chickenpox, measles or tuberculosis
- you have or develop symptoms of an infection such as a persistent fever
- you have any change in vision
- you have severe or frequent nosebleeds

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Read insert (inside package) on how to:

- use the spray
- clean the spray nozzle

ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER

- once daily, spray 2 times into each nostril while sniffing gently

CHILDREN 2 TO UNDER 12 YEARS OF AGE

- once your allergy symptoms improve, reduce to 1 spray in each nostril per day

Children 6 to under 12 years of age

- in an adult should supervise use
- once daily, spray 1 time into each nostril while sniffing gently

Drug Facts (continued)

CHILDREN 2 TO UNDER 12 YEARS OF AGE (continued)

children 6 to under 12 years of age

- if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.

children 2 to under 6 years of age

- an adult should supervise use
- once daily, spray 1 time into each nostril while sniffing gently

children under 2 years of age

- do not use

Do not use more than directed

- if you forget a dose, do not double the next dose
- do not spray into eyes or mouth
- if allergy symptoms do not improve after one week, stop using and talk to a doctor
- do not use for the common cold
- shake well before each use

Other information

- do not use if sealed package is torn or opened
- avoid package heat (heat). They contain important information.
- store between 20°-25°C (68°-77°F)

Inactive ingredients

benzalkonium chloride, carbonylmethylcellulose sodium, decahydrate disodium, hydrochloric acid or sodium hydroxide (for pH adjustment), microcrystalline cellulose, polyborate 80, purified water

Questions or comments? call toll-free 1-800-633-1610

Sanofi Regulatory Specifications		Verified Date: 11/13/17
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	14.00 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique / Helvetica Neue 57 Condensed	8.00 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8.00 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6.00 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6.00 point type
Bullets	Font name: Universal News with Commercial PI 97	6.00 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		Yes
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		Yes
Tracking:	0 to 16 Horizontal Scale:	94%-100%
Leading (Minimum space in body of Drug Facts):	6.3 pt. Maximum Characters/Inch (Characters plus spaces):	32
Barlines:	2.0 pt. Hairlines:	0.50 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		NA
PDP dimensions (in square inches)		NA
Font size of Statement of Identity (if not live text, to be measured in Helvetica capital "M")		NA
Font size of Logo/Largest Copy on PDP (if not live text, to be measured in Helvetica capital "M")		NA
Ratio of Statement of Identity to Logo/Largest Copy on PDP		NA
Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		NA

Nasacort Allergy 24 Hour Drug Facts Label

11.13.17

NAS_120+30sp_Multi-Pk_PeelAway_DFL_FNL2

Drug Facts

Active ingredient (in each spray) *Purpose*
Fluticasone propionate 50 mcg (1.0 mg/0.05 mL) Allergy symptoms reliever

Uses Temporarily relieves these symptoms of hay fever or other upper respiratory allergies: **nasal congestion** **runny nose** **sneezing** **itchy nose**

Warnings
Do not use
 • in children under 2 years of age
 • if you have ever had an allergic reaction to any of the ingredients

Ask a doctor before use if you:
 • have had recent nose ulcers or nose surgery
 • have had a nose injury that has not healed
 • are using a steroid medicine for asthma, allergies or skin rash
 • have an eye infection • have or had glaucoma or cataracts

When using this product:
 • the growth rate of some children may be slower
 • some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief.
 • do not share this bottle with anyone else as this may spread germs
 • remember to tell your doctor about all the medicines you take, including this one

Dist. by: Chatham, Inc., a Sanofi Company, Chatham, TN 37439-0219 ©2017 Origin France

000000-01

PEEL HERE

Drug Facts (continued)

Stop use and ask a doctor if:
 • you have, or come into contact with someone who has, chickenpox, measles or tuberculosis
 • you have or develop symptoms of an infection such as a persistent fever
 • you have any change in vision
 • you have severe or frequent nosebleeds

Directions
 Read insert (inside package) on how to use the spray.
 • **if pregnant or breast-feeding, ask a health professional before use.**
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Other information
 • **ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER**
 • once daily, spray 2 times into each nostril while sniffing gently
 • once your allergy symptoms improve, reduce to 1 spray in each nostril per day

CHILDREN 2 TO UNDER 12 YEARS OF AGE
 • the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.
 • once daily, spray 1 time into each nostril while sniffing gently

Drug Facts (continued)

CHILDREN 2 TO UNDER 12 YEARS OF AGE (continued)

children 6 to under 12 years of age (continued)
 • if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.

children 2 to under 6 years of age (continued)
 • an adult should supervise use
 • once daily, spray 1 time into each nostril while sniffing gently

children under 2 years of age (continued)
 • do not use
 • do not use more than directed
 • if you forget a dose, do not double the next dose
 • do not spray into eyes or mouth
 • if allergy symptoms do not improve after one week, stop using and talk to a doctor
 • do not use for the common cold
 • shake well before each use

Other information
 • do not use if sealed package is torn or opened
 • keep package and insert. They contain important information.
 • store between 20°-25°C (68°-77°F)

Inactive ingredients
 benzalkonium chloride, carbonylmethylcellulose sodium, dextrose, edetate disodium, hydrochloric acid or sodium hydroxide (for pH adjustment), microcrystalline cellulose, polyborate 80, purified water

Questions or comments? call toll-free 1-800-633-1610

Sanofi Regulatory Specifications		Verified Date: 11/13/17
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	14.00 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique / Helvetica Neue 57 Condensed	8.00 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8.00 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6.00 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6.00 point type
Bullets	Font name: Universal News with Commercial PI 97	6.00 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		Yes
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		Yes
Tracking:	0 to 16 Horizontal Scale:	94%-100%
Leading (Minimum space in body of Drug Facts):	6.3 pt. Maximum Characters/Inch (Characters plus spaces):	32
Barlines:	2.0 pt. Hairlines:	0.50 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		NA
PDP dimensions (in square inches)		NA
Font size of Statement of Identity (if not live text, to be measured in Helvetica capital "M")		NA
Font size of Logo/Largest Copy on PDP (if not live text, to be measured in Helvetica capital "M")		NA
Ratio of Statement of Identity to Logo/Largest Copy on PDP		NA
Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		NA

Nasacort Allergy 24 Hour Drug Facts Label

11.13.17

NAS_2x120sp_Multi-Pk_PeelAway DFL_FNL2

Drug Facts

Active ingredient (in each spray) *Fluticasone propionate* **Purpose** Allergy symptom reliever

Warnings

- Do not use
 - in children under 2 years of age
 - if you have ever had an allergic reaction to any of the ingredients
- Ask a doctor before use if you
 - have had recent nose ulcers or nose surgery
 - have had a nose injury that has not healed
 - are using a steroid medicine for asthma, allergies or skin rash
 - have an eye infection
 - have or had glaucoma or cataracts
- When using this product
 - the growth rate of some children may be slower
 - some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief.
 - do not share this bottle with anyone else as this may spread germs
 - remember to tell your doctor about all the medicines you take, including this one

Dist. by: Chatter, Inc., a Sanofi Company, Chattanooga, TN 37409-0219 ©2017 Origin France

00000-01

PEEL HERE

Drug Facts (continued)

Stop use and ask a doctor if

- you have, or come into contact with someone who has, chickenpox, measles or tuberculosis
- you have or develop symptoms of an infection such as a persistent fever
- you have any change in vision
- you have severe or frequent nosebleeds

Directions

Read insert (inside package) on how to use the spray

- if pregnant or breast-feeding, ask a health professional before use
- Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER

- once daily, spray 2 times into each nostril while sniffing gently

CHILDREN 2 TO UNDER 12 YEARS OF AGE

- once daily, spray 1 time into each nostril while sniffing gently

Drug Facts (continued)

CHILDREN 2 TO UNDER 12 YEARS OF AGE (continued)

children 6 to under 12 years of age

- if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.

children 2 to under 6 years of age

- an adult should supervise use
- once daily, spray 1 time into each nostril while sniffing gently

children under 2 years of age

- do not use
- do not use more than directed
- if you forget a dose, do not double the next dose
- do not spray into eyes or mouth
- if allergy symptoms do not improve after one week, stop using and talk to a doctor
- do not use for the common cold
- shake well before each use

Other information

- do not use if sealed package is torn or opened
- avoid package heat (heat). They contain important information.
- store between 20°-25°C (68°-77°F)

Inactive ingredients

benzalkonium chloride, carbonylmethylcellulose sodium, dextrose, edetate disodium, hydrochloric acid or sodium hydroxide (for pH adjustment), microcrystalline cellulose, polyborate 80, purified water

Questions or comments? call toll-free 1-800-633-1610

Sanofi Regulatory Specifications		Verified Date: 11/13/17
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	14.00 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique / Helvetica Neue 57 Condensed	8.00 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8.00 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6.00 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6.00 point type
Bullets	Font name: Universal News with Commercial PI 97	6.00 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		Yes
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		Yes
Tracking:	0 to 16 Horizontal Scale:	94%-100%
Leading (Minimum space in body of Drug Facts):	6.3 pt. Maximum Characters/Inch (Characters plus spaces):	32
Barlines:	2.0 pt. Hairlines:	0.50 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		NA
PDP dimensions (in square inches)		NA
Font size of Statement of Identity (if not live text, to be measured in Helvetica capital "M")		NA
Font size of Logo/Largest Copy on PDP (if not live text, to be measured in Helvetica capital "M")		NA
Ratio of Statement of Identity to Logo/Largest Copy on PDP		NA
Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		NA

Nasacort Allergy 24 Hour Drug Facts Label

11.13.17

NAS_2x120sp_Multi-PK-ALT_PeelAway DFL_FNL2

Drug Facts

Active ingredient (in each spray) Fluticasone propionate 50 mcg (1.0 mg/0.05 mL) **Purpose** Allergy symptom reliever

Uses Temporarily relieves these symptoms of hay fever or other upper respiratory allergies: **nasal congestion** **runny nose** **sneezing** **itchy nose**

Warnings

Do not use

- if you have ever had an allergic reaction to any of the ingredients

Ask a doctor before use if you:

- have had recent nose ulcers or nose surgery
- have had a nose injury that has not healed
- are using a steroid medicine for asthma, allergies or skin rash
- have an eye infection
- have or had glaucoma or cataracts

When using this product:

- the growth rate of some children may be slower
- some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief.
- do not share this bottle with anyone else as this may spread germs
- remember to tell your doctor about all the medicines you take, including this one

Dist. by: Chatter, Inc., a Sanofi Company, Chattanooga, TN 37409-0219 ©2017 Origin France

000000-01

PEEL HERE

Drug Facts (continued)

Stop use and ask a doctor if:

- you have, or come into contact with someone who has, chickenpox, measles or tuberculosis
- you have or develop symptoms of an infection such as a persistent fever
- you have any change in vision
- you have severe or frequent nosebleeds

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Read insert (inside package) on how to:

- use the spray
- clean the spray nozzle

ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER

- once daily, spray 2 times into each nostril while sniffing gently

CHILDREN 2 TO UNDER 12 YEARS OF AGE

- once your allergy symptoms improve, reduce to 1 spray in each nostril per day

Children 6 to under 12 years of age:

- an adult should supervise use
- once daily, spray 1 time into each nostril while sniffing gently

Drug Facts (continued)

CHILDREN 2 TO UNDER 12 YEARS OF AGE (continued)

children 6 to under 12 years of age (continued)

- if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.

children 2 to under 6 years of age (continued)

- an adult should supervise use
- once daily, spray 1 time into each nostril while sniffing gently

children under 2 years of age

- do not use

Do not use more than directed

- if you forget a dose, do not double the next dose
- do not spray into eyes or mouth
- if allergy symptoms do not improve after one week, stop using and talk to a doctor
- do not use for the common cold
- shake well before each use

Other information

- do not use if sealed package is torn or opened
- avoid package heat (heat). They contain important information.
- store between 20°-25°C (68°-77°F)

Inactive ingredients

benzalkonium chloride, carbonylmethylcellulose sodium, dextrose, edetate disodium, hydrochloric acid or sodium hydroxide (for pH adjustment), microcrystalline cellulose, polyborate 80, purified water

Questions or comments? call toll-free 1-800-633-1610

Sanofi Regulatory Specifications		Verified Date: 11/13/17
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	14.00 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique / Helvetica Neue 57 Condensed	8.00 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8.00 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6.00 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6.00 point type
Bullets	Font name: Universal News with Commercial PI 97	6.00 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		Yes
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		Yes
Tracking:	0 to 16 Horizontal Scale:	94%-100%
Leading (Minimum space in body of Drug Facts):	6.3 pt. Maximum Characters/Inch (Characters plus spaces):	32
Barlines:	2.0 pt. Hairlines:	0.50 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		NA
PDP dimensions (in square inches)		NA
Font size of Statement of Identity (if not live text, to be measured in Helvetica capital "M")		NA
Font size of Logo/Largest Copy on PDP (if not live text, to be measured in Helvetica capital "M")		NA
Ratio of Statement of Identity to Logo/Largest Copy on PDP		NA
Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		NA

Nasacort Allergy 24 Hour 1.75"x1" In Pack Coupon

12.14.17

NAS_1.75x1_In Pack Coupon_DSN1


Folded Composite



Opened Front of Booklet



Opened Back of Booklet



Sanofi Consumer Health Care Spec Box		Verified Date: 012/14/17
1.75" x 1" InPack Coupon		
Headline	Font name: Helvetica Neue 95 Black	27.00 point type
Sub Headline/Message	Font name: Myriad Pro Semi Bold	9.00 point type
Body Text	Font name: Helvetica Neue 57 Condensed	8.00 point type
Exp. Date	Font name: Helvetica Neue 57 Condensed	6.00 point type
Tracking:	0 to 16	Horizontal Scale: 95%-100%
Leading:	6 pt.	Maximum Characters/Inch: (Characters plus spaces) 36
Barrlines:	NA	Hairlines: NA

Nasacort Allergy 24 Hour 2.75"x1" In-Pack Coupon

01.10.18

NAS_2.75x1_In-Pack Coupon_FNL1



MANUFACTURER'S COUPON | EXPIRATION DATE: 00/00/00

Consumer: There is a limit of one coupon per purchase, and the coupon can't be combined with other offers. Good only in the U.S. and overseas commissaries on the purchase of the specified product/size. Void where prohibited or if altered, copied, transferred, auctioned, or sold. You must pay sales tax. Cash value: 1/100 of 1c. **Retailer:** We will reimburse you for the face value of this coupon plus 8¢ if redeemed in compliance with our Coupon Reimbursement Policy (available at address below). Invoices proving purchase of sufficient stock to cover coupons presented for redemption must be provided upon request. Failure to follow the policy may constitute fraud and may void all coupons submitted.

Mail to: Chattem, Inc. 1050,
NCH Marketing Services,
P.O. Box 880001,
El Paso, TX 88588-0001.
©2017 Chattem, Inc.

0041167-014501



000000-01