



VISINE®

ALLERGY EYE RELIEF
MULTI-ACTION



Drug Facts

Active ingredients	Purpose
Naphazoline HCl 0.025%	Redness reliever
Pheniramine maleate 0.3%	Antihistamine

Use
Temporarily relieves itchy, red eyes due to:
• pollen • ragweed • grass • animal hair and dander

Warnings
For external use only
Do not use if you are sensitive to any ingredient in this product.

Ask a doctor before use if you have
• heart disease • high blood pressure
• narrow angle glaucoma • trouble urinating

When using this product
• pupils may become enlarged temporarily causing light sensitivity
• do not touch tip of container to any surface to avoid contamination
• replace cap after each use
• remove contact lenses before using
• do not use if this solution changes color or becomes cloudy
• overuse may cause more eye redness
• some users may experience a brief tingling sensation

Stop use and ask a doctor if
• you feel eye pain
• changes in vision occur
• redness or irritation of the eye lasts
• condition worsens or lasts more than 72 hours

Keep out of reach of children.
If swallowed, get medical help or contact a Poison Control Center right away. Accidental swallowing by infants and children may lead to coma and marked reduction in body temperature.

Drug Facts (Continued)

Directions
• adults and children 6 years of age and over: put 1 or 2 drops in the affected eye(s) up to four times a day
• children under 6 years of age: consult a doctor

Other information
Store at 15° to 25°C (59° to 77°F)

Inactive ingredients
benzalkonium chloride, boric acid, edetate disodium, purified water, sodium borate, sodium hydroxide and/or hydrochloric acid

Questions?
call toll-free 888-734-7648 or 215-273-8755 (collect)

VISINE.

NAPHAZOLINE HCl 0.025%
AND PHENIRAMINE MALEATE 0.3%
OPHTHALMIC SOLUTION
ANTIHISTAMINE AND REDNESS RELIEVER

VISINE.

NAPHAZOLINE HCl 0.025%
AND PHENIRAMINE MALEATE 0.3%
OPHTHALMIC SOLUTION
ANTIHISTAMINE AND REDNESS RELIEVER

ALLERGY EYE RELIEF MULTI-ACTION



**ORIGINAL
PRESCRIPTION STRENGTH**

Clinically Proven to
Relieve Red,
Itchy Eyes



3 1254-749378 9

Caution: Do not use if VISINE® imprinted neckband on bottle is broken or missing.

STERILE EYE DROPS 1/2 FL. OZ. (15mL)

Distributed by: **ALCON & JOHNSON JOHNSON INC.**, Skillman, NJ 08588
Made in Belgium © 2014 2016 www.vision.com

USA
680644
955

LOT:

EXP:






VISINE® ALLERGY EYE RELIEF MULTI-ACTION

STERILE
EYE DROPS

VALUE TWIN PACK

VALUE TWIN PACK

VISINE.

VISINE.

NAPHAZOLINE HCl 0.025%
AND PHENIRAMINE MALEATE 0.3%
OPHTHALMIC SOLUTION
ANTIHISTAMINE AND REDNESS RELIEVER

NAPHAZOLINE HCl 0.025%
AND PHENIRAMINE MALEATE 0.3%
OPHTHALMIC SOLUTION
ANTIHISTAMINE AND REDNESS RELIEVER

ALLERGY EYE RELIEF MULTI-ACTION

ALLERGY EYE RELIEF MULTI-ACTION

ORIGINAL PRESCRIPTION STRENGTH

ORIGINAL PRESCRIPTION STRENGTH

Clinically Proven to
Relieve Red,
Itchy Eyes

Clinically Proven to
Relieve Red,
Itchy Eyes

STERILE
EYE DROPS

2 BOTTLES:
½ FL OZ (15 mL) EACH
NET 1 FL OZ (30 mL)

STERILE
EYE DROPS

2 BOTTLES:
½ FL OZ (15 mL) EACH
NET 1 FL OZ (30 mL)

Drug Facts (continued)

Directions

- adults and children 6 years of age and over: put 1 or 2 drops in the affected eye(s) up to four times a day
- children under 6 years of age: consult a doctor

Other information Store at 15° to 25°C (59° to 77°F)

Inactive ingredients benzalkonium chloride, boric acid, edetate disodium, purified water, sodium borate, sodium hydroxide and/or hydrochloric acid

Questions? call toll-free 888-734-7648 or 215-273-8755 (collect)

Drug Facts

Active ingredients	Purpose
Naphazoline HCl 0.025%	Redness reliever
Pheniramine maleate 0.3%	Antihistamine

Use
Temporarily relieves itchy, red eyes due to:
• pollen • ragweed • grass • animal hair and dander

Warnings
For external use only
Do not use if you are sensitive to any ingredient in this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- narrow angle glaucoma
- trouble urinating

When using this product

- pupils may become enlarged temporarily causing light sensitivity
- do not touch tip of container to any surface to avoid contamination
- replace cap after each use
- remove contact lenses before using
- do not use if this solution changes color or becomes cloudy
- overuse may cause more eye redness
- some users may experience a brief tingling sensation

Stop use and ask a doctor if

- you feel eye pain
- changes in vision occur
- redness or irritation of the eye lasts
- condition worsens or lasts more than 72 hours

Keep out of reach of children.
If swallowed, get medical help or contact a Poison Control Center right away. Accidental swallowing by infants and children may lead to coma and marked reduction in body temperature.

Caution: Do not use if VISINE® imprinted neckband on bottle is broken or missing.

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
Skillman, NJ 08558
Made in Belgium
© J&JCI 2018
www.visine.com
30041286

LOT:

EXP:



2D code is built with 0 BWR

(b) (4)

TO OPEN:
PUSH CAP
& TURN

VISINE®

NAPHAZOLINE HCl 0.025%
AND PHENIRAMINE MALEATE 0.3%
OPHTHALMIC SOLUTION
ANTIHISTAMINE AND REDNESS RELIEVER

**ALLERGY EYE RELIEF
MULTI-ACTION**

CLINICALLY PROVEN
TO RELIEVE RED,
ITCHY EYES

680642

STERILE N 3123417-000-78 9
EYE DROPS 1/2 FL. OZ. (15mL)

(b) (4)

Pull to Open

IMPORTANT: SEE CARTON FOR COMPLETE INFORMATION.

USE: Temporarily relieves itchy, red eyes due to pollen, ragweed, grass, animal hair and dander.

DIRECTIONS: Adults and children 6 years of age and over: Put 1 or 2 drops in the affected eye(s) up to four times a day. Some users may experience a brief stinging sensation.

ACTIVE INGREDIENTS: olopatadine HCl 0.025%, pheniramine maleate 0.3%.
INACTIVE INGREDIENTS: (See carton).
Store at 15° to 25° C (59° to 77° F).

Lot:

Exp:

680643

Keep this and all drugs out of the reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center right away.
CAUTION: Do not use if VISIBLE inverted redness on both is broken or missing.
Distributed by:
JOHNSON & JOHNSON
CONSUMER INC.
SILOAH, N.J. 08858
© ALLUCCI 2018
Made in Belgium
www.fish.com

WARNINGS:
(See carton for complete information):

If you experience eye pain, changes in vision, continued redness or irritation of the eye, or if the condition worsens or persists for more than 72 hours, discontinue use and consult a doctor. Accidental swallowing by infants and children may lead to coma AND MARKED REDUCTION IN BODY TEMPERATURE.
PARENTS NOTE: Before using with children under 6 years of age, consult a doctor.

(b) (4)

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

KAREN M MAHONEY
10/17/2018