

Important: Read all product information before using. Keep the box for important information. This product is intended for use in children.

Active ingredient (in each 5 mL) **Purpose**
 ibuprofen 100 mg (NSAID) Pain reliever/fever reducer

Uses: temporarily relieve minor aches and pains due to the common cold, flu, sore throat, headache, and toothache.

Warnings:
Allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** The chance is higher if your child has had stomach ulcers or bleeding problems, takes a blood thinning (anti-coagulant) or steroid drug, takes other drugs containing prescription nonsteroidal anti-inflammatory drugs (aspirin, ibuprofen, naproxen, or others), takes more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Soothe heat warning:** See the pediatrician if the child is under 6 months of age unless directed by doctor. **Do not use** if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer, or right before or after heart surgery. **Ask a doctor before use** if stomach bleeding was ever reported by your child, if the child has a history of stomach problems, such as heartburn, or if the child has

NDC 50580-184-50

Children's Motrin

contains **Ibuprofen**
Oral Suspension (NSAID)
100 mg per 5 mL 4 fl oz (120 mL)

Non-Staining Dye-Free • Original Berry Flavor

Directions: If the pain gets worse or lasts more than 3 days, or if redness or swelling is present in the painful area, or if any new symptoms appear. **Keep out of reach of children.** In case of an overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). **Directions:** This product does not contain directions or complete warnings for adult use. **Do not give more than directed.** Shake well before using. 5 mL = 1 teaspoon. Find right dose on chart below. If possible, use weight to dose; otherwise use age. Use only enclosed dosing cup. Do not use any other dosing device. If needed, repeat dose every 6-8 hours. Do not use more than 4 times a day. Replace original bottle cap to maintain child resistance.

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

*or as directed by a doctor

Other information: Each 5 mL contains: sodium 2 mg. Store between 20°C (68°F-77°F). Do not use if bottle wrap imprinted "SAFETY SEAL" is broken or missing. **Inactive ingredients:** sodium lauryl sulfate, anhydrous citric acid, flavor, glycerin, polyvinylpyrrolidone, polyethylene glycol, purified water, sodium benzoate, sucrose, xanthan gum.

Questions or comments? Call 1-877-895-3665 (toll-free) or 215-273-8755 (in PA).
 Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**
 McKel Consumer Healthcare Division
 Fort Washington, PA 19034 USA © J&JCI 2017 www.motrin.com
 30038104

Labeling Format Information: Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic

Drug Facts: NA pt	Body Text: 5.5 pt	Bullets: 4.36 pt	Horizontal Scale: 64%
Headers: NA pt	Leading: 6.00 pt	Barlines: NA pt	
Subheaders: 5.5 pt	Drug Facts (continued): NA pt	Hairlines: NA pt	Average Tracking: -25



Drug Facts (continued)

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver or kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
 - feels faint
 - vomits blood
 - has bloody or black stools
 - has stomach pain that does not get better
 - child has symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling
- child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.

Drug Facts (continued)

- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

* or as directed by a doctor

Other information

- each 5 mL contains: sodium 2 mg
- store between 20-25°C (68-77°F)
- do not use if carton is opened or if carton tape or bottle wrap imprinted "SAFETY SEAL" is broken or missing

Inactive ingredients acesulfame potassium, anhydrous citric acid, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum

Questions or comments?
Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
©J&JCI 2017
www.motrin.com 30038105

Children's Motrin
DYE-FREE
contains
Ibuprofen Oral Suspension (NSAID)
Non-Staining
Original Berry Flavor

See New Warnings
NDC 50580-184-50

For Ages 2 to 11 Years

Children's Motrin
contains **Ibuprofen Oral Suspension (NSAID)**
100 mg per 5 mL
Pain Reliever / Fever Reducer
DYE-FREE
Lasts up to 8 hours
Alcohol Free
Non-Staining
Original Berry Flavor
4 fl oz (120 mL)

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

Drug Facts

Active ingredient (in each 5 mL) Ibuprofen 100 mg (NSAID)*

Purpose Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses temporarily:
relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
hives, asthma (wheezing), skin reddening, facial swelling, shock, rash, blisters
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause stomach bleeding. The chance is higher if your child:
has had stomach ulcers or bleeding problems
takes a blood thinning (anticoagulant) or steroid drug
takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
takes more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use
if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
right before or after heart surgery

3 0045-0184-047

Labeling Format Information: Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic

Drug Facts: 8.00 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 90%
Headers: 6.00 pt	Leading: 5.50 pt	Barlines: 2.50 pt	
Subheaders: 5.00 pt	Drug Facts (continued): 8.00 pt	Hairlines: 0.50 pt	Average Tracking: -10

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

KAREN M MAHONEY
11/29/2018