



PAA105480.FDA01

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

**Do Not Use if breakable ring on bottle cap is separated or missing.**

**Children's  
Advil®  
Suspension**

**Ibuprofen Oral Suspension  
100 mg per 5 mL**

**Pain Reliever/Fever Reducer (NSAID)**

---

**Dye-Free / Alcohol-Free**

**Blue Raspberry-Flavored Liquid**

**4 FL OZ (120 mL)**

**Active ingredient (in each 5 mL)** ..... Ibuprofen 100 mg

**Uses** temporarily reduces fever, relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if the child has had stomach ulcers or bleeding problems, takes a blood thinning (anticoagulant) or steroid drug, takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), takes more or for a longer time than directed.

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult a doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by a doctor.

**Do not use if** the child has ever had an allergic reaction to any other fever reducer/pain reliever; right before or after heart surgery

**Ask a doctor before use** if stomach bleeding warning applies to the child; child has a history of stomach problems, such as heartburn; child has high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke; child has not been drinking fluids; child has lost a lot of fluid due to vomiting or diarrhea; child is taking a diuretic.

**Ask a doctor or pharmacist before use if the child is** under a doctor's care for any serious condition; taking any other drug

**When using this product** take with food or milk if upset stomach occurs

**Stop use and ask a doctor** if child experiences any of the following signs of stomach bleeding: feels faint; vomits blood; has bloody or black stools; or has stomach pain that does not get better; child has symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling; the child does not get any relief within first day (24 hours) of treatment; fever or pain gets worse or lasts more than 3 days; redness or swelling is present in the painful area; any new symptoms appear

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Each 5 mL contains: **sodium 10 mg.** Store at 20-25°C (68-77°F)

**Directions:** This product does not contain directions or complete warnings for adult use. Do not give more than directed. Shake well before using, mL = milliliter. Find right dose on chart. If possible, use weight to dose; otherwise use age. Repeat dose every 6-8 hours, if needed. Do not use more than 4 times a day. Measure only with the dosing cup provided. Dosing cup to be used with Children's Advil Suspension only. Do not use with other products. Dose lines account for product remaining in cup due to thickness of suspension.

Dosing Chart		
Weight (lb)	Age (yr)	Dose (mL)
under 24 lb	under 2 yr	ask a doctor
24-35 lb	2-3 yr	5 mL
36-47 lb	4-5 yr	7.5 mL
48-59 lb	6-8 yr	10 mL
60-71 lb	9-10 yr	12.5 mL
72-95 lb	11 yr	15 mL

**Questions or comments?** call 1-800-88-ADVIL  
Pizer, Madison, NJ 07940 USA © 2018 Pizer Inc.  
For most recent product information, visit [www.Advil.com](http://www.Advil.com)

PAA105481.FDA01

PAA105481.FDA01

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.6 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• WARNING BOX LINE, HAIRLINES	.5 pt