

**Do not use if neck wrap or hat liner seal is present. "Safety Seal" is broken or missing.** NDC 5240-110-XX  
**Children's Motrin**  
**Ibuprofen Chewable Tablets, 100 mg**  
**Pain Reliever/Fever Reducer (NSAID)**  
 Chew or crush tablets completely before swallowing.  
**24 Grape-Flavored Chewable Tablets**  
**Keep this box for important information.**  
 Active ingredient (in each tablet): Purpose  
 Ibuprofen 100 mg (NSAID) ... Pain reliever/fever reducer  
 Ibuprofen is a nonsteroidal anti-inflammatory drug (NSAID). It temporarily relieves pain and reduces fever. It relieves minor aches and pains due to the common cold, flu, sore throat, headaches, and toothaches. **Warnings: Allergy alert.**

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives; facial swelling; asthma (wheezing); shock; skin redness; rash; dizziness. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child: has had stomach ulcers or bleeding problems; takes a blood thinner (anticoagulant) or steroid drug; takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) or uses more or for a longer time than directed; **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor. **Do not use** if the child has ever had an allergic reaction to any other pain reliever/fever reducer; **right before or after heart surgery. Stop use and ask a doctor if** child experiences any of the following signs of stomach bleeding: **weak faint**; **vomits blood**; **has bloody or black stools**; **has stomach pain that does not get better**; **child has symptoms of heart problems or stroke**; **chest pain**; **trouble breathing**; **wheezes in one part or side of body**; **hoarse speech**; **leg swelling**; **the child does not get any relief within first day (24 hours) of treatment**; **stomach or pain gets worse or lasts more than 3 days**; **redness or swelling is present in the painful area**; **any new symptoms appear. Keep out of reach of children.**  
**Directions: Chew or crush tablets completely before swallowing. This product does not contain directions or complete warnings for adult use. Do not give more than directed and/or right before or after heart surgery. If possible, use weight to dose; otherwise use age. If needed, repeat dose every 6-8 hours. Do not use more than 4 times a day.**  
**Other information:** Phenylethylamine 2.8 mg per tablet; store between 20-25°C (68-77°F).

Weight (lb)	Age (yr)	Tablets
under 24	under 2	ask a doctor
24-35	2-3	1
36-47	4-5	1 1/2
48-59	6-8	2
60-71	9-10	2 1/2
72-95	11	3

048727-1-10  
 P. Washington, PA 19014 USA  
 EXP. DATE: 00000000  
 LOT: 00000000  
 00000000

LOT: EXP: 3 0045-060-244

NDC 50580-XXX-XX

# Children's Motrin®

**For Ages 2 to 11 Years**  
**Lasts up to 8 hours**

**Ibuprofen Chewable Tablets, 100 mg**  
**Pain Reliever / Fever Reducer (NSAID)**  
**Chewables**

Actual Size 


**24 Grape-Flavored Chewable Tablets**  
Chew or crush tablets completely before swallowing

**Drug Facts (continued)**

**Questions or comments?**  
Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Distributed by:  
**JOHNSON & JOHNSON CONSUMER INC.**  
McNeil Consumer Healthcare Division  
Fort Washington, PA 19034 USA  
©J&JCI 2018

[www.motrin.com](http://www.motrin.com)  
US\$3,320,855



**Drug Facts (continued)**

**Do not use**

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver disease, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

**Ask a doctor or pharmacist before use if the child is**


- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- do not use for throat burning that occurs, give with food or water
- do not take with food or milk. If stomach upset occurs

**Stop use and ask a doctor if**

- child experiences any of the following signs of stomach bleeding:
  - feels faint
  - vomits blood
  - has bloody or black stools
- has stomach pain that does not get better
- child has symptoms of heart problems or stroke
- trouble breathing
- swelled speech
- legs swelling
- the child does not feel any better within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear



**Important:** Read all product information before using. Keep this box or important information. This product is for children only.

**Drug Facts**

**Active ingredient (in each tablet)** Ibuprofen 100 mg (NSAID) ..... Pain reliever/fever reducer  
hydroxyzine antihistaminic drug

**Uses** temporarily:

- reduces fever
- relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

**Warnings**

**Allergy alert:** ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:


- hives
- facial swelling
- asthma (wheezing)
- swollen skin
- skin redness
- itch
- swelling
- stomach pain

**Stomach bleeding warning:** This product contains an NSAID, which may cause serious stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, naproxen, rofecoxib, or others)
- takes more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Sore throat warning:** Sore or swollen sore throat or sore throat accompanied by high fever, headache, nausea and vomiting may be serious. Consult doctor promptly. Do not use more than 7 days or administer to children under 3 years of age unless directed by doctor.



**Drug Facts (continued)**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-271-2122)

**Directions**

- chew or crush tablets completely before swallowing
- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- do not give more than directed
- if you give more than directed, if possible, use weight to dose, otherwise use age
- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day


**Dosing Chart**

Weight (lb)	Age (yr)	Tablets
under 24	under 2	ask a doctor
24-35	2-3	1
36-47	4-5	1 1/2
48-59	6-8	2
60-71	9-10	2 1/2
72-95	11	3

**Other information**

- phenylephrine contains phenylephrine 2.8 mg per tablet
- store between 20-25°C (68-77°F)
- do not use if neck wrap or foil inner seal is imprinted
- "Safety Seal" is broken or missing

**Inactive ingredients:** caseinamide potassium, ethyl alcohol, citric acid, aspartame, D-C-phenol, 7-calcium lactate, D-C-phenol, 30 aluminum lake, FD&C blue no. 1, aluminum lake, fumaric acid, hydroxyethyl cellulose, hypromellose, magnesium stearate, menthol, microcrystalline cellulose, povidone, sodium lauryl sulfate, sodium starch glycolate



-----  
**This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.**  
-----

/s/  
-----

KAREN M MAHONEY  
10/02/2018