This label may not be the latest approved by FDA. For current labeling information, please visit https://www.fda.gov/drugsatfda



EXP


X X X X X X X X X



For Complete Directions and Warnings
Do not use if foil seal under bottle cap printed "SAFETY SEAL®" is open or torn.

Drug Facts	
Active ingredients (in each chewable tablet)	Purposes
Famotidine 10 mg Calcium carbonate 800 mg Magnesium hydroxide 165 mg	Antacid
Use relieves heartburn associated wi indigestion and sour stomach	th acid

Distr buted by: OHNSON & JOHNSON CONSUMER INC. MoNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA @McNeil Consumer Pharmaceuticals Co. 20XX PEPCID® and PEPCID COMPLETE® are registered trademarks of McNeil CPC S Patent Nos. 5,989,588 and 6,814,978



Drug Facts (continued)

Other information

• each tablet contains: calcium 320 mg, magnesium 70 mg read the directions and warnings before use
 read the bottle. It contains important information. store at 20°-25°C (68°-77°F)

 protect from moisture
 do not use if foil seal under bottle cap printed "SAFETY SEAL®" is open or torn

Inactive ingredients cellulose acetate, corn starch, ospovidone, dextrose excipient, FD&C blue no. 1 aluminum lake, lavors, gum arabic, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, maltodextrin, mineral oil, sucralo yellow iron oxide

Questions or comments? call 1-800-755-4008 (toll-free) or 215-273-8755 (collect)

Tips for Managing Heartburn • Do not lie flat or bend over after eating • Do not wear tight-fitting clothing around the stomach • Do not eat before bedtime • Raise the head of your bed • Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables • Eat slowly and avoid big meals • If overweight, lose weight • Quit smoking



Famotidine 10 mg per tablet Calcium carbonate 800 mg per tablet

Magnesium hydroxide l65 mg per tablet Sho 25 Chewable Tablets Distributed by: JOHNSON & JOHNSON CONSUMER INC.


Distributed by OPD and OPD and Consoling the Consoling the Consoling that the Consoling the Consolin

Shown actual size

	Drug Facts (continued)	
er acid reducers	• nausea or vomiting • stomach pain	
th blood, or condition. See	Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.	
	Stop use and ask a doctor if •your heartburn continues or worsens •you need to take this product for more than 14 days	
re	If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)	
eating; dness	Directions • adults and children 12 years and over: • do not swallow tablet whole: chew completely • to relieve symptoms, chew 1 tablet before swallowing • do not use more than 2 chewable tablets in 24 hours • children under 12 years: ask a doctor	



This label may not be the latest approved by FDA. For current labeling information, please visit https://www.fda.gov/drugsatfda



EXP

X X X X X X X X X



NDC 16837-891-66 For Complete Directions and Warnings o not use if foil seal under bottle cap printed "SAFETY SEAL®" is open or torn.

Active ingredients (in each chewable tablet) Purposes Acid reducer . Famotidine 10 mg..... Calcium carbonate 800 mg.... Antacid Magnesium hydroxide 165 mg...Antacid

Use relieves heartburn associated with acid indigestion and sour stomach

motidine or other acid reducers if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools

These may be signs of a serious condition. See your doctor. with other acid reducers

SON & JOHNSON CONSUMER INC. Consumer Healthcare Division, Fort Washington, PA 19034 USA lef Consumer Pharmacouticals Co. 20XX. ID® and PEPCID COMPLETE® are registered trademarks of ICPC Lert Nos. 5,989,588 and 6,814,978

Drug Facts (continued)

• frequent chest pain frequent wheezing, particularly with heartburn
 unexplained weight loss nausea or vomiting stomach pain

prescription drugs.

Stop use and ask a doctor if vour heartburn continues or worsens

• you need to take this product for more than 14 days

Directions

 adults and children 12 years and over:
 do not swallow tablet whole: chew completely to relieve symptoms, chew 1 tablet before swallowing
do not use more than 2 chewable tablets in 24 hours children under 12 years: ask a doctor

Drug Facts (continued)

Other information

 Uther information

 • each tablet contains: calcium 320 mg, magnesium 70 mg

 • read the directions and warnings before use

 • read the bottle. It contains important information.

 • store at 20°-25°C (68°-77°F)

 • do not use if foil seal under bottle cap printed "SAFETY SEAL®" is open or torn

Inactive ingredients cellulose acetate, corn starch, crospovidone, dextrose excipient, FD&C blue no. 1 aluminum lake, flavo gum arabic, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, maltodextrin, mineral oil, sucralose vellow iron oxide

Questions or comments? call 1-800-755-4008 (toll-free) or 215-273-8755 (collect)



Famotidine 10 mg per tablet Calcium carbonate 800 mg per tablet Magnesium hydroxide 165 mg per tablet 50 Chewable Tablets



- **Tips for Managing Heartburn** Do not lie flat or bend over after eating Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
 Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals If overweight, lose weight
- Quit smoking
- SUCHREWIDDLE TRIDIETS Distributed by: JOHNSON & JOHNSON CONSUMER INC. McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA @McNeil Consumer Pharmaceuticals Co. 20XX PEPCDID® and PEPCDI COMPLETE® are registered trademarks of McNeil CPC US Patent Nos. 5,989,588 and 6,814,978

Ask a doctor before use if you have • had heartburn over 3 months. This may be a sign of a more serious condition. • heartburn with **lightheadedness, sweating, or dizziness** • chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)



Signature Page 1 of 1

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

KAREN M MAHONEY 12/21/2018