



2 FREE MIX-IN PAX Single Dose Packets SEE BACK

2 FREE MIX-IN PAX Single Dose Packets

MIRA LAX
Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

UNFLAVORED POWDER GRIT FREE

- Relieves Occasional Constipation/Irregularity
- Softens Stool

1 BOTTLE - NET WT 8.3 OZ (238 g)
2 PACKETS - NET WT 0.5 OZ (17 g) EACH
TOTAL NET WT 9.3 OZ (272 g)

16 ~~14~~ ONCE-DAILY DOSES



Drug Facts

Active ingredient (in each dose) Purpose
Polyethylene Glycol 3350, 17g (cap filled to line).....Osmotic Laxative

Use
relieves occasional constipation (irregularity)
generally produces a bowel movement in 1 to 3 days

Warnings
Allergy alert: Do not use if you are allergic to polyethylene glycol
Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have
nausea, vomiting or abdominal pain
a sudden change in bowel habits that lasts over 2 weeks
irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if
you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
you get diarrhea
you need to use a laxative for longer than 1 week

Drug Facts (continued)
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
do not take more than directed unless advised by your doctor
the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
adults and children 17 years of age and older:
fill to top of white section in cap which is marked to indicate the correct dose (17g)
stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
use once a day
use no more than 7 days
children 16 years of age or under: ask a doctor

Other information
store at 20° - 25°C (68° - 77°F)
tamper-evident: do not use if foil seal under cap, printed with "SEALED FOR YOUR PROTECTION" is missing, open or broken

Inactive ingredients none

Questions or comments?
1-800-MiraLAX (1-800-647-2529)

LOT/EXP AREA

Actual Size 100%



Bayer the Bayer Cross, and MiraLAX are registered trademarks of Bayer.
© 2018 Bayer,
Dist by: Bayer HealthCare LLC,
Whippany, NJ 07981
00000-00
Bayer

DRUG FACTS INFO		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt.**	8 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings*	8 pt.-bold italic ¹	8 pt.
Subheadings	≥6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	3.5 pt.
Leading	≥0.5 pt.	0.5 pt.
Hairlines	0.5 pt. rule	5 pt.
Box and Barlines†	>0.5 pt. rule	1 pt.
Characters Per Inch:	No more than 39	> 39

* Headings are to be at least 200% larger than Body Text.
** Drug Facts Title to be larger than all other type size used in DFP.
† FDA Recommends 2.5pt. rule





Drug Facts

Active ingredient (in each dose) **Purpose**
Polyethylene Glycol 3350,
17g (cap filled to line) Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed unless advised by your doctor**
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
 - fill to top of white section in cap which is marked to indicate the correct dose (17g)
 - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
 - use once a day
 - use no more than 7 days
- children 16 years of age or under: ask a doctor

Other information

- store at 20° - 25°C (68° - 77°F)
- tamper-evident: do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken

Inactive ingredients none

Questions or comments?
1-800-MiraLAX (1-800-647-2529)

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

KAREN M MAHONEY
05/04/2018