

<p>1A</p> <p>VAZALORE Aspirin Capsules, 325mg (NSAID)</p> <p>EXPIRES: MM/DD/YYYY LOT#: 012345</p> <p>Plx Pharma Inc. 8285 El Rio, Suite 130 Houston, TX 77054</p> <p>TO OPEN: Peel back paper from arrow</p>	<p>1B</p> <p>VAZALORE Aspirin Capsules, 325mg (NSAID)</p> <p>EXPIRES: MM/DD/YYYY LOT#: 012345</p> <p>Plx Pharma Inc. 8285 El Rio, Suite 130 Houston, TX 77054</p> <p>TO OPEN: Peel back paper from arrow</p>
<p>1C</p> <p>VAZALORE Aspirin Capsules, 325mg (NSAID)</p> <p>EXPIRES: MM/DD/YYYY LOT#: 012345</p> <p>Plx Pharma Inc. 8285 El Rio, Suite 130 Houston, TX 77054</p> <p>TO OPEN: Peel back paper from arrow</p>	<p>1D</p> <p>VAZALORE Aspirin Capsules, 325mg (NSAID)</p> <p>EXPIRES: MM/DD/YYYY LOT#: 012345</p> <p>Plx Pharma Inc. 8285 El Rio, Suite 130 Houston, TX 77054</p> <p>TO OPEN: Peel back paper from arrow</p>
<p>1E</p> <p>VAZALORE Aspirin Capsules, 325mg (NSAID)</p> <p>EXPIRES: MM/DD/YYYY LOT#: 012345</p> <p>Plx Pharma Inc. 8285 El Rio, Suite 130 Houston, TX 77054</p> <p>TO OPEN: Peel back paper from arrow</p>	<p>Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinner (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed</p>
<p>1F</p> <p>VAZALORE Aspirin Capsules, 325mg (NSAID)</p> <p>EXPIRES: MM/DD/YYYY LOT#: 012345</p> <p>Plx Pharma Inc. 8285 El Rio, Suite 130 Houston, TX 77054</p> <p>TO OPEN: Peel back paper from arrow</p>	<p>1G</p> <p>VAZALORE Aspirin Capsules, 325mg (NSAID)</p> <p>EXPIRES: MM/DD/YYYY LOT#: 012345</p> <p>Plx Pharma Inc. 8285 El Rio, Suite 130 Houston, TX 77054</p> <p>TO OPEN: Peel back paper from arrow</p>

VAZALORE™

Aspirin Capsules, 325 mg
Pain Reliever/Fever Reducer (NSAID)

28 Liquid-Filled Capsules

*Consult your healthcare provider before using this product for your heart.




NDC 00000-0000-0

VAZALORE™

Aspirin Capsules, 325 mg
Pain Reliever/Fever Reducer (NSAID)

325 mg

FIRST LIQUID-FILLED ASPIRIN CAPSULES



*Consult your healthcare provider before using this product for your heart.

28 Liquid-Filled Capsules

TAMPER EVIDENT: DO NOT USE IF MPR NTD SAFETY SEAL WITH REPEATED TEXT "SEALED FOR YOUR PROTECTION" UNDER CAP. 5 M'SS NG OR BROKEN.

VAZALORE™

Aspirin Capsules, 325 mg
Pain Reliever/Fever Reducer (NSAID)



Distributed by:
PLX Pharma Inc.
8285 El Rio, Suite 130
Houston, TX 77054

Visit us at: www.company.xxx.com



Area for
Lot & exp.

RETAIN CARTON FOR COMPLETE WARNINGS AND INFORMATION

*Consult your healthcare provider before using this product for your heart.

Drug Facts (continued)

Inactive ingredients
aminofrutic acid, carboxystyrene, colloidal silicon dioxide, FD&C Blue #1, hypromellose, medium chain triglycerides, pharmaceutical ink, potassium chloride, soy lecithin, titanium dioxide

Questions or comments

Call 1 888 842 3567 8:30 AM - 4:00 PM ET, Monday - Friday



FOR POSITION ONLY

FOR POSITION ONLY

FOR POSITION ONLY

Drug Facts

Active ingredient (in each capsule)

Aspirin (NSAID), 325mg
nonsteroidal anti-inflammatory drug
pain reliever/fever reducer

Purpose

You have high blood pressure, heart disease, liver cirrhosis, or kidney disease
You are taking a diuretic
You have asthma

Uses

For the temporary relief of minor aches and pains associated with:
• cold • headache • backache • muscular aches • toothache
• premenstrual & menstrual cramps • minor pain of arthritis
• temporarily reduces fever

Warnings

Key symptoms: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:
• hives • facial swelling • shock • asthma (wheezing)
This product contains soy.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

• are age 60 or older • have had stomach ulcers or bleeding problems

• take a blood thinning (anticoagulant) or steroid drug

• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

• have 5 or more alcoholic drinks every day while using this product

• take more or for a longer time than directed

Do not use if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer.

Ask doctor before use if

• stomach bleeding warning applies to you

• you have a history of stomach problems, such as heartburn

Drug Facts (continued)

Other information

• do not use if the blue band around the capsule is missing or broken
• avoid excessive heat above 40°C (104°F)
• store at 15°-30°C (59°-86°F)
• save carton for full directions and warnings

Directions
• drink a full glass of water with each dose
• adults and children 12 years and over:
• take 1 or 2 capsules every 4 hours or 3 capsules every 6 hours while symptoms persist
• do not exceed 12 capsules in 24 hours
• children under 12 years: consult a doctor

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

If pregnant or breast feeding, ask a health care professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Stop use and ask a doctor if

- you are experiencing any of the following signs of stomach bleeding:
• vomit blood • have bloody or black stools
- feel faint
- have stomach pain that doesn't get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts for more than 3 days
- ringing in the ears or loss of hearing occurs
- redness or swelling is present in the painful area

any new symptoms occur. These could be signs of a serious condition.

Ask doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout or arthritis
- you have asthma
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

NDC 1234-5678-90

VAZALORE™
Aspirin Capsules, 325 mg
Pain Reliever/Fever Reducer (NSAID)

*Consult your healthcare provider before using this product for your heart.

30 Liquid-filled Capsules

TAMPA EDITION: DO NOT USE IF PAIN MEDICATION SEALS WITH REPEATED TEXT
*EXCLUDED FROM PROTECTION UNDER CAP 335 SIGNATURE BANK

Drug Facts

Active ingredient (in each capsule)	Purpose
Aspirin (NSAID) ¹ 325mg, pain reliever/fever reducer ¹ nonsteroidal anti-inflammatory drug	

Uses

- For the temporary relief of minor aches and pains associated with:
 - a cold • headache • backache • muscular aches
 - toothache • premenstrual & menstrual cramps
 - minor pain of arthritis
- Temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:
• hives • facial swelling • shock • asthma (wheezing)
This product contains soy.

⬇ Continue Reading

Distributed By: Plx Pharma Inc.
8285 El Rio, Suite 130 Houston, TX 77054

PHE

LOT/EXP

Drug Facts (continued)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 5 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

If pregnant or breast-feeding, ask a health care professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.


Directions

- drink a full glass of water with each dose
- adults and children 12 years and over:
 - take 1 or 2 capsules every 4 hours or 3 capsules every 6 hours while symptoms persist
 - do not exceed 12 capsules in 24 hours
- children under 12 years: consult a doctor

Other information • save carton for full directions and warnings • store at 15-30°C (59-86°F) • avoid excessive heat above 40°C (104°F) • do not use if the blue band around the capsule is missing or broken

Inactive ingredients anhydrous citric acid, carrageenan, colloidal silicon dioxide, FD&C Blue #1, hypromellose, medium chain triglycerides, pharmaceutical ink, potassium chloride, soy lecithin, titanium dioxide

Questions or comments
Call 1-888-842-5567 8:30 AM - 4:00 PM ET, Monday - Friday



No Print
Glue Area

Drug Facts (continued)

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer.

Ask doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout or arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you are experiencing any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that doesn't get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts for more than 3 days
- ringing in the ears or loss of hearing occurs
- redness or swelling is present in the painful area
- any new symptoms occur. These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health care professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over:
 - take 1 or 2 capsules every 4 hours or 3 capsules every 6 hours while symptoms persist
 - do not exceed 12 capsules in 24 hours
- children under 12 years: consult a doctor

Other information

- save carton for full directions and warnings
- store at 15-30°C (59-86°F)
- avoid excessive heat above 40°C (104°F)
- do not use if the blue band around the capsule is missing or broken

Inactive ingredients

anhydrous citric acid, carrageenan, colloidal silicon dioxide, FD&C Blue #1, hypromellose, medium chain triglycerides, pharmaceutical ink, potassium chloride, soy lecithin, titanium dioxide

Questions or comments

Call 1-888-842-3567 8:30 AM - 4:00 PM ET, Monday - Friday

30 Liquid-filled Capsules
 Consult your healthcare provider before using this product for your heart.
 Pain Reliever/Fever Reducer (NSAID)
VAZALORE™
 Aspirin Capsules, 325 mg
 TAMPER EVIDENT- DO NOT USE IF SAFETY SEAL WITH REPEATED TEXT "SEALED FOR YOUR PROTECTION" UNDER CAP IS MISSING OR BROKEN.

30 Liquid-filled Capsules
325 mg FIRST LIQUID-FILLED ASPIRIN CAPSULES
 Aspirin Capsules, 325 mg
 Pain Reliever/Fever Reducer (NSAID)
VAZALORE™
 NDC 000000-0000-0
 Consult your healthcare provider before using this product for your heart.

RETAIN CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts

Active ingredient (in each capsule)	Purpose
Aspirin (NSAID)* 325mg pain reliever/fever reducer	
*nonsteroidal anti-inflammatory drug	

Uses

- For the temporary relief of minor aches and pains associated with:
 - a cold
 - headache
 - backache
 - muscular aches
 - toothache
 - premenstrual & menstrual cramps
 - minor pain of arthritis
- Temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
 - facial swelling
 - shock
 - asthma (wheezing)
- This product contains soy.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older



No Print
Glue Area

Distributed by: **PLx Pharma Inc.**
 8285 El Rio, Suite 130
 Houston, TX 77054
 Visit us at: www.companyxxx.com



LOT/EXP

NDC - 1234-5678-90

WAZALORE™
Aspirin Capsules, 325 mg
Pain Reliever/Fever Reducer (NSAID)

*Consult your healthcare provider before using this product for your heart.

120 Liquid-Filled Capsules

Drug Facts

Active ingredient (in each capsule) **Purpose**

Aspirin (NSAID)* 325 mg pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- For the temporary relief of minor aches and pains associated with:
 - a cold • headache • backache • muscular aches • toothache
 - premenstrual & menstrual cramps • minor pain of arthritis
- Temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert:

Aspirin may cause a severe allergic reaction which may include:
• hives • facial swelling • shock • asthma (wheezing)
This product contains soy.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
• are age 60 or older • have had stomach ulcers or bleeding problems
• Take a blood thinning (anticoagulant) or steroid drug
• Take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
• have 3 or more alcoholic drinks every day while using this product
• Take more or for a longer time than directed

If pregnant or breast feeding, ask a health care professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over:
 - Take 1 or 2 capsules every 4 hours or 3 capsules every 6 hours while symptoms persist
 - do not exceed 12 capsules in 24 hours
- children under 12 years: consult a doctor

Other information

- save carton for full directions and warnings
- store at 15-30°C (59-86°F)
- avoid excessive heat above 40°C (104°F)
- do not use if the blue band around the capsule is missing or broken

Inactive ingredients anhydrous citric acid, carrageenan, colloidal silicon dioxide, FD&C Blue #1, hypromellose, medium chain triglycerides, pharmaceutical ink, potassium chloride, soy lecithin, titanium dioxide

Questions or comments

Call 1 888 842 3567 8:30 AM - 4:00 PM ET, Monday - Friday
Distributed By: PLx Pharma Inc., 8285 El Rio, Suite 130, Houston, TX 77054

TAMPEN EVIDENCE: DO NOT USE IF IMPRINTED SAFETY SEAL WITH REPEATED TEXT "SEALED FOR YOUR PROTECTION" UNDER CAP IS MISSING OR BROKEN.

LOT/EXP

120 Liquid-Filled Capsules

*Consult your healthcare provider before using this product for your heart.

WAZALORE™
Aspirin Capsules, 325 mg
Pain Reliever/Fever Reducer (NSAID)

NDC 00000-0000-0

WAZALORE™

Aspirin Capsules, 325 mg
Pain Reliever/Fever Reducer (NSAID)

325mg

FIRST LIQUID-FILLED ASPIRIN CAPSULES

120 Liquid-Filled Capsules

*Consult your healthcare provider before using this product for your heart.

Distributed by:
PLx Pharma Inc.
8285 El Rio, Suite 130
Houston, TX 77054
Visit us at: www.compannyxxx.com

LOT/EXP

0 1000001000000

Drug Facts (continued)

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer.

Ask doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout or arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you are experiencing any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that doesn't get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts for more than 3 days
- ringing in the ears or loss of hearing occurs
- redness or swelling is present in the painful area
- any new symptoms occur. These could be signs of a serious condition.

Drug Facts (continued)

if pregnant or breast-feeding, ask a health care professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over:
 - take 1 or 2 capsules every 4 hours or 3 capsules every 6 hours while symptoms persist
 - do not exceed 12 capsules in 24 hours
- children under 12 years: consult a doctor

Other information

- save carton for full directions and warnings
- store at 15-30°C (59-86°F)
- avoid excessive heat above 40°C (104°F)
- do not use if the blue band around the capsule is missing or broken

Inactive ingredients

anhydrous citric acid, carraageenan, colloidal silicon dioxide, FD&C Blue #1, hypromellose, medium chain triglycerides, pharmaceutical ink, potassium chloride, soy lecithin, titanium dioxide

Questions or comments
Call 1-888-842-3567 8:30 AM – 4:00 PM ET, Monday – Friday

Drug Facts

Active Ingredient (in each capsule)
Aspirin (NSAID)* 325mg pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Purpose

For the temporary relief of minor aches and pains associated with:

- a cold
- headache
- backache
- muscular aches
- toothache
- premenstrual & menstrual cramps
- minor pain of arthritis
- Temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

This product contains soy.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older

REMAIN CARTON FOR COMPLETE WARNINGS AND INFORMATION

120 Liquid-Filled Capsules

*Consult your healthcare provider before using this product for your heart.

WAZALORE™
Aspirin Capsules, 325 mg
Pain Reliever/Fever Reducer (NSAID)

TAMPER EVIDENT: DO NOT USE IF SAFETY SEAL WITH REPEATED TEXT "SEALED FOR YOUR PROTECTION" UNDER CAP IS MISSING OR BROKEN.

NDC 00000-0000-0

WAZALORE™

Aspirin Capsules, 325 mg
Pain Reliever/Fever Reducer (NSAID)

325mg

FIRST LIQUID-FILLED ASPIRIN CAPSULES

120 Liquid-Filled Capsules

*Consult your healthcare provider before using this product for your heart.

Distributed by:
PLx Pharma Inc.
8285 El Rio, Suite 130
Houston, TX 77054
Visit us at: www.compannyxxx.com

LOT/EXP

0 1000001000000

Drug Facts (continued)

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer.

Ask doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout or arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you are experiencing any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that doesn't get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts for more than 3 days
- ringing in the ears or loss of hearing occurs
- redness or swelling is present in the painful area
- any new symptoms occur. These could be signs of a serious condition.

Drug Facts (continued)

if pregnant or breast-feeding, ask a health care professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over:
 - take 1 or 2 capsules every 4 hours or 3 capsules every 6 hours while symptoms persist
 - do not exceed 12 capsules in 24 hours
- children under 12 years: consult a doctor

Other information

- save carton for full directions and warnings
- store at 15-30°C (59-86°F)
- avoid excessive heat above 40°C (104°F)
- do not use if the blue band around the capsule is missing or broken

Inactive ingredients

anhydrous citric acid, carraageenan, colloidal silicon dioxide, FD&C Blue #1, hypromellose, medium chain triglycerides, pharmaceutical ink, potassium chloride, soy lecithin, titanium dioxide

Questions or comments
Call 1-888-842-3567 8:30 AM – 4:00 PM ET, Monday – Friday

Drug Facts

Active Ingredient (in each capsule)
Aspirin (NSAID)* 325mg pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Purpose

For the temporary relief of minor aches and pains associated with:

- a cold
- headache
- backache
- muscular aches
- toothache
- premenstrual & menstrual cramps
- minor pain of arthritis
- Temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

This product contains soy.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

THERESA M MICHELE
07/25/2018