

6.5 pt Helvetica Black Cond.

5 pt Helvetica Bold Cond.

6 pt Helvetica Bold Cond.

6 pt Helvetica Medium

Place tablet on tongue; tablet disintegrates, with or without water.
Do not crush or chew tablets.
First two doses of a 14-day course of treatment.
Take 1 tablet every day for 14 days.
Do not take for more than 14 days or more often than every 4 months unless directed by a doctor.

Omeprazole
Delayed Release Orally
Disintegrating Tablets 20 mg
acid reducer
Push tablet through foil
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam
2069200, Israel

Omeprazole
Delayed Release Orally
Disintegrating Tablets 20 mg
acid reducer
Push tablet through foil
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam
2069200, Israel

Reference ID: 4227553

6.5 pt Helvetica Black Cond.
5 pt Helvetica Bold Cond.
6 pt Helvetica Bold Cond.
6 pt Helvetica Medium

<p>Place tablet on tongue; tablet disintegrates, with or without water. Do not crush or chew tablets. One 14-day course of treatment. Take 1 tablet every day for 14 days. Do not take for more than 14 days or more often than every 4 months unless directed by a doctor.</p>	
<p>Omeprazole Delayed Release Orally Disintegrating Tablets 20 mg acid reducer Push tablet through foil Dexcel Pharma Technologies Ltd 10 Hakidma Street, Yokneam 2069200, Israel</p>	<p>Omeprazole Delayed Release Orally Disintegrating Tablets 20 mg acid reducer Push tablet through foil Dexcel Pharma Technologies Ltd 10 Hakidma Street, Yokneam 2069200, Israel</p>
<p>Omeprazole Delayed Release Orally Disintegrating Tablets 20 mg acid reducer Push tablet through foil Dexcel Pharma Technologies Ltd 10 Hakidma Street, Yokneam 2069200, Israel</p>	<p>Omeprazole Delayed Release Orally Disintegrating Tablets 20 mg acid reducer Push tablet through foil Dexcel Pharma Technologies Ltd 10 Hakidma Street, Yokneam 2069200, Israel</p>
<p>Omeprazole Delayed Release Orally Disintegrating Tablets 20 mg acid reducer Push tablet through foil Dexcel Pharma Technologies Ltd 10 Hakidma Street, Yokneam 2069200, Israel</p>	<p>Omeprazole Delayed Release Orally Disintegrating Tablets 20 mg acid reducer Push tablet through foil Dexcel Pharma Technologies Ltd 10 Hakidma Street, Yokneam 2069200, Israel</p>

<p>Omeprazole Delayed Release Orally Disintegrating Tablets 20 mg acid reducer</p> <p>Push tablet through foil</p> <p>Dexcel Pharma Technologies Ltd 10 Hakidma Street, Yokneam 2069200, Israel</p>	<p>Omeprazole Delayed Release Orally Disintegrating Tablets 20 mg acid reducer</p> <p>Push tablet through foil</p> <p>Dexcel Pharma Technologies Ltd 10 Hakidma Street, Yokneam 2069200, Israel</p>	<p>Omeprazole Delayed Release Orally Disintegrating Tablets 20 mg acid reducer</p> <p>Push tablet through foil</p> <p>Dexcel Pharma Technologies Ltd 10 Hakidma Street, Yokneam 2069200, Israel</p>
<p>Omeprazole Delayed Release Orally Disintegrating Tablets 20 mg acid reducer</p> <p>Push tablet through foil</p> <p>Dexcel Pharma Technologies Ltd 10 Hakidma Street, Yokneam 2069200, Israel</p>	<p>Omeprazole Delayed Release Orally Disintegrating Tablets 20 mg acid reducer</p> <p>Push tablet through foil</p> <p>Dexcel Pharma Technologies Ltd 10 Hakidma Street, Yokneam 2069200, Israel</p>	<p>Omeprazole Delayed Release Orally Disintegrating Tablets 20 mg acid reducer</p> <p>Push tablet through foil</p> <p>Dexcel Pharma Technologies Ltd 10 Hakidma Street, Yokneam 2069200, Israel</p>
<p>Omeprazole Delayed Release Orally Disintegrating Tablets 20 mg acid reducer</p> <p>Push tablet through foil</p> <p>Dexcel Pharma Technologies Ltd 10 Hakidma Street, Yokneam 2069200, Israel</p>	<p>Place tablet on tongue; tablet disintegrates, with or without water. Do not crush or chew tablets. One 14-day course of treatment. Do not take for more than 14 days or more often than every 4 months unless directed by a doctor.</p>	<p>Omeprazole Delayed Release Orally Disintegrating Tablets 20 mg acid reducer</p> <p>Push tablet through foil</p> <p>Dexcel Pharma Technologies Ltd 10 Hakidma Street, Yokneam 2069200, Israel</p>
<p>Omeprazole Delayed Release Orally Disintegrating Tablets 20 mg acid reducer</p> <p>Push tablet through foil</p> <p>Dexcel Pharma Technologies Ltd 10 Hakidma Street, Yokneam 2069200, Israel</p>	<p>Omeprazole Delayed Release Orally Disintegrating Tablets 20 mg acid reducer</p> <p>Push tablet through foil</p> <p>Dexcel Pharma Technologies Ltd 10 Hakidma Street, Yokneam 2069200, Israel</p>	<p>Omeprazole Delayed Release Orally Disintegrating Tablets 20 mg acid reducer</p> <p>Push tablet through foil</p> <p>Dexcel Pharma Technologies Ltd 10 Hakidma Street, Yokneam 2069200, Israel</p>
<p>Omeprazole Delayed Release Orally Disintegrating Tablets 20 mg acid reducer</p> <p>Push tablet through foil</p> <p>Dexcel Pharma Technologies Ltd 10 Hakidma Street, Yokneam 2069200, Israel</p>	<p>Omeprazole Delayed Release Orally Disintegrating Tablets 20 mg acid reducer</p> <p>Push tablet through foil</p> <p>Dexcel Pharma Technologies Ltd 10 Hakidma Street, Yokneam 2069200, Israel</p>	<p>Omeprazole Delayed Release Orally Disintegrating Tablets 20 mg acid reducer</p> <p>Push tablet through foil</p> <p>Dexcel Pharma Technologies Ltd 10 Hakidma Street, Yokneam 2069200, Israel</p>

5.5 pt Helvetica Black Cond. —
4.5 pt Helvetica Bold Cond. —
5 pt Helvetica Bold Cond. —
5 pt Helvetica Medium —

[illegible]

CODE AREA



Melts In Your Mouth

Actual size



- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight ■ Quit smoking

8-31-17

8-31-17

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue 57 Condensed
HEADINGS: 8 pt Helvetica Neue 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue 77 Bold Condensed
TEXT: 6 pt Helvetica Neue 57 Condensed
LEADING: 6.5 and 7.0 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt

8-31-17

9-13-17

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue 57 Condensed
HEADINGS: 8 pt Helvetica Neue 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue 77 Bold Condensed
TEXT: 6 pt Helvetica Neue 57 Condensed
LEADING: 6.5 and 7.0 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt

9-13-17

**Location for
→ lot number and
expiration date**

[illegible]

CODE AREA

1Y955 FA C2-K

TYPE SIZES:
TITLE: 9 pt Helvetica Neue 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue 77 Bold Condensed Oblique
 (continued) 8 pt Helvetica Neue 57 Condensed
HEADINGS: 8 pt Helvetica Neue 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue 77 Bold Condensed
TEXT: 6 pt Helvetica Neue 57 Condensed
LEADING: 6.5 and 6.7 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt

**Omeprazole Delayed Release Orally
Disintegrating Tablets 20 mg
14 Tablets**

Tips for Managing Heartburn

- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight ■ Quit smoking

9-13-17

STANDARD FORMAT

TYPE SIZES:

TITLE: 9 pt Helvetica Neue 77 Bold Condensed Oblique

TITLE (continued): 8 pt Helvetica Neue 77 Bold Condensed Oblique (continued)

8 pt Helvetica Neue 57 Condensed

HEADINGS: 8 pt Helvetica Neue 77 Bold Condensed Oblique

SUBHEADINGS: 6 pt Helvetica Neue 77 Bold Condensed

TEXT: 6 pt Helvetica Neue 57 Condensed

LEADING: 6.5 and 7.0 pt

BULLETS: 5 pt Square

TELEPHONE #: 6 pt Helvetica Neue 77 Bold Condensed

HEAVY LINES: 1.5 pt

HAIRLINES: 0.5 pt

9-13-17

The diagram shows a printed tablet blister pack with two compartments. The left compartment contains a printed tablet with a barcode and the number '0'. The right compartment is empty. Below the blister pack, there are instructions for managing heartburn, a safety feature warning, and a crosshair symbol.

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Safety Feature –
Do not use if printed tablet blister unit is open or torn.

8-31-17

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue 57 Condensed
HEADINGS: 8 pt Helvetica Neue 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue 77 Bold Condensed
TEXT: 6 pt Helvetica Neue 57 Condensed
LEADING: 6.5 and 7.0 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt

9-13-17

The diagram illustrates the layout of a product box with several key features and instructions:

- Barcode:** A barcode is located on the front of the box, with the number 0 00000 00000 0 printed below it.
- Tips for Managing Heartburn:** A list of instructions is provided on the front of the box:
 - Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking
- Safety Feature:** A warning is provided on the side of the box: "Do not use if printed tablet blister unit is open or torn."
- Registration Mark:** A registration mark (a circle with a crosshair) is located on the bottom of the box.

[illegible]

CODE AREA

1Y955 FA C2-A

Omeprazole

Delayed Release
Orally Disintegrating Tablets 20 mg

[illegible]

Use

- Treats frequent heartburn (occurs *2 or more* days a week)
- Not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Active ingredient (in each tablet)	Purpose
Omeprazole 20 mg.	Acid reducer

Directions

- for adult: 18 years or aged and older
- the product to be used once a day for 14 days
- in may take 10-14 days to feel effect; some people get complete relief

14-Day Course of Treatment

- take 1 tablet before eating in the morning
- do not crush or chew tablets
- place the tablet on tongue; let it disintegrate, with or without water.
- The tablet can also be swallowed whole with water.
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days
- do not take this medicine while you are still on it

Repeated (1-4 day Course if needed)

- you may repeat at 4-day course every 4 months
- do not take more than 1 and 1 day or more often in a day
- if you have this need checked by a doctor
- children under 18 years of age: ask a doctor before use. Hardium in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton, to contain important information.
- store at 20-25°C (68-77°F); keep product out of high heat and moisture

Drug Facts (continued)

- Stop use and ask a doctor if:
 - you have had a rash or joint pain
 - you need to take more than 1 course of treatment every 4 months
 - you need to take more than 14 days
 - you need to take more than 1 course of treatment every 4 months
 - you develop a rash or joint pain
- If you get an ear or nose -feeling, ask a health professional before use.
- Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222).

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Safety Feature –
Do not use if printed tablet
blister unit is open or torn.



NDC 64239-XXX-03

NEW DOSAGE FORM!

MELTech™
Melts In Your Mouth

FDA Approved

Omeprazole

Delayed Release
Orally Disintegrating Tablets 20 mg
Acid Reducer

Treats Frequent Heartburn!
Occurring 2 or More Days a Week

42 TABLETS

THREE 14-DAY COURSES OF TREATMENT

Melts in your mouth
Dissolves without water



Strawberry
Flavor



Actual size

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue 57 Condensed
HEADINGS: 8 pt Helvetica Neue 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue 77 Bold Condensed
TEXT: 6 pt Helvetica Neue 57 Condensed
LEADING: 6.5 and 6.7 pt
BULLETS: 5 pt Square with 2 M/S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt

STANDARD FORMAT

TYPE SIZES:

TITLE: 9 pt Helvetica Neue 77 Bold Condensed Oblique

TITLE (continued): 8 pt Helvetica Neue 77 Bold Condensed Oblique

(continued) 8 pt Helvetica Neue 57 Condensed

HEADINGS: 8 pt Helvetica Neue 77 Bold Condensed Oblique

SUBHEADINGS: 6 pt Helvetica Neue 77 Bold Condensed

TEXT: 6 pt Helvetica Neue 57 Condensed

LEADING: 6.5 and 7.0 pt

BULLETS: 5 pt Square

TELEPHONE #: 6 pt Helvetica Neue 77 Bold Condensed

HEAVY LINES: 1.5 pt

HAIRLINES: 0.5 pt

<p>Drug Facts</p> <p>Active ingredient (in each tablet)</p> <p>Unaspirin 20 mg, a low-dose aspirin to help reduce</p>	<p>Use</p> <p>• Take 1 or 2 tablets of heartburn (occur 2 or more days a week)</p> <p>• Not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p>	<p>Warnings</p> <p>Always alert: do not use if you are allergic to aspirin</p> <p>Do not use if you have:</p> <ul style="list-style-type: none"> • trouble (pain) swallowing food, vomiting with blood, or bloody or black stools • heartburn with light-headedness, swelling, or dizziness • dizziness or lightheadedness <p>These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before you use if you have:</p> <ul style="list-style-type: none"> • heartburn over 3 months, thinning or bleeding of the skin • frequent wheezing, particularly with heartburn • unexplained weight loss • nausea or vomiting • stomach pain <p>Ask a doctor or pharmacist before you use if you are taking:</p> <ul style="list-style-type: none"> • warfarin, clopidogrel, or other blood-thinning medicines • prescription anti-tingal or anti-seizure medicines • diazepam (anxiety medicine) • digoxin (heart medicine) • bromocriptine or any dopamine-related (immune system medicines) • prescription anti-ulcer or ulcer medicines for HIV infection • methotrexate (cancer medicine) 	<p>Other information</p> <ul style="list-style-type: none"> • Keep the carton, it contains important information. • Store at 20-25°C (68-77°F); keep product out of high heat and moisture.
<p>Drug Facts (continued)</p> <p>Stop use and ask a doctor if:</p> <ul style="list-style-type: none"> • your heartburn continues or worsens • you need to take this product for more than 14 days • you get dizziness • you develop a rash or joint pain <p>If you get a bad stomachache, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222).</p>	<p>Directions</p> <ul style="list-style-type: none"> • For adults 18 years of age and older • The product is to be used once daily (every 24 hours), every day for 14 days • It may take 1 to 4 days for full effect; some people get complete relief • Symptoms within 24 hours • Take 1 tablet once in the morning • Take 1 or 2 tablets with or without water • Place the tablet on tongue; tablet dissolves in water • Do not crush or chew tablets • Take every day for 14 days • Do not take more than 1 tablet a day • Do not use for more than 14 days or more often than every 4 months • Do not take for more than 14 days or more often than every 4 months unless directed by a doctor <p>Repeated 14-Day Course (if needed)</p> <ul style="list-style-type: none"> • You may repeat a 4-day course every 4 months • Do not take for more than 14 days or more often than every 4 months unless directed by a doctor 		

100

1Y930 FA C2-M

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking



NDC 64239-XXX-02

MELTech™
Melts In Your Mouth

NEW DOSAGE FORM!

Value 2 Pack

Omeprazole

Delayed Release

Orally Disintegrating Tablets 20 mg

Acid Reducer

Treats **Frequent** Heartburn!
Occurring **2 or More Days a Week**

28 TABLETS

TWO 14-DAY COURSES
OF TREATMENT

Melts in your mouth
Dissolves without water

Strawberry
Flavor

Actual
size

9-13-17

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue 57 Condensed
HEADINGS: 8 pt Helvetica Neue 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue 77 Bold Condensed
TEXT: 6 pt Helvetica Neue 57 Condensed
LEADING: 6.5 and 6.7 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt

CODE AREA

7-70 WJ CCB I I 1933 FA C2-L

Drug Facts (continue)

Inactive ingredients: anhydrous methacrylate copolymer, ascorbic acid, ethyl alcohol, colloidal silicon dioxide, croscollon, ferric oxide, flax oil, hypromellose, hydroxyethylcellulose, iron trioxide black, lecithin, microcrystalline cellulose, methylparaben, polyethylene glycol, polyethylene glycol stearate, sodium lauryl sulfate, sorbitol, sucrose, sugar spheres, waxes, xanthan gum.

Questions or comments? 1-800-799-2600; weekdays 7:30 AM to 5:00 PM EST

Drug Facts (continued)

■ **Do not use and ask a doctor if:**
■ you have heartburn, constipation or worsens
■ you need to take the product for more than 14 days
■ you need to take more than 1 course of treatment every 4 months
■ you get diarrhea
■ you develop a rash or joint pain

Keep out of reach of children. Ask a health professional before use, or ask a pharmacist for more information.

■ **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222).

Directions

■ **For adults 18 years of age and older**
■ this product is to be used once a day every 24 hours, every day for 14 days
■ if you take 1 to 4 days for full effect; some people get complete relief after 2 to 4 days

14-Day Course of Treatment

■ **Take 1 tablet before eating in the morning**
■ **do not eat or drink on an empty stomach**
■ **place the tablet on tongue; tablet dissolves with or without water.**
■ **take every day for 14 days**
■ **do not take more than 1 tablet a day**
■ **do not use for more than 14 days unless directed by your doctor.**

Repeated 14-Day Course (if needed)

■ **you may repeat a 14-day course every 4 months or often less than 14 days or more often as every 4 months as recommended by a doctor**
■ **children under 18 years of age ask a doctor before use.** Heartburn in children may sometimes be caused by a serious condition.
■ **read the directions and warnings before use**
■ **keep the carton, to contain important information.**
■ **store at 20°-25°C (68°-77°F); keep product out of high heat and moisture.**

Omeprazole

Delayed Release
Orally Disintegrating Tablets 20 mg

Safety Feature –
Do not use if printed tablet
blister unit is open or torn.

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

NDC 64239-XXX-03

NEW DOSAGE FORM!

Omeprazole

Delayed Release
Orally Disintegrating Tablets 20 mg
Acid Reducer

Treats Frequent Heartburn!
Occurring 2 or More Days a Week

42 TABLETS

THREE 14-DAY COURSES OF TREATMENT

Melts in your mouth
Dissolves without water

Strawberry Flavor

Actual
size

Reference ID: 4227553

9-13-17

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue 57 Condensed
HEADINGS: 8 pt Helvetica Neue 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue 77 Bold Condensed
TEXT: 6 pt Helvetica Neue 57 Condensed
LEADING: 6.5 and 7.0 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt

9-13-17

The diagram illustrates the layout of the blister pack. It is divided into four main sections by pink lines:

- Barcode Section:** A barcode with the numbers 0 00000 00000 0 below it.
- Tips for Managing Heartburn:**
 - Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking
- Safety Feature:**

Safety Feature –
Do not use if printed tablet blister unit is open or torn.
- Blister Unit:** A circular diagram representing the blister unit, with a crosshair in the center.

STANDARD FORMAT

TYPE SIZES:

TITLE: 9 pt Helvetica Neue 77 Bold Condensed Oblique

TITLE (continued): 8 pt Helvetica Neue 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue 57 Condensed

HEADINGS: 8 pt Helvetica Neue 77 Bold Condensed Oblique

SUBHEADINGS: 6 pt Helvetica Neue 77 Bold Condensed

TEXT: 6 pt Helvetica Neue 57 Condensed

LEADING: 6.5 and 7.0 pt


BULLETS: 5 pt Square

TELEPHONE #: 6 pt Helvetica Neue 77 Bold Condensed

HEAVY LINES: 1.5 pt

HAIRLINES: 0.5 pt


9-13-17



Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Safety Feature –
Do not use if printed tablet blister unit is open or torn.



STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue 57 Condensed
HEADINGS: 8 pt Helvetica Neue 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue 77 Bold Condensed
TEXT: 6 pt Helvetica Neue 57 Condensed
LEADING: 6.5 and 7.0 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt

9-13-17

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue 57 Condensed
HEADINGS: 8 pt Helvetica Neue 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue 77 Bold Condensed
TEXT: 6 pt Helvetica Neue 57 Condensed
LEADING: 6.5 and 7.0 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt

9-13-17

[illegible]

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue 57 Condensed
HEADINGS: 8 pt Helvetica Neue 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue 77 Bold Condensed
TEXT: 6 pt Helvetica Neue 57 Condensed
LEADING: 6.5 and 7.0 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt

9-13-17

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals

Safety Feature –
Do not use if printed tablet blister unit is open or torn.

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue 77 Bold Condensed Oblique
(continued) **8 pt Helvetica Neue 57 Condensed**
HEADINGS: 8 pt Helvetica Neue 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue 77 Bold Condensed
TEXT: 6 pt Helvetica Neue 57 Condensed
LEADING: 6.5 and 7.0 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt

9-13-17

The diagram illustrates the layout of the product box. It is divided into four main sections by pink lines:

- Top Left:** A large empty rectangular area.
- Top Right:** A large empty rectangular area.
- Bottom Left:** Contains a barcode with the numbers "0 00000 00000 0" below it.
- Bottom Center:** A section titled "Tips for Managing Heartburn" with a list of instructions:
 - Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking
- Bottom Right:** A section titled "Safety Feature –" with the instruction: "Do not use if printed tablet blister unit is open or torn." Below this text is a circular symbol with a crosshair inside.

<p>Drug Facts (continued)</p> <p>Inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscopollose, ferric oxide, flavoc, hypromellose, hypromellose phthalate, maleate, maleate codexin, mannitol, microcrystalline cellulose, propylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate</p>	<p>Questions or comments? 1-800-719-9280; weekdays 7:30 AM to 5:00 PM EST</p>
<p>Made in Israel</p> <p>Manufactured by:</p> <p>Dexcel Pharma Technologies Ltd</p> <p>10 Hakdima Street, Yokneam 2069200, Israel</p>	

1995 FA C2-E

<p>Drug Facts</p> <p>Active ingredient (in each tablet)</p> <p>Onepride 20 mg, Acid reducer</p>	<p>Use</p> <p>Take 1 frequent heartburn (occurs 2 or more days a week) and 1 bed at night for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p>	<p>Warnings</p> <p>Do not use if you have:</p> <ul style="list-style-type: none"> • a history of pain swallowing food, vomiting, or blood or black stools • heartburn with light-headedness, sweating, or dizziness • chest or shoulder pain with shortness of breath, sweating, pain spreading to jaw, neck, or shoulder • frequent chest pain • frequent heartburn signs <p>This may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before you use if you have:</p> <ul style="list-style-type: none"> • had heart burn over 3 months. This may be a sign of a more serious condition • frequent wheezing, particularly with heart burn • unexplained weight loss • nausea or vomiting • stomach pain <p>Ask a doctor if you are taking:</p> <ul style="list-style-type: none"> • warfarin, digoxin, or other blood-thinning medicines • prescription and/or over-the-counter medicines • cimetidine (antacid medicine) • ceftriaxone or any other oral antibiotics (antibiotics) • methoprene (heart medicine) 	<p>Directions</p> <ul style="list-style-type: none"> • If you are 18 years of age and older: <ul style="list-style-type: none"> • take 1 tablet once a day (every 24 hours), every day for 14 days • If you are 1 to 17 years of age for full effect: some people get complete relief of symptoms within 24 hours • 14-Day Course of 1 treatment • Take 1 tablet before eating in the morning • Do not eat or use alcohol • Place the tablet on tongue; tablet dissolves with or without water • Take every day for 14 days • Do not take more than 1 tablet a day • Do not use for more than 14 days unless directed by your doctor • Do not take this medicine with alcohol <p>Repeated 14-Day Course (if needed)</p> <ul style="list-style-type: none"> • You may repeat a 14-day course every 4 months • Do not take more than 14 days or more often than every 4 months unless directed by a doctor <p>Children under 18 years of age ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p>	<p>Other information</p> <ul style="list-style-type: none"> • Read the directions and warnings before use • Keep this out of reach of children. Important information • Store at 20-25°C (68-77°F); keep away from heat and moisture 	<p>Drug Facts (continued)</p> <p>Stop use and ask a doctor if:</p> <ul style="list-style-type: none"> • you need heartburn continues or worsens • you need take this product for more than 14 days • you need take more than 1 course of treatment every 4 months • you get diarrhea • you develop a rash or joint pain <p>Keep out of reach of children. Ask a health professional before use, in case of overdose or, get medical help or contact a Poison Control Center if necessary (1-800-222-1222).</p>	<p>Warnings</p> <p>At any time do not use if you are allergic to omeprazole</p>	<p>Directions</p> <ul style="list-style-type: none"> • If you are 18 years of age and older: <ul style="list-style-type: none"> • take 1 tablet once a day (every 24 hours), every day for 14 days • If you are 1 to 17 years of age for full effect: some people get complete relief of symptoms within 24 hours • 14-Day Course of 1 treatment • Take 1 tablet before eating in the morning • Do not eat or use alcohol • Place the tablet on tongue; tablet dissolves with or without water • Take every day for 14 days • Do not take more than 1 tablet a day • Do not use for more than 14 days unless directed by your doctor • Do not take this medicine with alcohol <p>Repeated 14-Day Course (if needed)</p> <ul style="list-style-type: none"> • You may repeat a 14-day course every 4 months • Do not take more than 14 days or more often than every 4 months unless directed by a doctor <p>Children under 18 years of age ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p>	<p>Other information</p> <ul style="list-style-type: none"> • Read the directions and warnings before use • Keep this out of reach of children. Important information • Store at 20-25°C (68-77°F); keep away from heat and moisture 	<p>Drug Facts</p> <p>Active ingredient (in each tablet)</p> <p>Onepride 20 mg, Acid reducer</p>	<p>Use</p> <p>Take 1 frequent heartburn (occurs 2 or more days a week) and 1 bed at night for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p>	<p>Warnings</p> <p>Do not use if you have:</p> <ul style="list-style-type: none"> • a history of pain swallowing food, vomiting, or blood or black stools • heartburn with light-headedness, sweating, or dizziness • chest or shoulder pain with shortness of breath, sweating, pain spreading to jaw, neck, or shoulder • frequent chest pain • frequent heartburn signs <p>This may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before you use if you have:</p> <ul style="list-style-type: none"> • had heart burn over 3 months. This may be a sign of a more serious condition • frequent wheezing, particularly with heart burn • unexplained weight loss • nausea or vomiting • stomach pain <p>Ask a doctor if you are taking:</p> <ul style="list-style-type: none"> • warfarin, digoxin, or other blood-thinning medicines • prescription and/or over-the-counter medicines • cimetidine (antacid medicine) • ceftriaxone or any other oral antibiotics (antibiotics) • methoprene (heart medicine) 	<p>Directions</p> <ul style="list-style-type: none"> • If you are 18 years of age and older: <ul style="list-style-type: none"> • take 1 tablet once a day (every 24 hours), every day for 14 days • If you are 1 to 17 years of age for full effect: some people get complete relief of symptoms within 24 hours • 14-Day Course of 1 treatment • Take 1 tablet before eating in the morning • Do not eat or use alcohol • Place the tablet on tongue; tablet dissolves with or without water • Take every day for 14 days • Do not take more than 1 tablet a day • Do not use for more than 14 days unless directed by your doctor • Do not take this medicine with alcohol <p>Repeated 14-Day Course (if needed)</p> <ul style="list-style-type: none"> • You may repeat a 14-day course every 4 months • Do not take more than 14 days or more often than every 4 months unless directed by a doctor <p>Children under 18 years of age ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p>	<p>Other information</p> <ul style="list-style-type: none"> • Read the directions and warnings before use • Keep this out of reach of children. Important information • Store at 20-25°C (68-77°F); keep away from heat and moisture 	<p>Drug Facts</p> <p>Active ingredient (in each tablet)</p> <p>Onepride 20 mg, Acid reducer</p>	<p>Use</p> <p>Take 1 frequent heartburn (occurs 2 or more days a week) and 1 bed at night for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p>	<p>Warnings</p> <p>Do not use if you have:</p> <ul style="list-style-type: none"> • a history of pain swallowing food, vomiting, or blood or black stools • heartburn with light-headedness, sweating, or dizziness • chest or shoulder pain with shortness of breath, sweating, pain spreading to jaw, neck, or shoulder • frequent chest pain • frequent heartburn signs <p>This may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before you use if you have:</p> <ul style="list-style-type: none"> • had heart burn over 3 months. This may be a sign of a more serious condition • frequent wheezing, particularly with heart burn • unexplained weight loss • nausea or vomiting • stomach pain <p>Ask a doctor if you are taking:</p> <ul style="list-style-type: none"> • warfarin, digoxin, or other blood-thinning medicines • prescription and/or over-the-counter medicines • cimetidine (antacid medicine) • ceftriaxone or any other oral antibiotics (antibiotics) • methoprene (heart medicine) 	<p>Directions</p> <ul style="list-style-type: none"> • If you are 18 years of age and older: <ul style="list-style-type: none"> • take 1 tablet once a day (every 24 hours), every day for 14 days • If you are 1 to 17 years of age for full effect: some people get complete relief of symptoms within 24 hours • 14-Day Course of 1 treatment • Take 1 tablet before eating in the morning • Do not eat or use alcohol • Place the tablet on tongue; tablet dissolves with or without water • Take every day for 14 days • Do not take more than 1 tablet a day • Do not use for more than 14 days unless directed by your doctor • Do not take this medicine with alcohol <p>Repeated 14-Day Course (if needed)</p> <ul style="list-style-type: none"> • You may repeat a 14-day course every 4 months • Do not take more than 14 days or more often than every 4 months unless directed by a doctor <p>Children under 18 years of age ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p>	<p>Other information</p> <ul style="list-style-type: none"> • Read the directions and warnings before use • Keep this out of reach of children. Important information • Store at 20-25°C (68-77°F); keep away from heat and moisture 	<p>Drug Facts</p> <p>Active ingredient (in each tablet)</p> <p>Onepride 20 mg, Acid reducer</p>	<p>Use</p> <p>Take 1 frequent heartburn (occurs 2 or more days a week) and 1 bed at night for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p>	<p>Warnings</p> <p>Do not use if you have:</p> <ul style="list-style-type: none"> • a history of pain swallowing food, vomiting, or blood or black stools • heartburn with light-headedness, sweating, or dizziness • chest or shoulder pain with shortness of breath, sweating, pain spreading to jaw, neck, or shoulder • frequent chest pain • frequent heartburn signs <p>This may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before you use if you have:</p> <ul style="list-style-type: none"> • had heart burn over 3 months. This may be a sign of a more serious condition • frequent wheezing, particularly with heart burn • unexplained weight loss • nausea or vomiting • stomach
--	--	--	---	--	---	--	---	--	--	--	--	---	--	--	--	--	---	--	--	--	---

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Safety Feature –
Do not use if printed tablet
blister unit is open or torn.



Actual size

9-13-17

[illegible]

1Y955 FA C2-F

[illegible]

**Safety Feature –
Do not use if printed tablet
blister unit is open or torn.**

00000000


9-13-17

	Drug Facts (continue d)
	Inactive ingredients: amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscarmellose, fennel oil, hypromellose, hypromellose phthalate, methylcellulose, mannitol, microcrystalline cellulose, polyethylene glycol, sodium lauryl sulfate, sodium stearoyl fumarate, sorbitol, sucrose, sugar spheres, etc., titanium dioxide, hydroxypropyl methylcellulose.
	Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5 PM EST
Made in Israel	
Manufactured by:	
Dexcel Pharma Technologies Ltd	
10 Hahadma Street, Yokneam 2069200, Israel	

CODE AREA

11955 FA CZ-G

<p>Drug Facts</p> <p>Active ingredient (in each tablet)</p> <p>Omeprazole 20 mg, ... acid reducer</p>	<p>Use</p> <p>■ Eat 4 equivalent heartburn (occurs 2 or more days a week)</p> <p>■ Take 1 tablet for immediate relief of heartburn (this drug may take 1 to 4 days for full effect)</p>	<p>Warnings</p> <p>■ Do not use if you have:</p> <ul style="list-style-type: none"> ■ a stomach or intestinal ulcer, or blood or ■ black stools <p>■ Heartburn with shortness of breath, sweating, pain spreading to shoulder pain with shortness of breath, or lightheadedness or dizziness</p> <p>■ 4 equivalent chest pain</p> <p>■ Stomach or intestinal ulcer, or lightheadedness or dizziness</p> <p>■ These may be signs of a serious condition. See your doctor.</p>	<p>Ask a doctor before you use if you have:</p> <ul style="list-style-type: none"> ■ had heartburn over 3 months. This may be a sign of a more serious condition ■ a recent wheezing, particularly with heart burn ■ unexplained weight loss ■ stomach pain <p>Ask a pharmacist before use if you are taking:</p> <ul style="list-style-type: none"> ■ a low iron, digoxin or a diuretic (blood-thinning medicines) ■ prescription anti-ulcer or anti-acid medicines ■ diazepam (anxiety medicine) ■ alcohol (heart medicine) ■ baclofen or any cytochrome P450 (immune system medicines) ■ prescription anti-ulcer or acid (medicines for HIV infection) ■ methoprostal (anti-inflammation medicine) 	<p>Other information</p> <ul style="list-style-type: none"> ■ Store at 20-25°C (68-77°F); keep product out of high heat and moisture ■ Keep the container closed until needed ■ Read the directions and warnings before use 	<p>Directions</p> <ul style="list-style-type: none"> ■ For adult 18 years of age and older ■ The product is to be used once a day (every 24 hours), every day for 14 days ■ If any pain, 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours ■ Take 1 dose of 1 tablet ■ Take 1 tablet before eating in the morning ■ Do not crush or chew tablets ■ Place the tablet on tongue; tablet dissolves, with or without water. ■ Take one glass of water ■ Take every day for 14 days ■ Do not take more than 1 tablet a day ■ Do not use for more than 14 days unless directed by your doctor ■ Do not take this medicine with alcohol <p>Repeated 14-Day Course (if needed)</p> <ul style="list-style-type: none"> ■ You may repeat a 14-day course every 4 months ■ Do not take more than 14 days or more often than every 4 months unless directed by a doctor 	<p>Children under 18 years of age: Ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p>
---	--	---	--	---	--	--



Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Safety Feature –

Do not use if printed tablet blister unit is open or torn.

NDC 64239-XXX-03

NEW DOSAGE FORM!

Bonus Size! 14 Free

Omeprazole

**Delayed Release
Orally Disintegrating Tablets 20 mg
Acid Reducer**

Treats **Frequent** Heartburn!
Occurring 2 or More Days a Week

42 TABLETS

THREE 14-DAY COURSES
OF TREATMENT

Melts in your mouth
Dissolves without water

**Strawberry
Flavor**

**Actual
size**

TYPE SIZES:
TITLE: 9 pt Helvetica Neue 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue 77 Bold Condensed Oblique
 (continued) **8 pt Helvetica Neue 57 Condensed**
HEADINGS: 8 pt Helvetica Neue 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue 77 Bold Condensed
TEXT: 6 pt Helvetica Neue 57 Condensed
LEADING: 6.5 and 6.7 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt

[illegible]

CODE AREA

H-70 VA 00611

Omeprazole

Delayed Release
Orally Disintegrating Tablets 20 mg

[illegible]

Drug Facts (continued)

Stop use and ask a doctor if:

- you're heartburn continues or worsens
- you need to take more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

Keep out of reach of children. Ask a health professional for use, if needed.

Directions

- For adults 18 years of age and older
- his product to be used once a day (every 24 hours), every day for 14 days
- his product to be used once a day (every 24 hours) for 14 days for full effects; some people get complete relief after 7 to 14 days

14-Day Course of Treatment

- Take 1 tablet before eating in the morning
- do not eat or drink on an empty stomach
- place the tablet on tongue; tablet dissolves with or without water.
- The tablet can also be swallowed whole with water.
- Take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- do not take this medicine with alcohol

Repeated 14-Day Course (if needed)

- You may repeat a 14-day course every 4 months
- do not take for more than 1 day or more often than every 4 months unless directed by a doctor

Children under 18 years of age ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton in a cool, dry place
- store at 20°-25°C (68°-77°F); keep product out of high heat and moisture

Drug Facts	
Active ingredient (in each tablet)	Omeprazole 20 mg
Purpose	Acid reducer

A barcode consisting of vertical black bars of varying widths, used for document identification. Below the bars are three groups of numbers: "0", "00000", and "00000".

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

**Safety Feature –
Do not use if printed tablet
blister unit is open or torn.**



NEW DOSAGE FORM!

Bonus! + 14 Free

Omeprazole

Delayed Release
Orally Disintegrating Tablets 20 mg
Acid Reducer

Treats Frequent Heartburn!
Occurring 2 or More Days a Week

42 TABLETS **THREE 14-DAY COURSES**
OF TREATMENT

Melts in your mouth
Dissolves without water



Strawberry
Flavor



Actual
si

TYPE SIZES:
TITLE: 9 pt Helvetica Neue 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue 57 Condensed
HEADINGS: 8 pt Helvetica Neue 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue 77 Bold Condensed
TEXT: 6 pt Helvetica Neue 57 Condensed
LEADING: 6.5 and 6.7 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt

<p>Drug Facts (continued)</p> <p>Inactive ingredients: amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscollon, ferric oxide, flavoc, hypromellose, hypromellose phthalate, methylcellulose, mannitol, microcrystalline cellulose, polyethylene glycol, sodium lauryl sulfate, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, hydroxypropyl methylcellulose.</p>	<p>Questions or comments? 1-800-719-9280; weekdays 7:30 AM to 5 PM EST</p> <p>Made in Israel</p> <p>Manufactured by: Dexcel Pharma Technologies Ltd 10 Hahadima Street, Yokneam 2069200, Israel</p>
---	---

1995 FA 022

Omeprazole

Delayed Release
Orally Disintegrating Tablets 20 mg

Drug Facts	Active ingredient (in each tablet)	Purpose
<p>Use</p> <p>Oneprazole 20 mg</p> <p>Acid reducer</p>	<p>Warnings</p> <p>1 to 4 days for full effect</p> <p>• Fast frequent heartburn (occurs 2 or more days a week)</p> <p>• Not intended for immediate relief of heartburn; this drug may take</p> <p>Any allergy do not use if you are allergic to omeprazole</p> <p>Do not use if you have:</p> <ul style="list-style-type: none"> • alcohol pain swallowing food, vomiting with blood, or blood or • black stools • heart or stomach pain with shortness of breath; sweating; pain spreading • heartburn with light-headedness, sweating or dizziness • hoarseness, neck or shoulder, or lightheadedness <p>1 frequent chest pain</p> <p>These may be signs of a serious condition. See your doctor.</p>	<p>Ask a doctor before you use it if you have:</p> <ul style="list-style-type: none"> • had heartburn over 3 months that may be a sign of a more serious condition • taken heartburn medicine with heartburn • unexplained weight loss • nausea or vomiting • stomach pain
	<p>Ask a doctor or pharmacist if you are taking:</p> <ul style="list-style-type: none"> • warfarin, digoxin, or other blood-thinning medicines • prescription and liquid or other medicines • dicyclanil (heart medicine) • glycolic (heart medicine) • ceftriaxone or any combination (immune system medicines) • prescription and oral or liquid medicines for HIV infection) • methotrexate (arthritis medicine) 	

Key facts (continued)

You see and ask a doctor if:

- your cat has vomited or worse
- you need to take his product for more than 14 days
- you need to take him product for more than 1 course of treatment every 4 months

you get diarrhea ■ you develop a rash or joint pain

For pet oral care - feeding, ask a health professional before use.

If your cat needs dental care, in case of overdose, get medical help or contact a Petco Control Center right away (toll-free 800-222-1222).

Infections

For adult 18 year old and older:
The product is to be used once a day (every 24 hours), every day for 14 days
It may take 1 to 4 weeks for full effect; some people get complete relief of symptoms within 24 hours

14-day courses of treatment

- Take 1 tablet before eating in the morning
- Do not eat until 1 hour after taking the tablet
- The tablets can also be swallowed whole with water.

Take every day for 14 days

- Do not take more than 1 tablet a day
- Do not use for more than 14 days unless told by your doctor
- Do not take his medicine with alcohol

Repeated 14-Day Course (if needed)

- You may repeat a 14-day course every 4 months
- Do not take for more than 14 days or more often than every 4 months if needed directed by a doctor

Children under 18 years of age as well as a doctor before use. A heartburn in children may sometimes be caused by a serious condition.

Other information

Read the directions and warnings before use.
Keep the cat in the area important information.
Store at 20-25°C (68-77°F); keep product out of high heat and moisture.

A barcode consisting of vertical black bars of varying widths on a white background. Below the bars are the numbers "000000000000".

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Safety Feature –
Do not use if printed tablet
blister unit is open or torn.

NDC 64239-XXX-03

NEW DOSAGE FORM!

Bonus! 14 Tablets FREE

Omeprazole

Delayed Release
Orally Disintegrating Tablets 20 mg
Acid Reducer

Treats Frequent Heartburn!
Occurring 2 or More Days a Week

42 TABLETS

THREE 14-DAY COURSES OF TREATMENT

Melts in your mouth
Dissolves without water



Strawberry
Flavor



**Actual
Size**

TYPE SIZES:
TITLE: 9 pt Helvetica Neue 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue 77 Bold Condensed Oblique
 (continued) 8 pt Helvetica Neue 57 Condensed
HEADINGS: 8 pt Helvetica Neue 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue 77 Bold Condensed
TEXT: 6 pt Helvetica Neue 57 Condensed
LEADING: 6.5 and 6.7 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt

Proposed Draft Labeling
NDA 209400
Omeprazole Delayed Release Orally
Disintegrating Tablets 20 mg
14 Tablets Peel Off Coupon

FRONT PANEL

peel off coupon

Instant Redeemable Coupon - Expires XX-XX-XX

SAVE \$X NOW!

On This Purchase

**MELTech™**
Melts In Your Mouth

NDC 64239-XXX-01

NEW DOSAGE FORM!

Omeprazole
Delayed Release
Orally Disintegrating Tablets 20 mg
Acid Reducer

Strawberry
Flavor

Treats Frequent Heartburn!
Occurring 2 or More Days a Week

Melts in your mouth
Dissolves without water

14 TABLETS ONE 14-DAY COURSE
OF TREATMENT

Actual
size

1Y974 FA CP1

BACK PANEL

SAVE \$X NOW!

MANUFACTURER'S COUPON EXPIRATION DATE XX/XX/XXXX

SAVE \$X^{XX} on your purchase of
14 COUNT SIZE
"Store Brand" Omeprazole
Delayed Release Orally Disintegrating Tablets 20 mg

**COUPON NOT SUBJECT
TO DOUBLING**
Customer pays any sales tax.

CONSUMER: Only one coupon is redeemable per purchase. RETAILER: This coupon will be redeemable for face value on specified products (plus a handling fee) if submitted in compliance with the terms of this offer. Cash value 1/100th of one cent. Limit one coupon per customer. Do Not Double. Coupon is non-reproducible. Coupons cannot be combined. Mail properly handled coupons to: L. Perrigo Company (#1566), NCH Marketing, PO Box 880001, El Paso, TX 88588-0001


0 000000 000000 0

Omeprazole
Delayed Release
Orally Disintegrating Tablets 20 mg

14 TABLETS

Proposed Draft Labeling
NDA 209400
Omeprazole Delayed Release Orally
Disintegrating Tablets 20 mg
28 Tablets Peel Off Coupon

FRONT PANEL

peel off coupon

Instant Redeemable Coupon - Expires XX-XX-XX

SAVE \$X NOW!

On This Purchase

**MELTech™**
Melts In Your Mouth

NDC 64239-XXX-02

NEW DOSAGE FORM!

Omeprazole
Delayed Release
Orally Disintegrating Tablets 20 mg
Acid Reducer

Strawberry
Flavor

Treats Frequent Heartburn!
Occurring 2 or More Days a Week

28 TABLETS TWO 14-DAY COURSES
OF TREATMENT

Melts in your mouth
Dissolves without water

Actual
Size

1Y930 FA CP1

BACK PANEL

SAVE \$X NOW!

MANUFACTURER'S COUPON EXPIRATION DATE XX/XX/XXXX

SAVE \$X^{XX} on your purchase of
28 COUNT SIZE
"Store Brand" Omeprazole
Delayed Release Orally Disintegrating Tablets 20 mg

CONSUMER: Only one coupon is redeemable per purchase. RETAILER: This coupon will be redeemable for face value on specified products (plus a handling fee) if submitted in compliance with the terms of this offer. Cash value 1/100th of one cent. Limit one coupon per customer. Do Not Double. Coupon is non-reproducible. Coupons cannot be combined. Mail properly handled coupons to: L. Perrigo Company (#1565), NCH Marketing, PO Box 880001, El Paso, TX 81589-0001

COUPON NOT SUBJECT TO DOUBLING
Customer pays any sales tax.


0 000000 000000 0

Omeprazole
Delayed Release
Orally Disintegrating Tablets 20 mg

28 TABLETS

Proposed Draft Labeling
NDA 209400
Omeprazole Delayed Release Orally
Disintegrating Tablets 20 mg
42 Tablets Peel Off Coupon

FRONT PANEL

peel off coupon

Instant Redeemable Coupon - Expires XX-XX-XX

SAVE \$X NOW!

On This Purchase

**MELTech™**
Melts In Your Mouth

NDC 64239-XXX-03

NEW DOSAGE FORM!

Omeprazole
Delayed Release
Orally Disintegrating Tablets 20 mg
Acid Reducer

Strawberry
Flavor

Treats Frequent Heartburn!
Occurring 2 or More Days a Week

42 TABLETS

THREE 14-DAY COURSES
OF TREATMENT

Melts in your mouth
Dissolves without water

Actual
size

1Y955 FA CP1

BACK PANEL

SAVE \$X NOW!

MANUFACTURER'S COUPON EXPIRATION DATE XX/XX/XXXX

SAVE \$X^{XX} on your purchase of
42 COUNT SIZE
"Store Brand" Omeprazole
Delayed Release Orally Disintegrating Tablets 20 mg

CONSUMER: Only one coupon is redeemable per purchase. RETAILER: This coupon will be redeemable for face value on specified products (plus a handling fee) if submitted in compliance with the terms of this offer. Cash value 1/100th of one cent. Limit one coupon per customer. Do Not Double. Coupon is non-reproducible. Coupons cannot be combined. Mail properly handled coupons to: L. Perrigo Company (#1566), NCH Marketing, PO Box 880001, St Paul, TX 88688-0001

COUPON NOT SUBJECT TO DOUBLING
Customer pays any sales tax.


0 000000 000000 0

Omeprazole
Delayed Release
Orally Disintegrating Tablets 20 mg

42 TABLETS

Reference ID: 4227553

9-13-17

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

KAREN M MAHONEY
02/28/2018