


PENTASA[®]
(mesalamine)
Controlled-Release Capsules 250 mg and 500 mg
Prescribing Information as of 07/2019
Rx only

DESCRIPTION

PENTASA (mesalamine) for oral administration is a controlled-release formulation of mesalamine, an aminosalicylate anti-inflammatory agent for gastrointestinal use.

Chemically, mesalamine is 5-amino-2-hydroxybenzoic acid. It has a molecular weight of 153.14.

The structural formula is:



Each 250-mg capsule contains 250 mg of mesalamine. It also contains the following inactive ingredients: acetylated monoglyceride, castor oil, colloidal silicon dioxide, ethylcellulose, hydroxypropyl methylcellulose, starch, stearic acid, sugar, talc, and white wax. The capsule shell contains D&C Yellow #10, FD&C Blue #1, FD&C Green #3, gelatin, titanium dioxide, and other ingredients.

Each 500-mg capsule contains 500 mg of mesalamine. It also contains the following inactive ingredients: acetylated monoglyceride, castor oil, colloidal silicon dioxide, ethylcellulose, hydroxypropyl methylcellulose, starch, stearic acid, sugar, talc, and white wax. The capsule shell contains FD&C Blue #1, gelatin, titanium dioxide, and other ingredients.

CLINICAL PHARMACOLOGY

Sulfasalazine is split by bacterial action in the colon into sulfapyridine (SP) and mesalamine (5-ASA). It is thought that the mesalamine component is therapeutically active in ulcerative colitis. The usual oral dose of sulfasalazine for active ulcerative colitis in adults is 2 to 4 g per day in divided doses. Four grams of sulfasalazine provide 1.6 g of free mesalamine to the colon.

The mechanism of action of mesalamine (and sulfasalazine) is unknown, but it appears to be topical rather than systemic. Mucosal production of arachidonic acid (AA) metabolites, both through the cyclooxygenase pathways, i.e., prostanooids, and through the lipoxygenase pathways, i.e., leukotrienes (LTs) and hydroxyeicosatetraenoic acids (HETEs), is increased in patients with chronic inflammatory bowel disease, and it is possible that mesalamine

diminishes inflammation by blocking cyclooxygenase and inhibiting prostaglandin (PG) production in the colon.

Human Pharmacokinetics and Metabolism

Absorption: PENTASA is an ethylcellulose-coated, controlled-release formulation of mesalamine designed to release therapeutic quantities of mesalamine throughout the gastrointestinal tract. Based on urinary excretion data, 20% to 30% of the mesalamine in PENTASA is absorbed. In contrast, when mesalamine is administered orally as an unformulated 1-g aqueous suspension, mesalamine is approximately 80% absorbed.

Plasma mesalamine concentration peaked at approximately 1 mcg/mL 3 hours following a 1-g PENTASA dose and declined in a biphasic manner. The literature describes a mean terminal half-life of 42 minutes for mesalamine following intravenous administration. Because of the continuous release and absorption of mesalamine from PENTASA throughout the gastrointestinal tract, the true elimination half-life cannot be determined after oral administration. N-acetyl-5-aminosalicylic acid, the major metabolite of mesalamine (5-aminosalicylic acid), peaked at approximately 3 hours at 1.8 mcg/mL, and its concentration followed a biphasic decline. Pharmacological activities of N-acetyl-5-aminosalicylic acid are unknown, and other metabolites have not been identified.

Oral mesalamine pharmacokinetics were nonlinear when PENTASA capsules were dosed from 250 mg to 1 g four times daily, with steady-state mesalamine plasma concentrations increasing about nine times, from 0.14 mcg/mL to 1.21 mcg/mL, suggesting saturable first-pass metabolism. N-acetyl-5-aminosalicylic acid pharmacokinetics were linear.

Elimination: About 130 mg free mesalamine was recovered in the feces following a single 1-g PENTASA dose, which was comparable to the 140 mg of mesalamine recovered from the molar equivalent sulfasalazine tablet dose of 2.5 g. Elimination of free mesalamine and salicylates in feces increased proportionately with PENTASA dose. N-acetyl-5-aminosalicylic acid was the primary compound excreted in the urine (19% to 30%) following PENTASA dosing.

CLINICAL TRIALS

In two randomized, double-blind, placebo-controlled, dose-response trials (UC-1 and UC-2) of 625 patients with active mild to moderate ulcerative colitis, PENTASA, at an oral dose of 4 g/day given 1 g four times daily, produced consistent improvement in prospectively identified primary efficacy parameters, PGA, Tx F, and SI as shown in the table below.

The 4-g dose of PENTASA also gave consistent improvement in secondary efficacy parameters, namely the frequency of trips to the toilet, stool consistency, rectal bleeding, abdominal/rectal pain, and urgency. The 4-g dose of PENTASA induced remission as assessed by endoscopic and symptomatic endpoints.

In some patients, the 2-g dose of PENTASA was observed to improve efficacy parameters measured. However, the 2-g dose gave inconsistent results in primary efficacy parameters across the two adequate and well controlled trials.

Parameter Evaluated	Clinical Trial UC-1			Clinical Trial UC-2		
	PL (n=90)	PENTASA		PL (n=83)	PENTASA	
		4 g/day (n=95)	2 g/day (n=97)		4 g/day (n=85)	2 g/day (n=83)
PGA	36%	59%*	57%*	31%	55%*	41%
Tx F	22%	9%*	18%	31%	9%*	17%*
SI	-2.5	-5.0*	-4.3*	-1.6	-3.8*	-2.6
Remission [†]	12%	26%*	24%*	12%	27%*	12%

*p<0.05 vs placebo.

PGA: Physician Global Assessment: proportion of patients with complete or marked improvement.

Tx F: Treatment Failure: proportion of patients developing severe or fulminant UC requiring steroid therapy or hospitalization or worsening of the disease at 7 days of therapy, or lack of significant improvement by 14 days of therapy.

SI: Sigmoidoscopic Index: an objective measure of disease activity rated by a standard (15-point) scale that includes mucosal vascular pattern, erythema, friability, granularity/ulcerations, and mucopus: improvement over baseline.

[†]Defined as complete resolution of symptoms plus improvement of endoscopic endpoints. To be considered in remission, patients had a “1” score for one of the endoscopic components (mucosal vascular pattern, erythema, granularity, or friability) and “0” for the others.

INDICATIONS AND USAGE

PENTASA is indicated for the induction of remission and for the treatment of patients with mildly to moderately active ulcerative colitis.

CONTRAINDICATIONS

PENTASA is contraindicated in patients who have demonstrated hypersensitivity to mesalamine, any other components of this medication, or salicylates.

PRECAUTIONS

General

Caution should be exercised if PENTASA is administered to patients with impaired hepatic function.

Mesalamine has been associated with an acute intolerance syndrome that may be difficult to distinguish from a flare of inflammatory bowel disease. Although the exact frequency of occurrence cannot be ascertained, it has occurred in 3% of patients in controlled clinical trials of mesalamine or sulfasalazine. Symptoms include cramping, acute abdominal pain and bloody diarrhea, sometimes fever, headache, and rash. If acute intolerance syndrome is suspected, prompt withdrawal is required. If a rechallenge is performed later in order to

validate the hypersensitivity, it should be carried out under close medical supervision at reduced dose and only if clearly needed.

Patients with pre-existing skin conditions such as atopic dermatitis and atopic eczema have reported more severe photosensitivity reactions.

Renal

Caution should be exercised if PENTASA is administered to patients with impaired renal function. Single reports of nephrotic syndrome and interstitial nephritis associated with mesalamine therapy have been described in the foreign literature. There have been rare reports of interstitial nephritis in patients receiving PENTASA. In animal studies, a 13-week oral toxicity study in mice and 13-week and 52-week oral toxicity studies in rats and cynomolgus monkeys have shown the kidney to be the major target organ of mesalamine toxicity. Oral daily doses of 2400 mg/kg in mice and 1150 mg/kg in rats produced renal lesions including granular and hyaline casts, tubular degeneration, tubular dilation, renal infarct, papillary necrosis, tubular necrosis, and interstitial nephritis. In cynomolgus monkeys, oral daily doses of 250 mg/kg or higher produced nephrosis, papillary edema, and interstitial fibrosis. Patients with preexisting renal disease, increased BUN or serum creatinine, or proteinuria should be carefully monitored, especially during the initial phase of treatment. Mesalamine-induced nephrotoxicity should be suspected in patients developing renal dysfunction during treatment.

Interference with Laboratory Tests

Use of mesalamine may lead to spuriously elevated test results when measuring urinary normetanephrine by liquid chromatography with electrochemical detection because of the similarity in the chromatograms of normetanephrine and mesalamine's main metabolite, N-acetyl-5-aminosalicylic acid (N-Ac-5-ASA). Consider an alternative, selective assay for normetanephrine.

Drug Interactions

No investigations of interactions between PENTASA and other drugs have been performed; however, the following drug-drug interactions have been reported for products containing mesalamine:

The concurrent use of mesalamine with known nephrotoxic agents, including non-steroidal anti-inflammatory drugs (NSAIDs), may increase the risk of renal reactions.

The concurrent use of mesalamine with azathioprine or 6-mercaptopurine and/or any other drugs known to cause myelotoxicity may increase the risk for blood disorders, bone marrow failure, and associated complications.

Carcinogenesis, Mutagenesis, Impairment of Fertility

In a 104-week dietary carcinogenicity study of mesalamine, CD-1 mice were treated with doses up to 2500 mg/kg/day and it was not tumorigenic. For a 50-kg person of average height (1.46 m² body surface area), this represents 2.5 times the recommended human dose on a body surface area basis (2960 mg/m²/day). In a 104-week dietary carcinogenicity study in

Wistar rats, mesalamine up to a dose of 800 mg/kg/day was not tumorigenic. This dose represents 1.5 times the recommended human dose on a body surface area basis.

No evidence of mutagenicity was observed in an in vitro Ames test and in an in vivo mouse micronucleus test.

No effects on fertility or reproductive performance were observed in male or female rats at oral doses of mesalamine up to 400 mg/kg/day (0.8 times the recommended human dose based on body surface area).

Semen abnormalities and infertility in men, which have been reported in association with sulfasalazine, have not been seen with PENTASA capsules during controlled clinical trials.

Pregnancy

Teratogenic Effects

Published data from meta-analyses, cohort studies, and case series on the use of mesalamine during pregnancy have not reliably informed an association with mesalamine and major birth defects, miscarriage, or adverse maternal or fetal outcomes. There is no clear evidence that mesalamine exposure in early pregnancy is associated with an increased risk of major congenital malformations, including cardiac malformations. Published epidemiologic studies have important methodological limitations which hinder interpretation of the data, including inability to control for confounders, such as underlying maternal disease, maternal use of concomitant medications, and missing information on the dose and duration of use for mesalamine products. PENTASA should only be used during pregnancy if the benefits outweigh the risks.

In animal reproduction studies, oral administration of mesalamine during organogenesis to pregnant rats at doses up to 1000 mg/kg/day (5900 mg/m²) and rabbits at doses of 800 mg/kg/day (6856 mg/m²) revealed no evidence of teratogenic effects or harm to the fetus due to mesalamine.

The estimated background risk of major birth defects and miscarriage for the indicated populations is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriages in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

Nonteratogenic Effects

Published data suggest that increased disease activity is associated with the risk of developing adverse pregnancy outcomes in women with ulcerative colitis. Adverse pregnancy outcomes include preterm delivery (before 37 weeks of gestation), low birth weight (less than 2500 g) infants, and small for gestational age at birth.

Nursing Mothers

Data from published literature report the presence of mesalamine and its metabolite, N-acetyl-5-aminosalicylic acid in human milk in small amounts with relative infant doses

(RID) of 0.1% or less for mesalamine. The clinical significance of this has not been determined and there is limited experience of nursing women using mesalamine.

There are reports of diarrhea observed in breastfed infants exposed to mesalamine. The effect of PENTASA on milk production is unknown. Exercise caution if PENTASA is administered to a nursing woman and monitor breastfed infants for diarrhea.

In published lactation studies, maternal mesalamine doses from various oral and rectal formulations and products ranged from 500 mg to 4.8 g daily. The average concentration of mesalamine in milk ranged from non-detectable to 0.5 mg/L. The average concentration of N-acetyl-5-aminosalicylic acid in milk ranged from 0.2 to 9.3 mg/L. Based on these concentrations, estimated infant daily dosages for an exclusively breastfed infant are 0 to 0.075 mg/kg/day of mesalamine (RID 0% to 0.1%) and 0.03 to 1.4 mg/kg/day of N-acetyl-5-aminosalicylic acid.

Pediatric Use

Safety and efficacy of PENTASA in pediatric patients have not been established.

ADVERSE REACTIONS

In combined domestic and foreign clinical trials, more than 2100 patients with ulcerative colitis or Crohn's disease received PENTASA therapy. Generally, PENTASA therapy was well tolerated. The most common events (i.e., greater than or equal to 1%) were diarrhea (3.4%), headache (2.0%), nausea (1.8%), abdominal pain (1.7%), dyspepsia (1.6%), vomiting (1.5%), and rash (1.0%).

In two domestic placebo-controlled trials involving over 600 patients with ulcerative colitis, adverse events were fewer in PENTASA (mesalamine)-treated patients than in the placebo group (PENTASA 14% vs placebo 18%) and were not dose-related. Events occurring in more than 1% are shown in the table below. Of these, only nausea and vomiting were more frequent in the PENTASA group. Withdrawal from therapy due to adverse events was more common on placebo than PENTASA (7% vs 4%).

Table 1. Adverse Events Occurring in More than 1% of Either Placebo or PENTASA Patients in Domestic Placebo-controlled Ulcerative Colitis Trials. (PENTASA Comparison to Placebo)		
Event	PENTASA n=451	Placebo n=173
Diarrhea	16 (3.5%)	13 (7.5%)
Headache	10 (2.2%)	6 (3.5%)
Nausea	14 (3.1%)	-----
Abdominal Pain	5 (1.1%)	7 (4.0%)
Melena (Bloody Diarrhea)	4 (0.9%)	6 (3.5%)
Rash	6 (1.3%)	2 (1.2%)
Anorexia	5 (1.1%)	2 (1.2%)
Fever	4 (0.9%)	2 (1.2%)
Rectal Urgency	1 (0.2%)	4 (2.3%)
Nausea and Vomiting	5 (1.1%)	-----
Worsening of Ulcerative Colitis	2 (0.4%)	2 (1.2%)
Acne	1 (0.2%)	2 (1.2%)

Clinical laboratory measurements showed no significant abnormal trends for any test, including measurement of hematological, liver, and kidney function.

The following adverse events, presented by body system, were reported infrequently (i.e., less than 1%) during domestic ulcerative colitis and Crohn’s disease trials. In many cases, the relationship to PENTASA has not been established.

Gastrointestinal: abdominal distention, anorexia, constipation, duodenal ulcer, dysphagia, eructation, esophageal ulcer, fecal incontinence, GGTP increase, GI bleeding, increased alkaline phosphatase, LDH increase, mouth ulcer, oral moniliasis, pancreatitis, rectal bleeding, SGOT increase, SGPT increase, stool abnormalities (color or texture change), thirst

Dermatological: acne, alopecia, dry skin, eczema, erythema nodosum, nail disorder, photosensitivity, pruritus, sweating, urticaria

Nervous System: depression, dizziness, insomnia, somnolence, paresthesia

Cardiovascular: palpitations, pericarditis, vasodilation

Other: albuminuria, amenorrhea, amylase increase, arthralgia, asthenia, breast pain, conjunctivitis, ecchymosis, edema, fever, hematuria, hypomenorrhea, Kawasaki-like syndrome, leg cramps, lichen planus, lipase increase, malaise, menorrhagia, metrorrhagia, myalgia, pulmonary infiltrates, thrombocythemia, thrombocytopenia, urinary frequency

One week after completion of an 8-week ulcerative colitis study, a 72-year-old male, with no previous history of pulmonary problems, developed dyspnea. The patient was subsequently diagnosed with interstitial pulmonary fibrosis without eosinophilia by one physician and

bronchiolitis obliterans with organizing pneumonitis by a second physician. A causal relationship between this event and mesalamine therapy has not been established.

Published case reports and/or spontaneous postmarketing surveillance have described infrequent instances of pericarditis, fatal myocarditis, chest pain and T-wave abnormalities, hypersensitivity pneumonitis, pancreatitis, nephrotic syndrome, interstitial nephritis, hepatitis, aplastic anemia, pancytopenia, leukopenia, agranulocytosis, or anemia while receiving mesalamine therapy. Anemia can be a part of the clinical presentation of inflammatory bowel disease. Allergic reactions, which could involve eosinophilia, can be seen in connection with PENTASA therapy.

Postmarketing Reports

The following events have been identified during post-approval use of the PENTASA brand of mesalamine in clinical practice. Because they are reported voluntarily from a population of unknown size, estimates of frequency cannot be made. These events have been chosen for inclusion due to a combination of seriousness, frequency of reporting, or potential causal connection to mesalamine:

Gastrointestinal: Reports of hepatotoxicity, including elevated liver enzymes (SGOT/AST, SGPT/ALT, GGT, LDH, alkaline phosphatase, bilirubin), hepatitis, jaundice, cholestatic jaundice, cirrhosis, and possible hepatocellular damage including liver necrosis and liver failure. Some of these cases were fatal. One case of Kawasaki-like syndrome which included hepatic function changes was also reported.

Other: Postmarketing reports of anaphylactic reaction, Stevens-Johnson syndrome (SJS), drug reaction with eosinophilia and systemic symptoms (DRESS), pneumonitis, granulocytopenia, systemic lupus erythematosus, lupus-like syndrome, acute renal failure, interstitial lung disease, hypersensitivity pneumonitis (including interstitial pneumonitis, allergic alveolitis, eosinophilic pneumonitis), chronic renal failure, nephrogenic diabetes insipidus, intracranial hypertension, angioedema and oligospermia (reversible) have been received in patients taking PENTASA.

OVERDOSAGE

Single oral doses of mesalamine up to 5 g/kg in pigs or a single intravenous dose of mesalamine at 920 mg/kg in rats were not lethal.

There is no clinical experience with PENTASA overdose. PENTASA is an aminosalicylate, and symptoms of salicylate toxicity may be possible, such as: tinnitus, vertigo, headache, confusion, drowsiness, sweating, hyperventilation, vomiting, and diarrhea. Severe intoxication with salicylates can lead to disruption of electrolyte balance and blood-pH, hyperthermia, and dehydration.

Treatment of Overdosage: Since PENTASA is an aminosalicylate, conventional therapy for salicylate toxicity may be beneficial in the event of acute overdose. This includes prevention of further gastrointestinal tract absorption by emesis and, if necessary, by gastric

lavage. Fluid and electrolyte imbalance should be corrected by the administration of appropriate intravenous therapy. Adequate renal function should be maintained.

DOSAGE AND ADMINISTRATION

The recommended dosage for the induction of remission and the symptomatic treatment of mildly to moderately active ulcerative colitis is 1 g (4 PENTASA 250-mg capsules or 2 PENTASA 500-mg capsules) 4 times a day for a total daily dosage of 4 g. Treatment duration in controlled trials was up to 8 weeks.

PENTASA capsules may be swallowed whole, or alternatively, the capsule may be opened and the entire contents sprinkled onto applesauce or yogurt. The entire contents should be consumed immediately. The capsules and capsule contents must not be crushed or chewed.

Safety and efficacy of PENTASA in pediatric patients have not been established.

HOW SUPPLIED

PENTASA controlled-release 250-mg capsules are supplied in bottles of 240 capsules (NDC 54092-189-81). Each green and blue capsule contains 250 mg of mesalamine in controlled-release beads. PENTASA controlled-release capsules are identified with a pentagonal starburst logo and the number 2010 on the green portion and S429 250 mg on the blue portion of the capsules.

PENTASA controlled-release 500-mg capsules are supplied in bottles of 120 capsules (NDC 54092-191-12). Each blue capsule contains 500 mg of mesalamine in controlled-release beads. PENTASA controlled-release capsules are identified with a pentagonal starburst logo and S429 500 mg on the capsules.

Store at 25°C (77°F); excursions permitted to 15 to 30°C (59 to 86°F) [see USP Controlled Room Temperature].

To report SUSPECTED ADVERSE REACTIONS, contact Shire US Inc. at 1-800-828-2088 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Manufactured for **Shire US Inc.**

300 Shire Way, Lexington, MA 02421, USA

PENTASA is a registered trademark of Ferring B.V.

© 2019 Takeda Pharmaceutical Company Limited

Rev. 07/2019