



Drug Interaction Warnings:

Tagamet HB 200® affects some prescription medicines causing slightly higher levels of those medicines in the blood. Higher blood levels could lead to side effects in rare situations. If you currently take Theophylline (oral asthma medicine), Warfarin (blood thinning medicine) or Phenytoin (seizure medicine), consult your doctor before taking Tagamet HB 200®.

Brand names of some medicines which contain one of these ingredients include:

THEOPHYLLINE	THEO-DUR® Uniphy®	Theochron® Theo-24®	Slo-Bid®
WARFARIN	Coumadin®		
PHENYTOIN	Dilantin®		

There may be other medicines that contain one of these ingredients. If in doubt, about this or about possible effects of Tagamet HB 200® on any other medicines you are taking, talk to your doctor or pharmacist.

Tips for Managing Heartburn

- Do not lie flat or bend over soon after eating
- Do not eat late at night, or just before bedtime
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables
- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose fitting clothing around your stomach

Questions? 1-800-482-4394

Read the directions and warnings before taking this medication.

In clinical studies, Tagamet HB 200® was significantly better than placebo tablets in relieving and preventing heartburn symptoms.











**Drug Interaction Warnings:**

Tagamet HB 200® affects some prescription medicines causing slightly higher levels of those medicines in the blood. Higher blood levels could lead to side effects in rare situations. If you currently take Theophylline (oral asthma medicine), Warfarin (blood thinning medicine) or Phenytoin (seizure medicine), consult your doctor before taking Tagamet HB 200®.

Brand names of some medicines which contain one of these ingredients include:

THEOPHYLLINE	THEO-DUR® Uniphy®	Theochron® Theo-24®	Slo-Bid®
WARFARIN	Coumadin®		
PHENYTOIN	Dilantin®		

There may be other medicines that contain one of these ingredients. If in doubt about this or about possible effects of Tagamet HB 200® on any other medicines you are taking, talk to your doctor or pharmacist.

**Tips for Managing Heartburn**

- Do not lie flat or bend over soon after eating
- Do not eat late at night, or just before bedtime
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables
- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose fitting clothing around your stomach

**Questions?** 1-800-482-4394

**Read the directions and warnings before taking this medication.**

**In clinical studies, Tagamet HB 200® was significantly better than placebo tablets in relieving and preventing heartburn symptoms.**



-----  
**This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.**  
-----

/s/  
-----

KAREN M MAHONEY  
03/26/2019 05:03:48 PM