



Do not use if foil inner seal imprinted with "SAFETY SEAL®" is broken or missing
Active ingredient in each tablet: Famotidine 10 mg **Purpose:** Acid Reducer
Uses: •relieves heartburn associated with acid indigestion and sour stomach •prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages. **No nings:** **Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers. **Do not use:** •if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. •with other acid reducers. **Ask a doctor before use if you have:** •had heartburn over 3 months. This may be a sign of a more serious condition. •heartburn with **lightheadedness, sweating, or dizziness** •chest pain or shoulder pain with shortness of breath, sweating, spreading to arms, neck or shoulders; or lightheadedness •**frequent chest pain** •frequent wheezing, particularly with heartburn •unexplained weight loss •nausea or vomiting •stomach pain •kidney disease. **Ask a doctor or pharmacist before use if you are taking a prescription drug.** Acid reducers may interact with certain prescription drugs. **Stop use and ask a doctor if:** •your heartburn continues or worsens •you need to take this product for more than 14 days. **If pregnant or breast feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1 800 222 1222). **Directions:** •adults and children 12 years and over: •to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew. •to prevent symptoms, swallow 1 tablet with a glass of water at any time from 15 to 60 minutes before eating food or drinking beverages that cause heartburn •do not use more than 2 tablets in 24 hours •children under 12 years: ask a doctor

ORIGINAL STRENGTH
NDC 16837-872-22 **See New Warnings**

Pepcid[®]

AC
Famotidine Tablets 10 mg Acid Reducer
Just One Tablet!
Prevents & Relieves Heartburn
Due to Acid Indigestion
90 Tablets

Famotidine (active ingredient) made in Hungary. Processed in Canada
Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**
McNeil Consumer Healthcare Division, Fort Washington, PA 19004 USA ©McNeil CPC 20XX
PEPCID® and PEPCID AC® are registered trademarks of McNeil CPC

Do not use if foil seal under bottle cap printed "SAFETY SEAL®" is broken or missing.

Active ingredient (in each tablet) Famotidine 10 mg	Purpose Acid reducer
---	--------------------------------

Uses • relieves heartburn associated with acid indigestion and sour stomach • prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages. **Warnings** **Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers. **Do not use** • if you have trouble or pain swallowing food; vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

use if you have • had heartburn over 3 months. This may be a sign of a more serious condition. • heartburn with **lightheadedness, sweating, or dizziness** • chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness • frequent **chest pain** • frequent wheezing, particularly with heartburn • unexplained weight loss • nausea or vomiting

See New Warnings

ORIGINAL STRENGTH

NOIC 10637-872-22

Pepcid[®]

AC

Famotidine Tablets 10 mg Acid Reducer

Just One Tablet!

Prevents & Relieves Heartburn Due to Acid Indigestion

90 Tablets

• stomach pain • kidney disease. **Ask a doctor or pharmacist before use if you are taking a prescription drug.** Acid reducers may interact with certain prescription drugs. **Stop use and ask a doctor if** • your heartburn continues or worsens • you need to take this product for more than 14 days. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). **Directions** • adults and children 12 years and over: • to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew. • to prevent symptoms, swallow 1 tablet with a glass of water at any time from 15 to 60 minutes before eating food or drinking beverages that cause heartburn • do not use more than 2 tablets in 24 hours • children under 12 years: ask a doctor. Famotidine (active ingredient) made in Hungary. Processed in Canada XXXXXXXX/XXXXXX

Printed by: JOHNSON & JOHNSON CONSUMER INC.
Manufactured by: Pharmacia Corporation, Kalamazoo, MI 49001 USA
© 2004 Pharmacia Corporation. All rights reserved. NDC 10637-872-22 contains 90 tablets.
EXP. DATE: 12/2005 LOT: 101



7 16837 18552 5



Do not use if foil inner seal imprinted with "SAFETY SEAL®" is broken or missing.
Active ingredient (in each tablet) **Purpose**
Famotidine 20 mg. **Acid reducers**
Uses •relieves heartburn associated with acid indigestion and sour stomach •prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages. **Warnings: Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers. **Do not use**
• if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. •with other acid reducers. **Ask a doctor before use if you have** •had heartburn over 3 months.



This may be a sign of a more serious condition. •heartburn with **high blood pressure, sweating, or dizziness** •chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders or lightheadedness •frequent **chest pain** •frequent wheezing, particularly with heartburn •unexplained weight loss •nausea or vomiting •stomach pain •kidney disease. **Ask a doctor or pharmacist before use if you are** taking a prescription drug. Acid reducers may interact with certain prescription drugs. **Stop use and ask a doctor if** •your heartburn continues or worsens •you need to take this product for more than 14 days. **If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away: (1 800 222-1222). **Directions** •adults and children 12 years and over: •to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew. •to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60 minutes before** eating food or drinking beverages that cause heartburn •do not use more than 2 tablets in 24 hours •children under 12 years: ask a doctor. Famotidine (active ingredient) made in Hungary. Processed in Canada. Distributed by: **JOHNSON & JOHNSON CONSUMER INC.** McNeil Consumer Healthcare Division Fort Washington, PA 19004 USA ©McNeil CPC 200X. PEPCID® and PEPCID AD® are registered trademarks of McNeil CPC



Do not use if foil seal under bottle cap printed "SAFETY SEAL®" is broken or missing.

Active ingredient (in each tablet) Famotidine 20 mg.

Purpose Acid reducer

Uses • relieves heartburn associated with acid indigestion and sour stomach • prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings **Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers

Do not use • if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. • with other acid reducers. **Ask a doctor before use if you have** • had heartburn over 3 months. This may be a sign of a more serious condition. • heartburn with **lightheadedness, sweating, or dizziness** • chest pain or shoulder pain with shortness of breath, sweating; pain spreading to arms, neck or shoulders; or lightheadedness • frequent **chest pain** • frequent wheezing, particularly with heartburn • unexplained weight loss • nausea or vomiting • stomach pain • kidney disease

See New Warnings

MAXIMUM STRENGTH

NDC 10437-833-16

Pepcid[®] AC

Famotidine Tablets 20 mg Acid Reducer

Just One Tablet!

Prevents & Relieves Heartburn Due to Acid Indigestion

50 Tablets

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs. **Stop use and ask a doctor if** • your heartburn continues or worsens • you need to take this product for more than 14 days. **If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose: get medical help or contact a Poison Control Center right away. (1-800-222-1222). **Directions** • adults and children 12 years and over: • to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew. • to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60 minutes before** eating food or drinking beverages that cause heartburn • do not use more than 2 tablets in 24 hours • children under 12 years: ask a doctor

Famotidine (active ingredient) made in Hungary, Processed in Canada

XXXXXXXXXXXX

Printed by: JANSSEN & JANSSEN, INC.
Manufactured by: Janssen Pharmaceutica, Inc.
NDC 10437-833-16
EXP. DATE: 12/2019
LOT: 10437-833-16