This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

PAA114536.FDA01

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS Title	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLETED SECTION	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Reference ID: 4416498

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda



PAA114537.FDA01

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.5 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• WARNING BOX LINE	.5 pt

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

- **AXID AR** is a non-prescription medicine that contains nizatidine, an ingredient that doctors have prescribed for years.
- **AXID AR** works by reducing the production of stomach acid that can cause heartburn.
- In clinical studies, **AXID AR** was significantly better than placebo in relieving and preventing heartburn symptoms.

Heartburn: a problem that can interfere with your lifestyle.

The stomach normally produces acid, especially following eating and drinking. Sometimes acid backing up into the esophagus can cause a burning pain and discomfort. This pain and discomfort is commonly known as heartburn.

When taken as directed, **AXID AR** relieves and/or prevents heartburn.

Tips for managing heartburn

- Avoid lying down flat or bending over soon after eating.
- Avoid eating late at night, or just before bedtime.
- Avoid certain foods or beverages more likely to cause heartburn, such as rich, spicy, fried foods; chocolate, caffeine, alcohol; even some fruits and vegetables.
- Eat slowly and do not eat big meals.
- If you are overweight, lose weight.
- If you smoke, stop or cut down.
- Elevate the head of your bed.
- Avoid wearing tight fitting clothing around your stomach.

PREVENTION
Benefit of AXID AR Compared to Placebo
Study A
Pills taken 60 minutes before eating



PREVENTION
Benefit of AXID AR Compared to Placebo
Study B
Pills taken 30 minutes before eating



PREVENTION
Benefit of AXID AR Compared to Placebo
Study C
Pills taken immediately before eating



RELIEF
Benefit of AXID AR Compared to Placebo
Study D
Pills taken after symptoms occur



Pfizer Consumer Healthcare
Madison, NJ 07940 USA

PAA114538.FDA01

BACK

AXID AR™

Nizatidine Tablets 75 mg
Acid Reducer

PRODUCT INFORMATION SHEET

Drug Facts

Active ingredient (in each tablet) — **Purpose**
Nizatidine 75 mg ..... Acid reducer

Uses
- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings
Allergy alert: Do not use if you are allergic to nizatidine or other acid reducers

Do not use
- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ heartburn with **lightheadedness, sweating or dizziness**
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
■ frequent **chest pain** ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain ■ kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
- adults and children 12 years and over:
  - to relieve symptoms, swallow 1 tablet with a full glass of water
  - to prevent symptoms, swallow 1 tablet with a full glass of water **right before eating or up to 60 minutes before** consuming food and beverages that cause you heartburn
  - do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information ■ store at 20-25°C (68-77°F) ■ protect from light ■ replace cap tightly after opening bottle
■ keep the carton and package insert. They contain important information.

Inactive ingredients colloidal silicon dioxide, corn starch, hypromellose, magnesium stearate, microcrystalline cellulose, pharmaceutical ink, polyethylene glycol, pregelatinized starch, propylene glycol, synthetic iron oxide, titanium dioxide

Questions or comments? Call **1-800-555-2943** Monday through Friday 9 AM to 5 PM EST

PLEASE SAVE THIS SHEET FOR FUTURE USE.

Bottle sealed with printed foil under cap. Do Not Use if foil imprinted "SEALED FOR YOUR PROTECTION" is open or torn.

FRONT

PAA114538.FDA01

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	N/A
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Reference ID: 4416498