


Actual Size 100%

FDA00068

CATEGORY	REQUIRED	ACTUAL IN FIL	
Drug Facts Title	>8 pt **	9 pt	
Drug Facts Continued	8 pt	8 pt	
Headings*	8 pt-bold italic*	8 pt	
Subheadings	≥6 pt	6 pt	
Body Text	≥6 pt	6 pt	
Bullets	5 pt	3.5 pt	
Leading	≥0.5 pt	6.5 pt	
Hairlines	0.5 pt rule	0.5 pt	
Box and Barlines <sup>+</sup>	>0.5 pt rule	1 pt	
Characters Per Inch:	No more than 39	≤ 39	



69000AGF Actual Size 100%

<sup>1</sup> aenihe8 bne xo	alun og 2.0<	tq C.f
senihis	eiun tiq 2.0	aq 87.0
6uibee	1q 2.0s	tq 01.0
stellu	tq C	tq C
aty Text	56 pt	tq 94.8
sgribeardu	26 pt	tq 94.8
*senibee	*oileti blod-tq 8	tq re.rr
beunitnoJ stoe7 gun	tq 8	tq re.rr
eltiT stoe7 gun	** tq 8<	tq E7.Sh
<b>KLEGOKA</b>	REQUIRED	VELOVE IN HEE





Sneezing
Runny Nose
Itchy, Watery Eyes
Itchy Throat or Nose

Drug Facts         Active ingredient (in each tablet)       Purpose         Loratadine 10 mg       Antihistamine         Uses       temporarily relieves these symptoms due to hay fever or other         upper respiratory allergies:       – itchy, watery eyes         – sneezing       – itching of the nose or throat		Drug Facts (continued)         Other information         • safety sealed: do not use if the individual blister unit imprinted with Claritin® RediTabs® is open or torn         • store between 20° to 25° C (68° to 77° F)         • use tablet immediately after opening individual blister         • complies with USP Procedure 2 for Assay and Organic Impurities and Test 2 for Disintegration		
its ingredients. Ask a doctor before use if you ha	an allergic reaction to this product or any of ave liver or kidney disease. Your doctor	Inactive ingredients anhydrous citric acid, gelatin, mannitol, mint flavor Questions or comments?		
should determine if you need a d When using this product do not directed may cause drowsiness. Stop use and ask a doctor if an a medical help right away. If pregnant or breast-feeding, as	ifferent dose. take more than directed. Taking more than allergic reaction to this product occurs. Seek sk a health professional before use. I case of overdose, get medical help or	Follow these d	-252-7484) or www.clarit lirections carefully it to push the table	
adults and children 6 years	disintegrates, with or without water   1 tablet daily; not more than 1 tablet in		GTOL	55
and over children under 6 years of age consumers with liver or kidney disease	24 hours ask a doctor ask a doctor	1. Peel back outer edge.	2. Gently push tablet out.	3. Place the tablet on tongue and close mouth. The tablet will disintegrate.

Bayer, the Bayer Cross, Claritin, the Claritin Vista and RediTabs are registered trademarks of Bayer. © 2018 Bayer. Dist. by: Bayer HealthCare LLC Whippany, NJ 07981 Product of Ireland. Bayer





This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

-----

/s/

THERESA M MICHELE 05/07/2019 12:21:34 PM