MAXIMUM STRENGTH

Pepcid® AC EZ CHEWS™

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
©McNeil Consumer Pharmaceuticals Co. 2019
Please visit our website at: www.pepcidac.com

The makers of Pepcid® AC do not manufacture store brands

7 16837 85425 8

MAXIMUM STRENGTH

Pepcid® AC

EZ CHEWS™

Can take without water!

25 Berries n Cream Flavor Chewable Tablets

Open other side of box for money saving offer

MAXIMUM STRENGTH

NDC 16837-854-25

Pepcid® AC

Famotidine Tablets 20 mg
Acid Reducer

EZ CHEWS™

Can take without water!

Just One Tablet!

Prevents & Relieves Heartburn
Due to Acid Indigestion

Berries 'n' Cream Flavor

25 Chewable Tablets

ACTUAL SIZE

Turn over for money saving offer

MAXIMUM STRENGTH

Pepcid® AC EZ CHEWS™

Do not use if carton is open or printed foil seal under bottle cap is open or torn.

- **1 tablet** relieves heartburn due to acid indigestion
- PEPCID® AC prevents heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.

The makers of Pepcid® AC do not manufacture store brands.

Drug Facts (continued)

Active ingredient (in each tablet)	Purpose
Famotidine 20 mg	Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.

Drug Facts (continued)

- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- adults and children 12 years and over:
- **do not swallow tablet whole: chew completely**
- to **relieve** symptoms, chew 1 tablet before swallowing
- to **prevent** symptoms, chew 1 tablet before swallowing at any time from **10 to 60 minutes before** eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- read the directions and warnings before use

Questions or comments?

1-800-755-4008 (toll-free) or **1-888-466-8746** (Spanish)

FPO

FPO

Fill Open Account™

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

LOT

EXP

XXXXXXXX

XXXXXXXX

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

VALERIE S PRATT
10/29/2019 01:50:32 PM