

FINAL RELEASE APPROVAL

Artwork No. 6-824480
HXXXXX

BD ChloraPrep™ Clear

Drug Facts	Purposes Antiseptic Antibiotic
Active ingredients Chlorhexidine gluconate 2% w/v Isopropyl alcohol 70% v/v	
Use For the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.	
Warnings For external use only. Flammable, keep away from fire or flame. • do not use with electrocautery procedures. Allergy alert: This product may cause a severe allergic reaction. Symptoms may include: • wheezing/difficulty breathing • shock • facial swelling • throat • rash If an allergic reaction occurs, stop use and seek medical help right away. Do not use • on patients allergic to chlorhexidine gluconate or any other ingredient in this product • for laceration puncture or in contact with the menses • on open skin wounds or as a general skin cleanser	
When using this product: Keep out of eyes, ears, nose, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor. Stop use and ask a doctor if: Irritation, sensitization, or allergic reaction occurs. There may be signs of a serious condition. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
Directions • use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. • maximal treatment area for one applicator is approximately 2.5 sq. x 2.5 in. (62 cm²) • remove applicator from package, do not touch sponge • hold the applicator with the sponge down. Flex the wings only once to activate the ampule and release the antiseptic. • wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin. • apply to the treatment area with single strokes. • dry surgical sites (e.g., abrasions or cuts): use gentle repeated back-and-forth strokes of the sponge for 30 seconds. Allow the area to air dry for approximately 30 seconds. • moist surgical sites (e.g., inguinal folds): use gentle repeated back-and-forth strokes of the sponge for 2 minutes. Allow the area to air dry for approximately 1 minute. • do not let it dry on skin. • discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.	
Other information • store between 15-30 °C (59-86 °F) • avoid freezing and excessive heat above 40 °C (104 °F)	
Inactive ingredient USP purified water	
Questions? • www.chloraprep.com • call 1-800-523-0502 M-F 8 a.m.-5 p.m. CST	

- Not made with natural rubber latex
- Do not reuse
- STERILE SOLUTION
- External Use Only
Professional Use Only



BD ChloraPrep™ Clear
1 mL Applicators
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Sterile Solution
Applicator is sterile if package is intact

NDC 054365-400-06



Store between 15-30 °C (59-86 °F)
Avoid freezing and excessive heat above 40 °C (104 °F)

0.03 fl. oz (1 mL) each
60 Applicators

BD ChloraPrep™ Clear
1 mL Applicators
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Sterile Solution
Applicator is sterile if package is intact

60 Applicators


NON-VARNISH AREA
LOT & EXP PRINTED IN LINE
DOES NOT PRINT

BARCODE PLACEHOLDER


1 1/16 12 7/8 5 9/8 12 7/8 5 19/32

37 19/32





FINAL RELEASE APPROVAL



4.75

3.8

BD ChloraPrep™ Clear REF 260480 / 260480NSB
NDC 054365-400-06

1 mL Applicator
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation

Sterile Solution
Applicator is sterile if package is intact

External Use Only
Professional Use Only

STERILE EO
STERILE SOLUTION

Drug Facts

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings
For external use only. Flammable, keep away from fire or flame.
• do not use with electrocautery procedures

Allergy alert:
This product may cause a severe allergic reaction. Symptoms may include:
• wheezing/difficulty breathing • shock • facial swelling • hives • rash
If an allergic reaction occurs, stop use and seek medical help right away.

Do not use
• on patients allergic to chlorhexidine gluconate or any other ingredient in this product
• for lumbar puncture or in contact with the meninges
• on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

FRONT

BD ChloraPrep™ Clear
1 mL Applicator
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

Drug Facts (continued)

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm²)
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wings **only once** to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for approximately 30 seconds. Allow the area to air dry for approximately 30 seconds.
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for approximately 2 minutes. Allow the area to air dry for approximately 1 minute.
- do not blot or wipe away
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15–30 °C (59–86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)


Inactive ingredient USP purified water

Questions? • www.chloraprep.com • call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

Becton, Dickinson and Company, El Paso, TX 79912 · Artwork No. 6-775480 · HXXXXXX
Made in the USA of US and imported parts · www.bd.com


REAR

FINAL RELEASE APPROVAL



(b) (4)

FINAL RELEASE APPROVAL



(b) (4)



Artwork No. 6-824299
WXXXXXX

BD ChloraPrep™ FREPP™ Clear

1.5 mL Applicators

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation


Sterile Solution
Applicator is sterile if package is intact

Store between 15-30 °C (59-86 °F)
Avoid freezing and excessive heat
above 40 °C (104 °F)

0.05 fl. oz (1.5 mL) each
20 Applicators

Clear

NDC 054365-400-02



REF 260299

BD ChloraPrep™ FREPP™ Clear

1.5 mL Applicators

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation

Sterile Solution
Applicator is sterile if package is intact

20 Applicators

NON-VARNISH AREA
LOT & EXP PRINTED IN-LINE
DOES NOT PRINT

Clear

Drug Facts

Active ingredients
Chlorhexidine gluconate 2% w/v Antiseptic
Isopropyl alcohol 70% v/v Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings
For external use only. Flammable, keep away from fire or flame.
• do not use with electrocautery procedures

Allergy alert: This product may cause a severe allergic reaction. Symptoms may include:
• wheezing/difficulty breathing • shock • facial swelling • hives • rash
If an allergic reaction occurs, stop use and seek medical help right away.

Do not use
• on patients allergic to chlorhexidine gluconate or any other ingredient in this product
• for lumbar puncture or in contact with the meninges
• on open skin wounds or as a general skin cleanser

When using this product, keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Other information
• store between 15-30 °C (59-86 °F) • avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredient USP purified water

Questions?
• www.chloraprep.com • call 1-800-523-0502 (M-F 8 a.m.-5 p.m. CST)

External Use Only
Professional Use Only

Becton, Dickinson and Company, El Paso, TX 79912 www.bd.com Made in the USA of US and imported parts
© 2017 BD, BD, the BD Logo and BD ChloraPrep are trademarks of Becton, Dickinson and Company.

Drug Facts (continued)

Directions
• use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
• maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm²)

• remove applicator from package; do not touch sponge
• hold the applicator horizontally with the sponge down. Pinch wings **only once** to activate the ampule and release the antiseptic.

• wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
• completely wet the treatment area with antiseptic

• **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds. Allow the area to air dry for approximately 30 seconds.
• **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes. Allow the area to air dry for approximately 1 minute.

• do not blot or wipe away
• discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information
• store between 15-30 °C (59-86 °F) • avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredient USP purified water

Questions?
• www.chloraprep.com • call 1-800-523-0502 (M-F 8 a.m.-5 p.m. CST)

External Use Only
Professional Use Only



2.484

0.625

8.500

3.500

8.500

3.469

24.594

0.969


2.219

3.500


1.734

10.422






FINAL RELEASE APPROVAL





FINAL RELEASE APPROVAL



2.5

**BD ChloraPrep™
FREPP™ Clear**
1.5 mL Applicator
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Sterile Solution
Applicator is sterile if package is intact

Drug Facts

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use For the preparation of the patient's skin prior to surgery.
Helps to reduce bacteria that potentially can cause skin infection.

Warnings
For external use only.
Flammable, keep away from fire or flame.
• do not use with electrocautery procedures

Allergy alert:
This product may cause a severe allergic reaction. Symptoms may include:
• wheezing/difficulty breathing • shock • facial swelling
• hives • rash
If an allergic reaction occurs, stop use and seek medical help right away.

Do not use
• on patients allergic to chlorhexidine gluconate or any other ingredient in this product
• for lumbar puncture or in contact with the meninges
• on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions
• use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
• maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm²)
• remove applicator from package; do not touch sponge
• hold the applicator horizontally with the sponge down. Pinch wings **only once** to activate the ampule and release the antiseptic.
• wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
• completely wet the treatment area with antiseptic
• **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for approximately 30 seconds. Allow the area to air dry for approximately 30 seconds.
• **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for approximately 2 minutes. Allow the area to air dry for approximately 1 minute.
• do not blot or wipe away
• discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information
• store between 15-30 °C (59-86 °F)
• avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredient USP purified water

Questions? • www.chloraprep.com
• call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

See other side for additional information



7.5


FRONT


REF 260299NSB / 260599NSB
NDC 054365-400-02

**BD ChloraPrep™
FREPP™ Clear**
1.5 mL Applicator
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Sterile Solution
Applicator is sterile if package is intact

See Drug Facts for Full Labeling Information

  Not made with natural rubber latex

 Do not reuse


 **STERILE SOLUTION**


External Use Only
Professional Use Only


Becton, Dickinson and Company,
El Paso, TX 79912
Artwork No. 6-775403F
SXXXXXX
Made in the USA of US and imported parts
www.bd.com
© 2017 BD. BD, the BD Logo and all other trademarks are property of Becton, Dickinson and Company.

REAR


FINAL RELEASE APPROVAL







FINAL RELEASE APPROVAL



4.75

3.8

⚠ WARNING
FLAMMABLE

Keep away from fire or flame. To reduce risk of fire, **PREP CAREFULLY:**

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (**minimum of 3 minutes on hairless skin; up to 1 hour in hair**)
- do not allow solution to pool
- remove wet materials from prep area

REF 260400 / 260400NS / 260500NS NDC 054365-400-01
REF 260415 / 260415NS / 260415NSB NDC 054365-400-11

BD ChloraPrep™ Clear
BD ChloraPrep™ Hi-Lite Orange™
3 mL Applicator
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Sterile Solution
Applicator is sterile if package is intact

STERILE SOLUTION External Use Only Professional Use Only

Drug Facts

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings
For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, **PREP CAREFULLY:**

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (**minimum of 3 minutes on hairless skin; up to 1 hour in hair**)
- do not allow solution to pool
- remove wet materials from prep area

Allergy alert:
This product may cause a severe allergic reaction. Symptoms may include:
• wheezing/difficulty breathing • shock • facial swelling • hives • rash
If an allergic reaction occurs, stop use and seek medical help right away.

FRONT

BD ChloraPrep™ Clear / BD ChloraPrep™ Hi-Lite Orange™
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

Drug Facts (continued)

Do not use

- on patients allergic to chlorhexidine gluconate or any other ingredient in this product
- for lumbar puncture or in contact with the meninges • on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 4 in. x 5 in. (130 cm²)
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wings **only once** to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes of the sponge for approximately 30 seconds. Allow solution to air dry for approximately 30 seconds. Do not blot or wipe away.
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes of the sponge for approximately 2 minutes. Allow solution to air dry for approximately 1 minute. Do not blot or wipe away.
- **if using an ignition source, allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair).** Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information • store between 15–30 °C (59–86 °F) • avoid freezing and excessive heat above 40 °C (104 °F)
• the tint will slowly fade from the skin. Soap and water, or alcohol may be used to remove the tint if desired.


Inactive ingredients FD&C yellow #6 dye (Hi-Lite Orange applicators only), USP purified water

Questions? • www.chloraprep.com • call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

Becton, Dickinson and Co., El Paso, TX 79912 · Artwork No. 6-775400 · LXXXXXX · Made in the USA of US and imported parts · www.bd.com

REAR

FINAL RELEASE APPROVAL



This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

THERESA M MICHELE
04/03/2019 12:00:00 AM