

NDC 16837-891-25
See New Warnings

DUAL ACTION
Pepcid Complete

Famotidine 10 mg—Acid reducer
Calcium carbonate 800 mg—Antacid
Magnesium hydroxide 165 mg—Antacid

Just One Tablet!
Relieves Heartburn Due to Acid Indigestion
Cool Mint Flavor

25 Chewable Tablets

actual size


For Complete Directions and Warnings

Do not use if foil seal under bottle cap printed "SAFETY SEAL®" is open or torn.

Active ingredients (in each chewable tablet)	Purposes
Famotidine 10 mg.....	Acid reducer
Calcium carbonate 800 mg.....	Antacid
Magnesium hydroxide 165 mg.....	Antacid

Use relieves heartburn associated with acid indigestion and sour stomach

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA
©McNeil Consumer Pharmaceuticals Co. 20XX
PEPCID and PEPCID COMPLETE are registered trademarks of McNeil CPC
US Patent Nos. 5,989,588 and 6,814,978



EXP
XXXXXXX

Drug Facts (continued)

Other information

- each tablet contains: calcium 320 mg, magnesium 70 mg
- read the directions and warnings before use
- read the bottle. It contains important information.
- store at 20°-25° C (68°-77° F)
- protect from moisture
- do not use if foil seal under bottle cap printed "SAFETY SEAL®" is open or torn

Inactive ingredients: cellulose acetate, corn starch, crospovidone, D&C yellow no. 10 aluminum lake, dextrose excipient, FD&C blue no. 1 aluminum lake, flavors, gum arabic, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, maltodextrin, mineral oil, sucralose

Questions or comments?
call 1-800-755-4008 (toll-free) or 215-273-8755 (collect)

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking



Shown actual size

Pepcid Complete

Acid Reducer + Antacid

Famotidine 10 mg per tablet
Calcium carbonate 800 mg per tablet
Magnesium hydroxide 165 mg per tablet

25 Chewable Tablets

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**
McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA
©McNeil Consumer Pharmaceuticals Co. 20XX
PEPCID and PEPCID COMPLETE are registered trademarks of McNeil CPC
US Patent Nos. 5,989,588 and 6,814,978

<p>Drug Facts (continued)</p> <p>Warnings Allergy alert: Do not use if you are allergic to famotidine or other acid reducers</p> <p>Do not use</p> <ul style="list-style-type: none"> • if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. • See your doctor. • with other acid reducers <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> • had heartburn over 3 months. This may be a sign of a more serious condition. • heartburn with lightheadedness, sweating, or dizziness • chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness <p>Frequent chest pain</p> <ul style="list-style-type: none"> • frequent wheezing, particularly with heartburn • unexplained weight loss • stomach pain 	<p>Drug Facts (continued)</p> <p>Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids and acid reducers may interact with certain prescription drugs.</p> <p>Step use and ask a doctor if</p> <ul style="list-style-type: none"> • your heartburn continues or worsens • you need to take this product for more than 14 days <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)</p> <p>Directions</p> <ul style="list-style-type: none"> • adults and children 12 years and over: <ul style="list-style-type: none"> • do not swallow tablet whole; chew completely • to relieve symptoms, chew 1 tablet before swallowing • do not use more than 2 chewable tablets in 24 hours • children under 12 years: ask a doctor
---	--

Labeling Format Information: Fonts: Univers LT Std-57 Cond, 67 Cond Bold, 67 Bold Cond Oblique

Drug Facts: 8.50 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 85%
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: 2.50 pt	
Subheaders: 6.00 pt	Drug Facts (continued): 8.00 pt	Hairlines: 0.50 pt	Average Tracking: 0



EXP

XXXXXXX

Drug Facts (continued)

Other information

- each tablet contains: calcium 320 mg, magnesium 70 mg
- read the directions and warnings before use
- read the bottle. It contains important information.
- store at 20°-25°C (68°-77°F) • protect from moisture
- do not use if foil seal under bottle cap printed "SAFETY SEAL®" is open or torn

Inactive ingredients cellulose acetate, corn starch, crospovidone, D&C yellow no. 10 aluminum lake, dextrose excipient, FD&C blue no. 1 aluminum lake, flavors, gum arabic, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, maltodextrin, mineral oil, sucralose

Questions or comments? call 1-800-755-4008 (toll-free) or 215-273-8755 (collect)

Pepcid Complete
Acid Reducer + Antacid
 Famotidine 10 mg per tablet
 Calcium carbonate 800 mg per tablet
 Magnesium hydroxide 165 mg per tablet
50 Chewable Tablets



Shown actual size

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**
 McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA
 ©McNeil Consumer Pharmaceuticals Co. 20XX
 PEPCID® and PEPCID COMPLETE® are registered trademarks of McNeil CPC
 US Patent Nos. 5,989,588 and 6,814,978

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

See New Warnings NDC 16837-891-50

DUAL ACTION
Pepcid Complete
 Famotidine 10 mg—Acid reducer
 Calcium carbonate 800 mg—Antacid
 Magnesium hydroxide 165 mg—Antacid

Just One Tablet!
 Relieves Heartburn Due to Acid Indigestion
 Cool Mint Flavor

50 Chewable Tablets

For Complete Directions and Warnings
 Do not use if foil seal under bottle cap printed "SAFETY SEAL®" is open or torn.

Drug Facts

Active ingredients (in each chewable tablet)

Active ingredients	Purposes
Famotidine 10 mg	Acid reducer
Calcium carbonate 800 mg	Antacid
Magnesium hydroxide 165 mg	Antacid

Use relieves heartburn associated with acid indigestion and sour stomach

Warnings
Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Distributed by:
 JOHNSON & JOHNSON CONSUMER INC.
 McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA
 ©McNeil Consumer Pharmaceuticals Co. 20XX
 PEPCID® and PEPCID COMPLETE® are registered trademarks of
 McNeil CPC
 US Patent Nos. 5,989,588 and 6,814,978

Drug Facts (continued)

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids and acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults and children 12 years and over:
 - do not swallow tablet whole: **chew completely**
 - to relieve symptoms, **chew 1 tablet** before swallowing
 - do not use more than 2 chewable tablets in 24 hours
- children under 12 years: ask a doctor

Labeling Format Information: Fonts: Univers LT Std-57 Cond, 67 Cond Bold, 67 Bold Cond Oblique

Drug Facts: 8.50 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 85/100%
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: 2.50 pt	
Subheaders: 6.00 pt	Drug Facts (continued): 8.00 pt	Hairlines: 0.50 pt	Average Tracking: 0



Drug Facts (continued)
Other information
<ul style="list-style-type: none"> each tablet contains: calcium 320 mg, magnesium 70 mg read the directions and warnings before use read the bottle. It contains important information. store at 20°-25° C (68°-77° F) protect from moisture do not use if foil seal under bottle cap printed "SAFETY SEAL®" is open or torn
Inactive ingredients: cellulose acetate, corn starch, crospovidone, dextrose excipient, FD&C blue no. 1 aluminum lake, flavors, gum arabic, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, maltodextrin, mineral oil, sucralose, yellow iron oxide
Questions or comments? call 1-800-755-4008 (toll-free) or 215-273-8755 (collect)

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Pepcid Complete
Acid Reducer + Antacid
Famotidine 10 mg per tablet
Calcium carbonate 800 mg per tablet
Magnesium hydroxide 165 mg per tablet
25 Chewable Tablets

Shown actual size

Distributed by **JOHNSON & JOHNSON CONSUMER INC.**
McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA
©McNeil Consumer Pharmaceuticals Co. 20XX
PEPCID® and PEPCID COMPLETE® are registered trademarks of McNeil CPC
US Patent Nos. 5,989,588 and 6,814,978

NDC 16837-891-65
See New Warnings

DUAL ACTION
Pepcid Complete

Famotidine 10 mg—Acid reducer
Calcium carbonate 800 mg—Antacid
Magnesium hydroxide 165 mg—Antacid

Just One Tablet!
Relieves Heartburn Due to Acid Indigestion
Cool Mint Flavor

25 Chewable Tablets

For Complete Directions and Warnings

Do not use if foil seal under bottle cap printed "SAFETY SEAL®" is open or torn.

Drug Facts	
Active ingredients (in each chewable tablet)	Purposes
Famotidine 10 mg.....	Acid reducer
Calcium carbonate 800 mg.....	Antacid
Magnesium hydroxide 165 mg.....	Antacid
Use	relieves heartburn associated with acid indigestion and sour stomach

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA
©McNeil Consumer Pharmaceuticals Co. 20XX
PEPCID® and PEPCID COMPLETE® are registered trademarks of McNeil CPC
US Patent Nos. 5,989,588 and 6,814,978

Drug Facts (continued)	Drug Facts (continued)
Warnings Allergy alert: Do not use if you are allergic to famotidine or other acid reducers	Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids and acid reducers may interact with certain prescription drugs.
Do not use • if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. • with other acid reducers	Step use and ask a doctor if • your heartburn continues or worsens • you need to take this product for more than 14 days
Ask a doctor before use if you have • had heartburn over 3 months. This may be a sign of a more serious condition. • heartburn with lightheadedness, sweating, or dizziness • chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness	If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)
Directions • frequent chest pain • frequent wheezing, particularly with heartburn • unexplained weight loss • stomach pain	Directions • adults and children 12 years and over: • do not swallow tablet whole; chew completely • to relieve symptoms, chew 1 tablet before swallowing • do not use more than 2 chewable tablets in 24 hours • children under 12 years: ask a doctor

Labeling Format Information: Fonts: Univers LT Std-57 Cond, 67 Cond Bold, 67 Bold Cond Oblique

Drug Facts: 8.50 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 85%
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: 2.50 pt	
Subheaders: 6.00 pt	Drug Facts (continued): 8.00 pt	Hairlines: 0.50 pt	Average Tracking: 0

(b) (4)



EXP
XXXXXXX

Drug Facts (continued)
Other information
 • each tablet contains: calcium 320 mg, magnesium 70 mg
 • read the directions and warnings before use
 • read the bottle. It contains important information.
 • store at 20°-25°C (68°-77°F) • protect from moisture
 • do not use if foil seal under bottle cap printed "SAFETY SEAL®" is open or torn
Inactive ingredients cellulose acetate, corn starch, crospovidone, dextrose excipient, FD&C blue no. 1 aluminum lake, flavors, gum arabic, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, maltodextrin, mineral oil, sucralose, yellow iron oxide
Questions or comments? call 1-800-755-4008 (toll-free) or 215-273-8755 (collect)

Pepcid Complete
Acid Reducer + Antacid
 Famotidine 10 mg per tablet
 Calcium carbonate 800 mg per tablet
 Magnesium hydroxide 165 mg per tablet
50 Chewable Tablets



Shown actual size

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**
 McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA
 ©McNeil Consumer Pharmaceuticals Co. 20XX
 PEPCID® and PEPCID COMPLETE® are registered trademarks of McNeil CPC
 US Patent Nos. 5,989,588 and 6,814,978

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

See New Warnings NDC 16837-891-66

DUAL ACTION
Pepcid Complete
 Famotidine 10 mg—Acid reducer
 Calcium carbonate 800 mg—Antacid
 Magnesium hydroxide 165 mg—Antacid

Just One Tablet!
 Relieves Heartburn Due to Acid Indigestion
 Cool Mint Flavor

50 Chewable Tablets

For Complete Directions and Warnings
 Do not use if foil seal under bottle cap printed "SAFETY SEAL®" is open or torn.

Drug Facts
Active ingredients (in each chewable tablet)
 Famotidine 10 mg..... Acid reducer
 Calcium carbonate 800 mg..... Antacid
 Magnesium hydroxide 165 mg..... Antacid
Purposes
 • relieves heartburn associated with acid indigestion and sour stomach
Use
 • relieves heartburn associated with acid indigestion and sour stomach
Warnings
Allergy alert: Do not use if you are allergic to famotidine or other acid reducers
Do not use
 • if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
 • with other acid reducers

Distributed by:
 JOHNSON & JOHNSON CONSUMER INC.
 McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA
 ©McNeil Consumer Pharmaceuticals Co. 20XX
 PEPCID® and PEPCID COMPLETE® are registered trademarks of McNeil CPC
 US Patent Nos. 5,989,588 and 6,814,978

Drug Facts (continued)
Ask a doctor before use if you have
 • had heartburn over 3 months. This may be a sign of a more serious condition.
 • heartburn with **lightheadedness, sweating, or dizziness**
 • chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness
 • frequent **chest pain**
 • frequent wheezing, particularly with heartburn
 • unexplained weight loss
 • nausea or vomiting
 • stomach pain
 • kidney disease
Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids and acid reducers may interact with certain prescription drugs.
Stop use and ask a doctor if
 • your heartburn continues or worsens
 • you need to take this product for more than 14 days
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions
 • adults and children 12 years and over:
 • do not swallow tablet whole: **chew completely**
 • to relieve symptoms, **chew 1 tablet** before swallowing
 • do not use more than 2 chewable tablets in 24 hours
 • children under 12 years: ask a doctor

Labeling Format Information: Fonts: Univers LT Std-57 Cond, 67 Cond Bold, 67 Bold Cond Oblique

Drug Facts: 8.50 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 85/100%
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: 2.50 pt	
Subheaders: 6.00 pt	Drug Facts (continued): 8.00 pt	Hairlines: 0.50 pt	Average Tracking: 0

NDC 16837-298-25
See New Warnings

DUAL ACTION
Pepcid Complete

Famotidine 10 mg—Acid reducer
Calcium carbonate 800 mg—Antacid
Magnesium hydroxide 165 mg—Antacid

Just One Tablet!
Relieves Heartburn Due to Acid Indigestion
Berry Flavor

25 Chewable Tablets

actual size

For Complete Directions and Warnings

Do not use if foil seal under bottle cap printed "SAFETY SEAL®" is open or torn.

Drug Facts	
Active ingredients (in each chewable tablet)	Purposes
Famotidine 10 mg	Acid reducer
Calcium carbonate 800 mg	Antacid
Magnesium hydroxide 165 mg	Antacid

Use relieves heartburn associated with acid indigestion and sour stomach

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
©McNeil Consumer Pharmaceuticals Co., 2018
PEPCID® and PEPCID COMPLETE® are registered trademarks of McNeil-PPC
US Patent Nos. 5,991,588 and 6,814,978



EXP
XXXXXXX

Drug Facts (continued)

Other information

- each tablet contains: calcium 320 mg, magnesium 70 mg
- read the directions and warnings before use
- read the bottle. It contains important information.
- store at 20°-25° C (68°-77° F)
- protect from moisture
- do not use if foil seal under bottle cap printed "SAFETY SEAL®" is open or torn

Inactive ingredients cellulose acetate, corn starch, croscopolidone, D&C red no. 7 calcium lake, dextrose excipient, FD&C blue no. 1 aluminum lake, FD&C red no. 40 aluminum lake, flavors, gum arabic, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, maltodextrin, mineral oil, modified starch, sucralose

Questions or comments?
call 1-800-755-4088 (toll-free) or 215-273-8755 (collect)

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Pepcid Complete
Acid Reducer + Antacid
Famotidine 10 mg per tablet
Calcium carbonate 800 mg per tablet
Magnesium hydroxide 165 mg per tablet
25 Chewable Tablets

Shown actual size

Developed by: JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA
©McNeil Consumer Pharmaceuticals Co., 2018
PEPCID® and PEPCID COMPLETE® are registered trademarks of McNeil-PPC
US Patent Nos. 5,991,588 and 6,814,978



<p>Drug Facts (continued)</p> <p>Warnings Allergy alert: Do not use if you are allergic to famotidine or other acid reducers</p> <p>Do not use</p> <ul style="list-style-type: none"> if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. with other acid reducers <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> had heartburn over 3 months. This may be a sign of a more serious condition. heartburn with lightheadedness, sweating, or dizziness chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness frequent chest pain frequent wheezing, particularly with heartburn unexplained weight loss stomach pain nausea or vomiting kidney disease 	<p>Drug Facts (continued)</p> <p>Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids and acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> your heartburn continues or worsens you need to take this product for more than 14 days <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)</p> <p>Directions</p> <ul style="list-style-type: none"> adults and children 12 years and over: <ul style="list-style-type: none"> do not swallow tablet whole: chew completely to relieve symptoms, chew 1 tablet before swallowing do not use more than 2 chewable tablets in 24 hours children under 12 years: ask a doctor
--	---

Labeling Format Information: Fonts: Univers LT Std-57 Cond, 67 Bold Cond, 67 Bold Cond Oblique

Drug Facts: 8.50 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 85%
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: 2.50 pt	
Subheaders: 6.00 pt	Drug Facts (continued): 8.00 pt	Hairlines: 0.50 pt	Average Tracking: 0



7
10837291501
3

EXP

XXXXXXXXX

Drug Facts (continued)

Other information

- each tablet contains: calcium 320 mg, magnesium 70 mg
- read the directions and warnings before use
- read the bottle. It contains important information.
- store at 20°-25°C (68°-77°F) • protect from moisture
- **do not use if foil seal under bottle cap printed "SAFETY SEAL®" is open or torn**

Inactive ingredients cellulose acetate, corn starch, croscopovidone, D&C red no. 7 calcium lake, dextrose excipient, FD&C blue no. 1 aluminum lake, FD&C red no. 40 aluminum lake, flavors, gum arabic, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, maltodextrin, mineral oil, modified starch, sucralose

Questions or comments? call 1-800-755-4008 (toll-free) or 215-273-8755 (collect)

Pepcid Complete

Acid Reducer + Antacid

Famotidine 10 mg per tablet
Calcium carbonate 800 mg per tablet
Magnesium hydroxide 165 mg per tablet

50 Chewable Tablets

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**
McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA
©McNeil Consumer Pharmaceuticals Co. 20XX
PEPCID® and PEPCID COMPLETE® are registered trademarks of McNeil CPC
US Patent Nos. 5,989,588 and 6,814,978



Shown actual size

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

See New Warnings **DUAL ACTION** NDC 16837-288-50

Do not use if foil seal under bottle cap printed "SAFETY SEAL®" is open or torn.

Pepcid Complete

Famotidine 10 mg—Acid reducer
Calcium carbonate 800 mg—Antacid
Magnesium hydroxide 165 mg—Antacid

Just One Tablet!
Relieves Heartburn Due to Acid Indigestion
Berry Flavor

50 Chewable Tablets

Drug Facts

Active ingredients (in each chewable tablet)	Purposes
Famotidine 10 mg.....	Acid reducer
Calcium carbonate 800 mg.....	Antacid
Magnesium hydroxide 165 mg.....	Antacid

Use relieves heartburn associated with acid indigestion and sour stomach

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**
McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA
©McNeil Consumer Pharmaceuticals Co. 20XX
PEPCID® and PEPCID COMPLETE® are registered trademarks of McNeil CPC
US Patent Nos. 5,989,588 and 6,814,978

Drug Facts (continued)

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids and acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults and children 12 years and over:
 - **do not swallow tablet whole; chew completely**
 - to relieve symptoms, **chew 1** tablet before swallowing
 - do not use more than 2 chewable tablets in 24 hours
- children under 12 years: ask a doctor

Labeling Format Information: Fonts: Univers LT Std-57 Cond, 67 Bold Cond, 67 Bold Cond Oblique

Drug Facts: 8.50 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 85/100%
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: 2.50 pt	
Subheaders: 6.00 pt	Drug Facts (continued): 8.00 pt	Hairlines: 0.50 pt	Average Tracking: 0



7
1603729112
1
EXP
XXXXXXXX

Drug Facts (continued)

Other information

- each tablet contains: calcium 320 mg, magnesium 70 mg
- read the directions and warnings before use
- read the bottle. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture
- do not use if foil seal under bottle cap printed "SAFETY SEAL®" is open or torn

Inactive ingredients cellulose acetate, corn starch, crospovidone, D&C red no. 7 calcium lake, dextrose excipient, FD&C blue no. 1 aluminum lake, FD&C red no. 40 aluminum lake, flavors, gum arabic, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, maltodextrin, mineral oil, modified starch, sucralose

Questions or comments? call 1-800-755-4008 (toll-free) or 215-273-8755 (collect)

Pepcid Complete
Acid Reducer + Antacid

Famotidine 10 mg per tablet
Calcium carbonate 800 mg per tablet
Magnesium hydroxide 165 mg per tablet

100 Chewable Tablets

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**
McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA
©McNeil Consumer Pharmaceuticals Co. 20XX
PEPCID® and PEPCID COMPLETE® are registered trademarks of McNeil CPC
US Patent Nos. 5,999,588 and 6,814,978



Shown actual size

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

See New Warnings **DUAL ACTION** NDC 16837-298-12

Pepcid Complete

Famotidine 10 mg—Acid reducer
Calcium carbonate 800 mg—Antacid
Magnesium hydroxide 165 mg—Antacid

Just One Tablet!
Relieves Heartburn Due to Acid Indigestion
Berry Flavor

100 Chewable Tablets

For Complete Directions and Warnings
Do not use if foil seal under bottle cap printed "SAFETY SEAL®" is open or torn.

Drug Facts

Active ingredients (in each chewable tablet)	Purposes
Famotidine 10 mg.....	Acid reducer
Calcium carbonate 800 mg.....	Antacid
Magnesium hydroxide 165 mg.....	Antacid

Use relieves heartburn associated with acid indigestion and sour stomach

Warnings
Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
©McNeil Consumer Pharmaceuticals Co. 20XX
PEPCID® and PEPCID COMPLETE® are registered trademarks of McNeil CPC
US Patent Nos. 5,999,588 and 6,814,978

Drug Facts (continued)

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids and acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults and children 12 years and over:
 - do not swallow tablet whole; **chew completely**
 - to relieve symptoms, **chew** 1 tablet before swallowing
 - do not use more than 2 chewable tablets in 24 hours
- children under 12 years: ask a doctor

Labeling Format Information: Fonts: Univers LT Std-57 Cond, 67 Bold Cond, 67 Bold Cond Oblique

Drug Facts: 8.50 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 95/100%
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: 2.50 pt	
Subheaders: 6.00 pt	Drug Facts (continued): 8.00 pt	Hairlines: 0.50 pt	Average Tracking: 0



(b) (4)



EXP
XXXXXXX

<p>Drug Facts (continued)</p> <p>Other information</p> <ul style="list-style-type: none"> • each tablet contains: calcium 320 mg, magnesium 70 mg • contains FD&C yellow no. 5 (tartrazine) as a color additive • read the directions and warnings before use • read the bottle. It contains important information. • store at 20°-25° C (68°-77°F) • protect from moisture • do not use if foil seal under bottle cap printed "SAFETY SEAL®" is open or torn <p>Inactive ingredients cellulose acetate, corn starch, corn syrup solids, croscopolidone, dextrose excipient, FD&C yellow no. 5 aluminum lake (tartrazine), FD&C yellow no. 6 aluminum lake, flavors, gum arabic, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, maltodextrin, mineral oil, sucralose, triacetin</p> <p>Questions or comments? call 1-800-755-4008 (toll-free) or 215-273-8755 (collect)</p>	<p>Tips for Managing Heartburn</p> <ul style="list-style-type: none"> • Do not lie flat or bend over after eating • Do not wear tight-fitting clothing around the stomach • Do not eat before bedtime • Raise the head of your bed • Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables • Eat slowly and avoid big meals • If overweight, lose weight • Quit smoking <p style="text-align: center;">Pepcid Complete</p> <p style="text-align: center;">Acid Reducer + Antacid</p> <p style="text-align: center;">Famotidine 10 mg per tablet Calcium carbonate 800 mg per tablet Magnesium hydroxide 165 mg per tablet</p> <p style="text-align: center;">25 Chewable Tablets</p> <p style="text-align: center; font-size: small;">Distributed by: JOHNSON & JOHNSON CONSUMER INC. McNeil Consumer Pharmaceuticals Co. 20XX PEPCID® and PEPCID COMPLETE® are registered trademarks of McNeil CPC US Patent Nos. 5,989,588 and 6,814,978</p>	 <p style="font-size: x-small;">Shown actual size</p>
---	--	--





For Complete Directions and Warnings

Do not use if foil seal under bottle cap printed "SAFETY SEAL®" is open or torn.

Drug Facts	Purposes
Active ingredients (in each chewable tablet)	
Famotidine 10 mg.....	Acid reducer
Calcium carbonate 800 mg.....	Antacid
Magnesium hydroxide 165 mg.....	Antacid

Use relieves heartburn associated with acid indigestion and sour stomach

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA
©McNeil Consumer Pharmaceuticals Co. 20XX
PEPCID® and PEPCID COMPLETE® are registered trademarks of McNeil CPC
US Patent Nos. 5,989,588 and 6,814,978

<p>Drug Facts (continued)</p> <p>Warnings</p> <p>Allergy alert: Do not use if you are allergic to famotidine or other acid reducers</p> <p>Do not use</p> <ul style="list-style-type: none"> • if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. • with other acid reducers <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> • had heartburn over 3 months. This may be a sign of a more serious condition; • heartburn with lightheadedness, sweating, or dizziness • chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness • frequent chest pain • frequent wheezing, particularly with heartburn • unexplained weight loss • stomach pain • nausea or vomiting • kidney disease 	<p>Drug Facts (continued)</p> <p>Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids and acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> • your heartburn continues or worsens • you need to take this product for more than 14 days <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)</p> <p>Directions</p> <ul style="list-style-type: none"> • adults and children 12 years and over: <ul style="list-style-type: none"> • do not swallow tablet whole; chew completely • do not use more than 2 chewable tablets in 24 hours • children under 12 years: ask a doctor
---	---

Labeling Format Information: Fonts: Univers LT Std-57 Cond, 67 Bold Cond, 67 Bold Cond Oblique			
Drug Facts: 8.50 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 85%
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: 2.50 pt	
Subheaders: 6.00 pt	Drug Facts (continued): 8.00 pt	Hairlines: 0.50 pt	Average Tracking: 0



EXP
XXXXXXX

Drug Facts (continued)

Other information

- each tablet contains: calcium 320 mg, magnesium 70 mg
- contains FD&C yellow no. 5 (tartrazine) as a color additive
- read the directions and warnings before use
- store at 20°-25° C (68°-77°F)
- do not use if foil seal under bottle cap printed "SAFETY SEAL®" is open or torn

Inactive ingredients cellulose acetate, corn starch, corn syrup solids, crospovidone, dextrose excipient, FD&C yellow no. 5 aluminum lake (tartrazine), FD&C yellow no. 6 aluminum lake, flavors, gum arabic, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, maltodextrin, mineral oil, sucralose, triacetin

Questions or comments? call 1-800-755-4008 (toll-free) or 215-273-8755 (collect)

Pepcid Complete

Acid Reducer + Antacid
Famotidine 10 mg per tablet
Calcium carbonate 800 mg per tablet
Magnesium hydroxide 165 mg per tablet



Shown actual size

50 Chewable Tablets
Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**
McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA
©McNeil Consumer Pharmaceuticals Co. 20XX
PEPCID® and PEPCID COMPLETE® are registered trademarks of McNeil CPC
US Patent Nos. 5,989,588 and 6,814,978

- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

See New Warnings NDC 16837-246-50

DUAL ACTION

Pepcid Complete

Famotidine 10 mg—Acid reducer
Calcium carbonate 800 mg—Antacid
Magnesium hydroxide 165 mg—Antacid

Just One Tablet!
Relieves Heartburn Due to Acid Indigestion
Tropical Fruit Flavor

50 Chewable Tablets

For Complete Directions and Warnings
Do not use if foil seal under bottle cap printed "SAFETY SEAL®" is open or torn.

Drug Facts

Active ingredients (in each chewable tablet)	Purposes
Famotidine 10 mg	Acid reducer
Calcium carbonate 800 mg	Antacid
Magnesium hydroxide 165 mg	Antacid

Use relieves heartburn associated with acid indigestion and sour stomach

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**
McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA
©McNeil Consumer Pharmaceuticals Co. 20XX
PEPCID® and PEPCID COMPLETE® are registered trademarks of McNeil CPC
US Patent Nos. 5,989,588 and 6,814,978

Drug Facts (continued)

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids and acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults and children 12 years and over:
 - do not swallow tablet whole; chew completely
 - to relieve symptoms, **chew** 1 tablet before swallowing
 - do not use more than 2 chewable tablets in 24 hours
- children under 12 years: ask a doctor

Labeling Format Information: Fonts: Univers LT Std-57 Cond, 67 Bold Cond, 67 Bold Cond Oblique

Drug Facts: 8.50 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 85/100%
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: 2.50 pt	
Subheaders: 6.00 pt	Drug Facts (continued): 8.00 pt	Hairlines: 0.50 pt	Average Tracking: 0

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

KAREN M MAHONEY on behalf of VALERIE S PRATT
04/09/2019 12:00:00 AM