

This label may not be the latest approved by FDA.  
For current labeling information, please visit <https://www.fda.gov/drugsatfda>



Do not use if pouch is open or torn.

**Active ingredients (in each chewable tablet)**

Famotidine 10 mg.....Acid reducer

Calcium carbonate 800 mg.....Antacid

Magnesium hydroxide 165 mg.....Antacid

**Use:** relieves heartburn associated with acid indigestion and sour stomach

**Warnings: Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers

To Open: While Folded  
on Line, Tear At Slit

**Do not use** • if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. • with other acid reducers. **Ask a doctor before use if you have** • had heartburn over 3 months. This may be a sign of a more serious condition. • heartburn with **lightheadedness, sweating, or dizziness** • chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness • frequent **chest pain** • frequent wheezing, particularly with heartburn • unexplained weight loss • nausea or vomiting • stomach pain • kidney disease. **Ask a doctor or pharmacist before use if you are** taking a prescription drug. Antacids and acid reducers may interact with certain prescription drugs. **Stop use and ask a doctor if** • your heartburn continues or worsens • you need to take this product for more than 14 days. **If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) **Directions** • adults and children 12 years and over: • **do not swallow tablet whole: chew completely** • to relieve symptoms, **chew** 1 tablet before swallowing • do not use more than 2 chewable tablets in 24 hours • children under 12 years: ask a doctor. **Other information** • **each tablet contains:** calcium 320 mg; magnesium 70 mg • read the directions and warnings before use • keep the carton. It contains important information. • store at 20°-25°C (68°-77°F) • protect from moisture. • **Do not use if pouch is torn or damaged.** **Inactive ingredients** cellulose acetate, corn starch, croscopolone, D&C red no. 7 calcium lake, dextrose excipient, FD&C blue no. 1 aluminum lake, FD&C red no. 40 aluminum lake, flavors, gum arabic, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, maltodextrin, mineral oil, modified starch, sucralose

**Questions or comments?** 1-800-755-4008 (toll free) or 215-273-8755 (collect)

Distributed by: Johnson & Johnson CONSUMER INC.

MCNEIL CONSUMER HEALTHCARE DIVISION, FORT WASHINGTON, PA 19034 USA

©McNeil Consumer Pharmaceuticals Co. 2018



Heartburn Relief

on-the GO

Individually wrapped tablets perfect for:

NDC 16837-298-25

DUAL ACTION

Pepcid Complete

Famotidine 10mg Acid reducer

Calcium carbonate 800mg Antacid

Magnesium hydroxide 165mg Antacid

Just One Tablet!

Relieves Heartburn Due to Acid Indigestion

Berry Flavor

actual size

8 Chewable Tablets

<b>Drug Facts</b> (continued)	<b>Stop use and ask a doctor if</b> <ul style="list-style-type: none"><li>• your heartburn continues or worsens</li><li>• you need to take this product for more than 14 days</li></ul> <b>If pregnant or breast-feeding</b> , ask a health professional before use.
<b>Other information</b> <ul style="list-style-type: none"><li>• children under 12 years: ask a doctor</li></ul>	<b>Directions</b> • Adults and children 12 years and older: take 1 tablet twice daily with meals and at bedtime. If symptoms do not improve after 2 weeks of use, ask your doctor.
<b>Questions or comments?</b> 1-800-755-4008 (toll-free) or 215-273-8755 (collect)	<b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)
<b>Drug Facts</b> (continued)	<b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)
<b>Drug Facts</b> (continued)	<b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

<b>Drug Facts</b> (continued)	<b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)
<b>Drug Facts</b> (continued)	<b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)
<b>Drug Facts</b> (continued)	<b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)
<b>Drug Facts</b> (continued)	<b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)
<b>Drug Facts</b> (continued)	<b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Pepcid Complete

Consumer

Distributed by:

(Johnson & Johnson)

MICHEL CONSUMER HEALTHCARE DIVISION

FORT WASHINGTON, PA 19034 USA

©McNeil Consumer Pharmaceuticals Co. 2018

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Have the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty, or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

7 16837 29108 4

-----  
**This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.**  
-----

/s/  
-----

THERESA M MICHELE  
06/11/2019 02:32:03 PM