



Contains: 100 Individual Swabs, 0.034 fl. oz. (1 mL) Each

*equivalent to 32 mg of Chlorhexidine Gluconate per pouch

Non-sterile Solution
ANTISEPTIC

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]



REORDER NO. B10800
NDC 10819-1080-1

5PB10800

Drug Facts

Active ingredients

Chlorhexidine gluconate 3.15% w/v Antiseptic
Isopropyl alcohol 70% v/v Antiseptic

Purpose

Antiseptic
Antiseptic

Use For the preparation of the patient's skin prior to an injection

Warnings

- for external use only • flammable, keep away from fire or flame
- Do not use with electrocautery procedures

Allergy alert: This product may cause a severe allergic reaction.

- Symptoms may include:
- wheezing/difficulty breathing • shock • facial swelling
 - hives • rash
- If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- on patients allergic to chlorhexidine gluconate or any other ingredient in this product
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser
- under occlusive patch

Drug Facts (continued)

When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- maximum treatment area for one swab is approximately 2.5 by 2.5 inches (6 by 6 cm)
- tear open packet and remove swab. Do not unfold swab.
- prior to injection, apply swab to the procedure site by holding swab between thumb and index finger. Apply swab to skin using repeated back-and-forth strokes for 15 seconds.
- allow the prepped area to air dry for 30 seconds
- do not blot or wipe dry
- discard after a single use

Drug Facts (continued)

Other information

- store between 68-77°F (20-25°C)
- avoid freezing and excessive heat above 104°F (40°C)
- for hospital and professional use only

Inactive ingredient

USP purified water

Questions?

call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)

NDC 10819-1080-1

REORDER NO. B10800



[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]

ANTISEPTIC
Non-sterile Solution

*equivalent to 32 mg of Chlorhexidine Gluconate per pouch

FOR EXTERNAL USE ONLY

- Single step, broad spectrum, persistent antiseptic that significantly reduces the number of microorganisms on intact skin
- Use for the preparation of the patient's skin prior to an injection
- Helps reduce bacteria that potentially can cause skin infection
- Latex free
- For hospital and professional use only

Contains: 100 Individual Swabs, 0.034 fl.oz. (1 mL) Each



(01)10310819000014



SWAB

STORAGE

- store between 68-77°F (20-25°C)
- avoid freezing and excessive heat above 104°F (40°C)

CONTENTS:

100 packets of swabs premoistened with 1 mL of a solution of 3.15% (w/v) Chlorhexidine gluconate, 70% (v/v) Isopropyl alcohol, and USP purified water

Do not use hypochlorite bleach on fabrics exposed to chlorhexidine gluconate.

LOT

EXP.

SWAB

Professional Disposables International, Inc.
Orangeburg, NY
10962-1376 USA
1-800-999-6423
www.pdihc.com



Prevantics® Swab

NET CONTENTS: 10 BOXES OF 100 INDIVIDUAL SWABS

REORDER NO. B10800

NDC 10819-1080-1

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]

Antiseptic

Non-sterile Solution

*equivalent to 32 mg of Chlorhexidine Gluconate per pouch

Drug Facts		Drug Facts (continued)	
Active ingredients	Purpose	Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
Chlorhexidine gluconate 3.15% w/v.	Antiseptic	Directions <ul style="list-style-type: none"> • use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. • maximum treatment area for one swab is approximately 2.5 by 2.5 inches (6 by 6 cm) • tear open packet and remove swab. Do not unfold swab. • prior to injection, apply swab to the procedure site by holding swab between thumb and index finger. Apply swab to skin using repeated back-and-forth strokes for 15 seconds. • allow the prepped area to air dry for 30 seconds • do not blot or wipe dry • discard after a single use 	
Isopropyl alcohol 70% v/v.	Antiseptic		
Use For the preparation of the patient's skin prior to an injection		Other information <ul style="list-style-type: none"> • store between 68-77°F (20-25°C) • avoid freezing and excessive heat above 104°F (40°C) • for hospital and professional use only 	
Warnings <ul style="list-style-type: none"> • For external use only • Flammable, keep away from fire or flame • Do not use with electrocautery procedures 			
Allergy alert: This product may cause a severe allergic reaction. Symptoms may include: <ul style="list-style-type: none"> • wheezing/difficulty breathing • shock • facial swelling • hives • rash If an allergic reaction occurs, stop use and seek medical help right away.		Inactive ingredient USP purified water	
Do not use <ul style="list-style-type: none"> • on patients allergic to chlorhexidine gluconate or any other ingredient in this product • for lumbar puncture or in contact with the meninges • on open skin wounds or as a general skin cleanser • under occlusive patch 			
When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.		Questions? call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)	
Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.			

8NB10800

Each swab contains 0.034 fl. oz. (1 mL)

 Professional Disposables International, Inc.
 Orangeburg, NY 10962-1376 USA
 1-800-999-6423 www.pdihc.com



(01)20310819000011



NDC 10819-1080-2 REORDER NO. B11400

Prevantics®

[Chlorhexidine Gluconate (3.15%)*
and Isopropyl Alcohol (70%) Swab]

ANTISEPTIC

Non-sterile Solution

Packaging is sterile when in sterile kit

*equivalent to 32 mg of Chlorhexidine Gluconate per pouch

FOR EXTERNAL USE ONLY

- Use for the preparation of the patient's skin prior to an injection
- Helps reduce bacteria that potentially can cause skin infection
- For hospital and professional use only • Latex free
- Do not use with electrocautery procedures

IMPORTANT: See insert for complete **Drug Facts** and safety information.

One Swab, 0.034 fl. oz. (1 mL) Each



(01)00310819000864



SWAB



TEAR BELOW

CONTENTS: One swab premoistened with 1 mL of a solution of 3.15% (w/v) Chlorhexidine gluconate, 70% (v/v) Isopropyl alcohol, and USP purified water. **DIRECTIONS:** Use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. Maximum treatment area for one swab is approximately 2.5 by 2.5 inches (6 by 6 cm). Tear open packet and remove swab. Do not unfold swab. Prior to injection, apply swab to the procedure site by holding swab between thumb and index finger. Apply swab to skin using repeated back-and-forth strokes for 15 seconds.

Allow the prepped area to air dry for 30 seconds. Do not blot or wipe dry. Discard after a single use. **WARNINGS: For external use only. Flammable, keep away from fire or flame.**

Allergy alert: This product may cause a severe allergic reaction. Symptoms may include: • wheezing/difficulty breathing • shock

• facial swelling • hives • rash **Keep out of reach**

of children. If swallowed, get medical help or

contact a Poison Control Center right away.

Professional Disposables International, Inc.

Orangeburg, NY 10962-1376 USA

1-800-999-6423 www.pdihc.com



2RB114X

LOT

EXP.

Prevantics® Swab

REORDER NO. B11400
NDC 10819-1080-2

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]

Antiseptic

Non-sterile Solution

Packaging is sterile when in sterile kit

*equivalent to 32 mg of Chlorhexidine Gluconate per pouch

Contains: 3000 Individual Swabs, 0.034 fl. oz. (1 mL) Each

<p>Drug Facts</p>	<p>Drug Facts (continued)</p>
<p>Active ingredients</p> <p>Chlorhexidine gluconate 3.15% w/v Antiseptic Isopropyl alcohol 70% v/v Antiseptic</p>	<p>Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p> <p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p>
<p>Use For the preparation of the patient's skin prior to an injection</p>	
<p>Warnings</p> <ul style="list-style-type: none"> • For external use only • Flammable, keep away from fire or flame • Do not use with electrocautery procedures 	<p>Directions</p> <ul style="list-style-type: none"> • use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. • maximum treatment area for one swab is approximately 2.5 by 2.5 inches (6 by 6 cm) • tear open packet and remove swab. Do not unfold swab. • prior to injection, apply swab to the procedure site by holding swab between thumb and index finger. Apply swab to skin using repeated back-and-forth strokes for 15 seconds. • allow the prepped area to air dry for 30 seconds • do not blot or wipe dry • discard after a single use
<p>Allergy alert: This product may cause a severe allergic reaction. Symptoms may include: • wheezing/difficulty breathing • shock • facial swelling • hives • rash If an allergic reaction occurs, stop use and seek medical help right away.</p>	<p>Other information</p> <ul style="list-style-type: none"> • store between 68-77°F (20-25°C) • avoid freezing and excessive heat above 104°F (40°C) • for hospital and professional use only
<p>Do not use</p> <ul style="list-style-type: none"> • on patients allergic to chlorhexidine gluconate or any other ingredient in this product • for lumbar puncture or in contact with the meninges • on open skin wounds or as a general skin cleanser • under occlusive patch 	<p>Inactive ingredient USP purified water</p>
<p>When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.</p>	<p>Questions? call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)</p>

8RB11400

 Professional Disposables International, Inc.
Orangeburg, NY 10962-1376 USA
1-800-999-6423 www.pdihc.com



(01)20310819000868



+Prevantics® Swab REORDER NO. B11400
 [Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab] Antiseptic NDC 10819-1080-2
 Non-sterile Solution • Packaging is sterile when in sterile kit
 *equivalent to 32 mg of Chlorhexidine Gluconate per pouch

Drug Facts	
Active ingredients	Purpose
Chlorhexidine gluconate 3.15% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic
Use For the preparation of the patient's skin prior to an injection	
Warnings	
<ul style="list-style-type: none"> • For external use only • Flammable, keep away from fire or flame • Do not use with electrocautery procedures 	
Allergy alert: This product may cause a severe allergic reaction. Symptoms may include: • wheezing/difficulty breathing • shock • facial swelling • hives • rash If an allergic reaction occurs, stop use and seek medical help right away.	
Do not use <ul style="list-style-type: none"> • on patients allergic to chlorhexidine gluconate or any other ingredient in this product • for lumbar puncture or in contact with the meninges • on open skin wounds or as a general skin cleanser • under occlusive patch 	
When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent ▶	

Drug Facts (continued)
injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.
Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.
Directions
<ul style="list-style-type: none"> • use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. • maximum treatment area for one swab is approximately 2.5 by 2.5 inches (6 by 6 cm) • tear open packet and remove swab. Do not unfold swab. • prior to injection, apply swab to the procedure site by holding swab between thumb and index finger. Apply swab to skin using repeated back-and-forth strokes for 15 seconds. • allow the prepped area to air dry for 30 seconds • do not blot or wipe dry • discard after a single use
Other information
<ul style="list-style-type: none"> • store between 68-77°F (20-25°C) • avoid freezing and excessive heat above 104°F (40°C) • for hospital and professional use only
Inactive ingredient USP purified water
Questions? call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)
Each Swab contains 0.034 fl. oz. (1 mL) +PDI Professional Disposables International, Inc. Orangeburg, NY 10962-1376 USA 1-800-999-6423 www.pdihc.com  
5QB1141S

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

THERESA M MICHELE
01/16/2019 01:15:10 PM