

(b) (4)

Carton Size : 83.00 x 22.00 x 112.00
 Track : 15/11/2018, 23/11/2018

Drug Facts

Active ingredient (in each capsule)	Purpose
Loperamide HCl 2 mg.....	Anti-diarrheal
Use controls symptoms of diarrhea, including Travelers' Diarrhea	
Warnings	
Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl	
Heart alert: Taking more than directed can cause serious heart problems or death	
Do not use ■if you have bloody or black stool ■in children under 12 years of age	
Ask a doctor before use if you have ■ fever ■mucus in the stool ■ a history of liver disease ■ a history of abnormal heart rhythm	
Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.	
When using this product tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.	
Stop use and ask a doctor if ■symptoms get worse ■diarrhea lasts for more than 2 days ■you get abdominal swelling or bulging. These may be signs of a serious condition.	
if pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions	
■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea.	
■ not for use in children under 12 years of age	
■ adults and children 12 years and over: 2 softgels after the first loose stool; 1 softgel after each subsequent loose stool; but no more than 4 softgels in 24 hours	
Other information	
■ store at 20°-25°C (68°-77°F). Protect from Light.	■ do not use if carton or blister unit is open or torn
■ avoid excessive heat above 40°C (104°F)	■ see side panel for lot number and expiration date
Inactive ingredients butylated hydroxyanisole, edible ink, FD&C Blue #1, gelatin, glycerin, glyceryl caprylate, polyoxyl 40 hydrogenated castor oil, purified water	
Questions or comments? call toll free 1-888-235-2466	

Loperamide 2 mg Softgels*

*All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Imodium® A-D

4 Softgels*
 each Liquid-filled capsule contains 2 mg loperamide HCl



Suitable for adults and children 12 years and over

Anti-Diarrheal

Soft Gelatin Capsules, 2 mg


Loperamide HCl

Compare to the active ingredient in Imodium® A-D

NDC 0000-000-00

THIS PRODUCT IS PACKAGED IN A CHILD-RESISTANT AND TAMPER EVIDENT PACKAGE. USE ONLY IF BLISTERS ARE INTACT.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.



1 000001 000001 3

Loperamide 2 mg Softgels*

Lot #
 Expiration Date
 Rev # 1118



Manufactured for:
BIONPHARMA
 600 Alexander Road,
 Princeton, NJ 08540

BIONPHARMA

Labeling Format Information:
 Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique

Drug Facts:	14pt	Leading:	6.5pt
Header:		Bullets:	
Subheader:	6pt	Barlines:	
Body Text:	8pt 6pt	Hairlines:	5pt .5pt
Drug Facts (continued):	N/A		

Horizontal Scale: 100% Average Kerning: 0%

(b) (4)

Carton Size : 83.00 x 22.00 x 112.00
Track : 02/12/2018

Drug Facts	
Active ingredient (in each capsule) Loperamide HCl 2 mg.....	Purpose Anti-diarrheal
Use controls symptoms of diarrhea, including Travelers' Diarrhea	
Warnings	
Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl	
Heart alert: Taking more than directed can cause serious heart problems or death	
Do not use ■if you have bloody or black stool ■in children under 12 years of age	
Ask a doctor before use if you have ■ fever ■mucus in the stool ■ a history of liver disease ■ a history of abnormal heart rhythm	
Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.	
When using this product tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.	
Stop use and ask a doctor if ■symptoms get worse ■diarrhea lasts for more than 2 days ■you get abdominal swelling or bulging. These may be signs of a serious condition.	
if pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions	
■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea.	
■ not for use in children under 12 years of age	
■ adults and children 12 years and over: 2 softgels after the first loose stool; 1 softgel after each subsequent loose stool; but no more than 4 softgels in 24 hours	
Other information	
■ store at 20°-25°C (68°-77°F). Protect from Light.	■ do not use if carton or blister unit is open or torn
■ avoid excessive heat above 40°C (104°F)	■ see side panel for lot number and expiration date
Inactive ingredients butylated hydroxyanisole, edible ink, FD&C Blue #1, gelatin, glycerin, glyceryl caprylate, polyoxyol 40 hydrogenated castor oil, purified water	
Questions or comments? call toll free 1-888-235-2466	

Loperamide 2 mg Softgels*

*All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Imodium® A-D

6 Softgels*
each Liquid-filled capsule contains 2 mg loperamide HCl



Suitable for adults and children 12 years and over

Anti-Diarrheal

Soft Gelatin Capsules, 2 mg

Loperamide HCl

Compare to the active ingredient in Imodium® A-D

NDC 0000-000-00

THIS PRODUCT IS PACKAGED IN A CHILD-RESISTANT AND TAMPER EVIDENT PACKAGE. USE ONLY IF BLISTERS ARE INTACT.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.



1 000001 000001 3

Lot #
Expiration Date
Rev # 1118



Manufactured for:
BIONPHARMA
600 Alexander Road,
Pinebluff, NJ 08540

Loperamide 2 mg Softgels*

BIONPHARMA

Labeling Format Information:

Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique

Drug Facts:	14pt	Leading:	6.5pt
Header:		Bullets:	
Subheader:	6pt	Barlines:	
Body Text:	8pt 6pt	Hairlines:	5pt .5pt
Drug Facts (continued):	N/A		

Horizontal Scale: 100% Average Kerning: 0%

(b) (4)

Carton Size : 83.00 x 22.00 x 112.00
 Track : 15/11/2018, 23/11/2018

Drug Facts

Active ingredient (in each capsule)
 Loperamide HCl 2 mg

Purpose
 Anti-diarrheal

Use controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings
Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl
Heart alert: Taking more than directed can cause serious heart problems or death
Do not use if you have bloody or black stool in children under 12 years of age
Ask a doctor before use if you have fever mucus in the stool a history of liver disease a history of abnormal heart rhythm
Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.
When using this product tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.
Stop use and ask a doctor if symptoms get worse diarrhea lasts for more than 2 days you get abdominal swelling or bulging. These may be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 drink plenty of clear fluids to help prevent dehydration caused by diarrhea.
 not for use in children under 12 years of age
 adults and children 12 years and over: 2 softgels after the first loose stool; 1 softgel after each subsequent loose stool; but no more than 4 softgels in 24 hours

Other information
 store at 20°-25°C (68°-77°F). Protect from Light. do not use if carton or blister unit is open or torn
 avoid excessive heat above 40°C (104°F) see side panel for lot number and expiration date

Inactive ingredients butylated hydroxyanisole, edible ink, FD&C Blue #1, gelatin, glycerin, glyceryl caprylate, polyoxy 40 hydrogenated castor oil, purified water

Questions or comments? call toll free 1-888-235-2466

Loperamide 2 mg Softgels*

*All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Imodium® A-D

8 Softgels*
 each Liquid-filled capsule contains 2 mg loperamide HCl

Suitable for adults and children 12 years and over

Anti-Diarrheal

Soft Gelatin Capsules, 2 mg

Loperamide HCl

Compare to the active ingredient in Imodium® A-D


NDC 00000-000-00



Loperamide 2 mg Softgels*

THIS PRODUCT IS PACKAGED IN A CHILD-RESISTANT AND TAMPER EVIDENT PACKAGE. USE ONLY IF BLISTERS ARE INTACT.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.



1
000000 000000
3

Lot #
 Expiration Date
 Rev # 11/18

Manufactured for:
BIONPHARMA
 600 Alexander Road,
 Princeton, NJ 08540

BIONPHARMA

Labeling Format Information:
 Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique

Drug Facts:	14pt	Leading:	6.5pt
Header:		Bullets:	
Subheader:	6pt	Barlines:	5pt
Body Text:	8pt 6pt	Hairlines:	.5pt
Drug Facts (continued):	N/A		

Horizontal Scale: 100%
 Average Kerning: 0%

(b) (4)

Carton Size : 83.00 x 22.00 x 112.00
Track : 15/11/2018, 23/11/2018

Drug Facts	
Active ingredient (in each capsule) Loperamide HCl 2 mg.....	Purpose Anti-diarrheal
Use controls symptoms of diarrhea, including Travelers' Diarrhea	
Warnings Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl Heart alert: Taking more than directed can cause serious heart problems or death Do not use ■ if you have bloody or black stool ■ in children under 12 years of age Ask a doctor before use if you have ■ fever ■ mucus in the stool ■ a history of liver disease ■ a history of abnormal heart rhythm Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs. When using this product tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery. Stop use and ask a doctor if ■ symptoms get worse ■ diarrhea lasts for more than 2 days ■ you get abdominal swelling or bulging. These may be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea. ■ not for use in children under 12 years of age ■ adults and children 12 years and over: 2 softgels after the first loose stool; 1 softgel after each subsequent loose stool; but no more than 4 softgels in 24 hours	
Other information ■ store at 20°-25°C (68°-77°F). Protect from Light. ■ do not use if carton or blister unit is open or torn ■ avoid excessive heat above 40°C (104°F) ■ see side panel for lot number and expiration date	
Inactive ingredients butylated hydroxyanisole, edible ink, FD&C Blue #1, gelatin, glycerin, glyceryl caprylate, polyoxyl 40 hydrogenated castor oil, purified water	
Questions or comments? call toll free 1-888-235-2466	

Loperamide 2 mg Softgels*

*All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Imodium® A-D



12 Softgels*
each Liquid-filled capsule contains 2 mg loperamide HCl

Suitable for adults and children 12 years and over

Anti-Diarrheal

Soft Gelatin Capsules, 2 mg


Loperamide HCl

Compare to the active ingredient in Imodium® A-D

NDC 00000-000-00

THIS PRODUCT IS PACKAGED IN A CHILD RESISTANT AND TAMPER EVIDENT PACKAGE. USE ONLY IF BLISTER IS UNHARMED.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.



1 000000 000000 3

Loperamide 2 mg Softgels*

Lot #
Expiration Date
Rev # 11/18

Manufactured for:
BIONPHARMA
600 Alexander Road,
Princeton, NJ 08540

BIONPHARMA

Labeling Format Information:

Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique

Drug Facts:	14pt	Leading:	6.5pt
Header:	6pt	Bullets:	5pt
Subheader:	8pt	Barlines:	5pt
Body Text:	N/A	Hairlines:	5pt

Horizontal Scale: 100%
Average Kerning: 0%

(b) (4)

Label Size: 3+9/32 x 7/8 x 3+7/8
 Dieline: J80619A
 Track: 15/11/2018, 23/11/2018

Lot #
Expiration Date
Rev # 11/18

Manufactured for:
BIONPHARMA
600 Alexander Road,
Princeton, NJ 08540

Loperamide 2 mg
Soft Gelatin Capsules, 2 mg

Anti-Diarrheal

Suitable for adults and children 12 years and over

Loperamide HCl

Compare to the active ingredient in Imodium® A-D

NDC 00000-000-00

18 Softgels*
*each Liquid-filled capsule contains 2 mg loperamide HCl

THIS PRODUCT IS PACKAGED IN A CHILD-RESISTANT AND TAMPER EVIDENT PACKAGE. USE ONLY IF BLISTERS ARE INTACT.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

1

3 00000 00000 1

All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Imodium® A-D

Loperamide 2 mg
Softgels*

Drug Facts	Purpose
Active ingredient (in each capsule) Loperamide HCl 2 mg.....	Anti-diarrheal
Use controls symptoms of diarrhea, including Travelers' Diarrhea	
Warnings Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl Heart alert: Taking more than directed can cause serious heart problems or death	
Do not use ■ if you have bloody or black stool ■ in children under 12 years of age	
Ask a doctor before use if you have ■ fever ■ mucus in the stool ■ a history of liver disease ■ a history of abnormal heart rhythm	
Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.	
When using this product tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.	
Stop use and ask a doctor if ■ symptoms get worse ■ diarrhea lasts for more than 2 days ■ you get abdominal swelling or bulging. These may be signs of a serious condition.	
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea. ■ not for use in children under 12 years of age ■ adults and children 12 years and over: 2 softgels after the first loose stool; 1 softgel after each subsequent loose stool; but no more than 4 softgels in 24 hours	
Other information ■ store at 20°-25°C (68°-77°F). Protect from Light. ■ do not use if carton or blister unit is open or torn ■ avoid excessive heat above 40°C (104°F) ■ see side panel for lot number and expiration date	
Inactive ingredients butylated hydroxyanisole, edible ink, FD&C Blue #1, gelatin, glycerin, glyceryl caprylate, polyoxyl 40 hydrogenated castor oil, purified water	
Questions or comments? call toll free 1-888-235-2466	

NO INK NO VARNISH

PRINT SIDE GLUE


BIONPHARMA


Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique

Drug Facts:	14pt	Leadings:	6.5pt
Header:		Bullets:	
Subheader:	6pt	Barlines:	7pt
Body Text:	8pt 6pt	Hairlines:	.5pt
Drug Facts (continued):	N/A		


Horizontal Scale: 100%

Average Kerning: 0%






NOT OFFICIAL SIZE/DIELINE



Carton Size : 0.75" x 3.375" x 4.375"
Track : 15/11/2018, 23/11/2018



Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique			
Drug Facts:	14pt	Leading:	6.5pt
Header:	8pt	Bullets:	5pt
Subheader:	6pt	Barlines:	1.5pt
Body Text:	6pt	Hairlines:	.5pt
Drug Facts (continued):	N/A		
Horizontal Scale: 100%		Average Kerning: 0%	



Track : 16/11/2018