



Actual Size 100%

FDA00037-r5

DRUG FACTS INFO (updated 5/18/18)		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt **	9 pt
Drug Facts Continued	8 pt	-
Headings [†]	8 pt-bold italic [†]	8 pt
Subheadings	≥6 pt	6 pt
Body Text	≥6 pt	6 pt
Bullets	5 pt	3.5 pt
Leading	≥0.5 pt	6.5 pt
Hairlines	0.5 pt rule	0.5 pt
Box and Barlines [†]	>0.5 pt rule	1 pt
Characters Per Inch:	No more than 39	≤39

[†] Headings are to be at least 2pts larger than Body Text.
^{**} Drug Facts Title to be larger than all other type size used in DFP
[†] FDA Recommends 2.5pt rule



Actual Size 100%

FDA00039_r6

DRUG FACTS INFO (updated 5/18/18)		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt **	9 pt
Drug Facts Continued	8 pt	-
Headings*	8 pt-bold italic*	8 pt
Subheadings	≥6 pt	6 pt
Body Text	≥8 pt	6 pt
Bullets	5 pt	3.5 pt
Leading	≥0.5 pt	6.5 pt
Hairlines	0.5 pt rule	0.5 pt
Box and Barlines†	>0.5 pt rule	1 pt
Characters Per Inch:	No more than 39	≤39

* Headings are to be at least 2xts larger than Body Text.
 ** Drug Facts Title to be larger than all other type size used in DFP
 † FDA Recommends 2.5pt rule

CATEGORY		REQUIRED	ACTUAL IN FILE
Drug Facts Title	> 8 pt **	9 pt	
Drug Facts Continued	8 pt	8 pt	
Headings*	8 pt-bold italic*	8 pt	
Subheadings	> 6 pt	6 pt	
Body Text	> 6 pt	6 pt	
Bullets	5 pt	3.5 pt	
Leading	> 0.5 pt	0.5 pt	
Headings	0.5 pt rule	0.5 pt	
Box and Barlines	> 0.5 pt rule	1 pt	
Character's Per Inch:	No more than 39	< 39	

* Headings are to be at least 2pts larger than other type size used in DFP
 ** Drug Facts Title to be larger than all other type size used in DFP
 † FDA Recommends 2-3pt rule

FDA00149-3

Actual Size 100%

Glue Area

Drug Facts

Active ingredient (in each tablet) **Purpose**
 Loratadine 5 mg.....Antihistamine

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 ■ runny nose ■ itchy, watery eyes
 ■ sneezing ■ itching of the nose or throat

Warnings
Do not use if you have ever had an allergic reaction to this product or any of its ingredients.
Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.
When using this product do not take more than directed. Taking more than directed may cause drowsiness.
Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

adults and children 6 years and over	chew 2 tablets daily; not more than 2 tablets in 24 hours
children 2 to under 6 years of age	chew 1 tablet daily; not more than 1 tablet in 24 hours
children under 2 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information

- phenyleketonurics: contains phenylalanine 1.4 mg per tablet
- safety sealed: do not use if the individual blister unit imprinted with Children's Claritin® is open or torn
- store between 20° to 25°C (68° to 77°F)

Inactive ingredients
 aspartame, citric acid anhydrous, colloidal silicon dioxide, D&C red No. 27 aluminum lake, FD&C blue No. 2 aluminum lake, flavor, magnesium stearate, mannitol, microcrystalline cellulose, sodium starch glycolate, stearic acid

Questions or comments?
 1-800-CLARITIN (1-800-252-7484) or www.claritin.com

LOT/EXP/AREA
NO VARNISH
NO INK

Bayer
 Product of Ireland.
 Whippany, NJ 07981
 Dist by: Bayer Healthcare LLC
 © 2019 Bayer.
 Claritin & Balloon Design and the Claritin Vista are registered trademarks of Bayer.

00000000


00000000

10 CHEWABLE TABLETS

Children's
Claritin
Chewables

Children's
Claritin
Chewables

LOT/EXP/AREA
NO VARNISH
NO INK



CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	> 8 pt **	9 pt
Drug Facts Continued	8 pt	8 pt
Headings*	8 pt-bold italic*	8 pt
Subheadings	> 6 pt	6 pt
Body Text	> 6 pt	6 pt
Bullets	5 pt	3.5 pt
Leading	> 0.5 pt	6.5 pt
Headings	0.5 pt rule	0.5 pt
Box and Barlines†	> 0.5 pt rule	1 pt
Characters Per Inch:	No more than 39	< 39

† FDA recommends ≥ 6 pt rule
 ** Drug Facts Title to be larger than all other type used in DFP
 * Headings are to be at least 2 pts larger than Body Text

DRUG FACTS INFO (updated 5/18/13)
 FDA00147-3

Actual Size 100%

Glue Area

Children's Claritin Chewables

10 CHEWABLE TABLETS

NO VARNISH

Children's Claritin Chewables

LOT/EXP AREA NO VARNISH NO INK

Children's Claritin Chewables

2 years and older

5 mg/antihistamine loratadine chewable tablets

Indoor & Outdoor Allergies

24 Hour Relief of:

- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itchy Throat or Nose

Non-Drowsy*

BubbleGum Flavored

10 CHEWABLE TABLETS

See Drug Facts Panel.

* When taken as directed.

Drug Facts		Drug Facts (continued)	
Active ingredient (in each tablet)	Purpose	Directions	
Loratadine 5 mg.....	Antihistamine	<ul style="list-style-type: none"> chew or crush tablets completely before swallowing. adults and children 6 years and over: chew 2 tablets daily; not more than 2 tablets in 24 hours children 2 to under 6 years of age: chew 1 tablet daily; not more than 1 tablet in 24 hours children under 2 years of age: ask a doctor consumers with liver or kidney disease: ask a doctor 	
Uses	temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:	Other information	
<ul style="list-style-type: none"> runny nose sneezing 	<ul style="list-style-type: none"> itchy, watery eyes itching of the nose or throat 	<ul style="list-style-type: none"> phenylketonurics: contains phenylalanine 1.9 mg per tablet safety sealed: do not use if the individual blister unit imprinted with Children's Claritin® is open or torn store between 20° to 25°C (68° to 77°F) 	
Warnings	Do not use if you have ever had an allergic reaction to this product or any of its ingredients.	Inactive ingredients	
Ask a doctor before use if you have liver or kidney disease.	Your doctor should determine if you need a different dose.	aspartame, carmine, citric acid, colloidal silicon dioxide, flavor, magnesium stearate, mannitol, microcrystalline cellulose, sodium starch glycolate, stearic acid	
When using this product	do not take more than directed. Taking more than directed may cause drowsiness.	Questions or comments?	
Stop use and ask a doctor	if an allergic reaction to this product occurs. Seek medical help right away.	1-800-CLARITIN (1-800-252-7484) or www.claritin.com	
If pregnant or breast-feeding,	ask a health professional before use.		
Keep out of reach of children.	In case of overdose, get medical help or contact a Poison Control Center right away.		

Children's Claritin Chewables

© 2019 Bayer. are registered trademarks of Bayer.

Claritin & Balloon Design and the Claritin Vial are registered trademarks of Bayer.

Dist by: Bayer Healthcare LLC

Whisper, NJ 07981

Product of Ireland.

00000000

Bayer




Actual Size 100%

FDA00156-3


DRUG FACTS INFO (updated 5/18/18)		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt **	9 pt
Drug Facts Continued	8 pt	8 pt
Headings*	8 pt-bold italic*	8 pt
Subheadings	≥6 pt	6 pt
Body Text	≥6 pt	6 pt
Bullets	5 pt	3.5 pt
Leading	≥0.5 pt	≥0.5 pt
Hairlines	0.5 pt rule	.5 pt
Box and Barlines†	>0.5 pt rule	1 pt
Characters Per Inch:	No more than 39	≤ 39

* Headings are to be at least 2pts larger than Body Text.
 ** Drug Facts Title to be larger than all other type size used in DFP
 † FDA Recommends 2.5pt rule




<p>Drug Facts Active ingredient (in each tablet) Purpose Loratadine 5 mg Antihistamine</p> <p>Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ itchy, watery eyes ■ sneezing ■ itching of the nose or throat</p> <p>Warnings Do not use if you have ever had an allergic reaction to this product or any of its ingredients. Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. When using this product do not take more than directed. Taking more than directed may cause drowsiness. Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>	<p>Drug Facts (continued) Directions ■ chew or crush tablets completely before swallowing.</p> <table border="1"> <tr> <td>adults and children 6 years and over</td> <td>chew 2 tablets daily; not more than 2 tablets in 24 hours</td> </tr> <tr> <td>children 2 to under 6 years of age</td> <td>chew 1 tablet daily; not more than 1 tablet in 24 hours</td> </tr> <tr> <td>children under 2 years of age</td> <td>ask a doctor</td> </tr> <tr> <td>consumers with liver or kidney disease</td> <td>ask a doctor</td> </tr> </table> <p>Other information ■ phenylketonurics: contains phenylalanine 1.4 mg per tablet ■ safety sealed: do not use if the individual blister unit imprinted with Children's Claritin® is open or torn ■ store between 20° to 25°C (68° to 77°F)</p> <p>Inactive ingredients aspartame, citric acid anhydrous, colloidal silicon dioxide, D&C red No. 27 aluminum lake, FD&C blue No. 2 aluminum lake, flavor, magnesium stearate, mannitol, microcrystalline cellulose, sodium starch glycolate, stearic acid</p> <p>Questions or comments? 1-800-CLARITIN (1-800-252-7484) or www.claritin.com</p>	adults and children 6 years and over	chew 2 tablets daily; not more than 2 tablets in 24 hours	children 2 to under 6 years of age	chew 1 tablet daily; not more than 1 tablet in 24 hours	children under 2 years of age	ask a doctor	consumers with liver or kidney disease	ask a doctor
adults and children 6 years and over	chew 2 tablets daily; not more than 2 tablets in 24 hours								
children 2 to under 6 years of age	chew 1 tablet daily; not more than 1 tablet in 24 hours								
children under 2 years of age	ask a doctor								
consumers with liver or kidney disease	ask a doctor								

Bayer, the Bayer Cross, Claritin, Children's Claritin & Balloon Design and the Claritin Vista are registered trademarks of Bayer.
 © 2019 Bayer.
 Dist. by: Bayer HealthCare LLC
 Whippany, NJ 07981
 Product of Ireland.
 0000000



Bayer



Actual Size 100%

FDA00155-3

DRUG FACTS INFO (updated 5/18/18)		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt **	8 pt
Drug Facts Continued	8 pt	8 pt
Headings*	8 pt-bold italic*	8 pt
Subheadings	≥8 pt	8 pt
Body Text	≥8 pt	8 pt
Bullets	5 pt	3.5 pt
Leading	≥0.5 pt	≥0.5 pt
Headlines	0.5 pt rule	.5 pt
Box and Barlines†	>0.5 pt rule	1 pt
Characters Per Inch	No more than 38	≤38

* Headings are to be at least 2xpt larger than Body Text.
 ** Drug Facts Title to be larger than all other type size used in DFP.
 † FDA Recommends 2.5xpt rule

Drug Facts		Drug Facts (continued)	
Active Ingredient (in each tablet) Loratadine 5 mg.....	Purpose Antihistamine	Directions ■ chew or crush tablets completely before swallowing.	
Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ itchy, watery eyes ■ sneezing ■ itching of the nose or throat		adults and children 6 years and over	chew 2 tablets daily; not more than 2 tablets in 24 hours
		children 2 to under 6 years of age	chew 1 tablet daily; not more than 1 tablet in 24 hours
Warnings Do not use if you have ever had an allergic reaction to this product or any of its ingredients.		children under 2 years of age	ask a doctor
Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.		consumers with liver or kidney disease	ask a doctor
When using this product do not take more than directed. Taking more than directed may cause drowsiness.		Other information ■ phenylketonurics: contains phenylalanine 1.9 mg per tablet ■ safety sealed: do not use if the individual blister unit imprinted with Children's Claritin® is open or torn ■ store between 20° to 25°C (68° to 77°F)	
Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.		Inactive ingredients aspartame, camille, citric acid, colloidal silicon dioxide, flavor, magnesium stearate, mannitol, microcrystalline cellulose, sodium starch glycolate, stearic acid	
If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.		Questions or comments? 1-800-CLARITIN (1-800-252-7484) or www.claritin.com	



Actual Size 100%

FDA00146-3

DRUG FACTS INFO (updated 5/18/18)		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt**	9 pt
Drug Facts Continued	8 pt	8 pt
Headings*	8 pt-bold italic*	8 pt
Subheadings	≥6 pt	6 pt
Body Text	≥6 pt	6 pt
Bullets	5 pt	3.5 pt
Leading	≥0.5 pt	6.5 pt
Hairlines	0.5 pt rule	0.5 pt
Box and Barlines [†]	>0.5 pt rule	1 pt
Characters Per Inch:	No more than 39	≤ 39

* Headings are to be at least 2pts larger than Body Text.
 ** Drug Facts Title to be larger than all other type size used in DFP
 † FDA Recommends 2.5pt rule



Actual Size 100%

FDA00154-3

DRUG FACTS INFO (Updated 5/18/18)		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt **	9 pt
Drug Facts Continued	8 pt	8 pt
Headings*	8 pt-bold italic*	8 pt
Subheadings	≥6 pt	6 pt
Body Text	≥6 pt	6 pt
Bullets	5 pt	3.5 pt
Leading	≥0.5 pt	6.5 pt
Hairlines	0.5 pt rule	0.5 pt
Box and Barlines†	>0.5 pt rule	1 pt
Characters Per Inch:	No more than 39	≤ 39

* Headings are to be at least 2pts larger than Body Text.
 ** Drug Facts Title to be larger than all other type size used in DFP.
 † FDA Recommends 2.5pt rule.

Children's Claritin Chewables
loratadine chewable tablets
5 mg/antihistamine

50 CHEWABLE TABLETS



Indoor & Outdoor Allergies

Grape Flavored

50 CHEWABLE TABLETS



Bayer, the Bayer Cross, Claritin, Children's Claritin & Balloon Design and the Claritin Vista are registered trademarks of Bayer. © 2019 Bayer. Dist by: Bayer HealthCare LLC Whippany, NJ 07981 Product of Ireland. 00000000 Bayer



Drug Facts (continued)

Active ingredient (in each tablet) Loratadine 5 mg

Purpose Antihistamine

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: runny nose, itchy, watery eyes, sneezing, itching of the nose or throat

Warnings Do not use if you have ever had an allergic reaction to this product or any of its ingredients. Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. When using this product do not take more than directed. Taking more than directed may cause drowsiness. Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Questions or comments? 1-800-CLARITIN (1-800-252-7484) or www.claritin.com

Inactive ingredients aspartame, citric acid anhydrous, colloidal silicon dioxide, D&C red No. 27 aluminum lake, FD&C blue No. 2 aluminum lake, flavor, magnesium stearate, mannitol, microcrystalline cellulose, sodium starch glycolate, stearic acid

Other information ■ phenylethanolamines: contains phenylethanolamine 1.4 mg per tablet ■ safety sealed: do not use if the individual blister unit is imprinted with Children's Claritin® is open or torn ■ store between 20° to 25°C (68° to 77°F)

Directions ■ chew or crush tablets completely before swallowing. ■ chew 2 tablets in 24 hours; not more than 2 tablets daily. ■ children 2 to under 6 years and over: chew 1 tablet in 24 hours; not more than 1 tablet in 24 hours ■ children under 2 years of age: ask a doctor ■ consumers with liver or kidney disease: ask a doctor

Drug Facts (continued)

Active ingredient (in each tablet) Loratadine 5 mg

Purpose Antihistamine

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: runny nose, itchy, watery eyes, sneezing, itching of the nose or throat

Warnings Do not use if you have ever had an allergic reaction to this product or any of its ingredients. Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. When using this product do not take more than directed. Taking more than directed may cause drowsiness. Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Questions or comments? 1-800-CLARITIN (1-800-252-7484) or www.claritin.com

Inactive ingredients aspartame, citric acid anhydrous, colloidal silicon dioxide, D&C red No. 27 aluminum lake, FD&C blue No. 2 aluminum lake, flavor, magnesium stearate, mannitol, microcrystalline cellulose, sodium starch glycolate, stearic acid

Other information ■ phenylethanolamines: contains phenylethanolamine 1.4 mg per tablet ■ safety sealed: do not use if the individual blister unit is imprinted with Children's Claritin® is open or torn ■ store between 20° to 25°C (68° to 77°F)

Directions ■ chew or crush tablets completely before swallowing. ■ chew 2 tablets in 24 hours; not more than 2 tablets daily. ■ children 2 to under 6 years and over: chew 1 tablet in 24 hours; not more than 1 tablet in 24 hours ■ children under 2 years of age: ask a doctor ■ consumers with liver or kidney disease: ask a doctor

Actual Size 100%

FDA00148-3

DRUG FACTS INFO (updated 5/18/18)		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt.**	13 pt.
Drug Facts Continued	8 pt.	11.5 pt.
Headings*	8 pt.-bold italic*	11.5 pt.
Subheadings	≥6 pt.	8.5 pt.
Body text	≥6 pt.	8.5 pt.
Bullets	5 pt.	5 pt.
Leading	≥0.5 pt.	9.5 pt.
Headlines	0.5 pt. rule	0.5 pt.
Box and Barlines*	>0.5 pt. rule	1 pt.
Characters Per Inch:	No more than 38	≤ 38

* Headings are to be at least 20% larger than Body text.
 ** Drug Facts Title to be larger than all other type size used in DPP.
 † FDA recommends 2-24 pt. rule.

ages 2 years and older

Children's Claritin Chewables

loratadine chewable tablets 5 mg/antihistamine

Non-Drowsy* Allergy Relief

Indoor & Outdoor Allergies

Grape Flavored

50 2-Count Samples
Total 100 Chewable Tablets

*When taken as directed. See Drug Facts Panel.

PROFESSIONAL SAMPLES - NOT FOR SALE

ages 2 years and older

Children's Claritin Chewables

loratadine chewable tablets 5 mg/antihistamine

Non-Drowsy* Allergy Relief

Indoor & Outdoor Allergies

Grape Flavored

50 2-Count Samples
Total 100 Chewable Tablets

*When taken as directed. See Drug Facts Panel.

PROFESSIONAL SAMPLES - NOT FOR SALE

ages 2 years and older

Children's Claritin Chewables

loratadine chewable tablets 5 mg/antihistamine

Non-Drowsy* Allergy Relief

Indoor & Outdoor Allergies

Grape Flavored

50 2-Count Samples
Total 100 Chewable Tablets

*When taken as directed. See Drug Facts Panel.

PROFESSIONAL SAMPLES - NOT FOR SALE

ages 2 years and older

Children's Claritin Chewables

loratadine chewable tablets 5 mg/antihistamine

Non-Drowsy* Allergy Relief

Indoor & Outdoor Allergies

Grape Flavored

50 2-Count Samples
Total 100 Chewable Tablets

*When taken as directed. See Drug Facts Panel.

PROFESSIONAL SAMPLES - NOT FOR SALE

ages 2 years and older

Children's Claritin Chewables

loratadine chewable tablets 5 mg/antihistamine

Non-Drowsy* Allergy Relief

Indoor & Outdoor Allergies

Grape Flavored

50 2-Count Samples
Total 100 Chewable Tablets

*When taken as directed. See Drug Facts Panel.

PROFESSIONAL SAMPLES - NOT FOR SALE

ages 2 years and older

Children's Claritin Chewables

loratadine chewable tablets 5 mg/antihistamine

Drug Facts

Active ingredient (in each tablet) Purpose
Loratadine 5mg Antihistamine

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 ■ runny nose ■ itchy, watery eyes
 ■ sneezing ■ itching of the nose or throat

Warnings
 Do not use if you have ever had an allergic reaction to this product or any of its ingredients.
 Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.
 When using this product do not take more than directed. Taking more than directed may cause drowsiness.
 Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.
 If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 ■ chew or crush tablets completely before swallowing.

adults and children 6 years and over	chew 2 tablets daily; not more than 2 tablets in 24 hours
children 2 to under 6 years of age	chew 1 tablet daily; not more than 1 tablet in 24 hours
children under 2 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Drug Facts (continued)

Other information
 ■ phenylketonurics: contains phenylalanine 1.4 mg per tablet
 ■ safety sealed: do not use if the individual blister unit imprinted with Children's Claritin® is open or torn
 ■ store between 20° to 25°C (68° to 77°F)

Inactive ingredients
 aspartame, citric acid anhydrous, colloidal silicon dioxide, D&C red No. 27 aluminum lake, FD&C blue No. 2 aluminum lake, flavor, magnesium stearate, mannitol, microcrystalline cellulose, sodium starch glycolate, stearic acid

Questions or comments?
 1-800-CLARITIN (1-800-252-7484) or www.claritin.com

Bayer, the Bayer Cross, Claritin, Children's Claritin & Balloon Design and the Claritin Vista are registered trademarks of Bayer.
 © 2019 Bayer.
 Dist. by: Bayer HealthCare LLC, Whippany, NJ 07981
 Product of Ireland. 00000000 Bayer

**LOT/EXP AREA
NO VARNISH**

Actual Size 100%

FDA00153-3

CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt.**	12 pt.
Drug Facts Continued	8 pt.	10.5 pt.
Headings*	8 pt.-bold italic†	10.5 pt.
Subheadings	>6 pt.	8.5 pt.
Body Text	>6 pt.	8.5 pt.
Bullets	5 pt.	5 pt.
Leading	>0.5 pt.	0.5 pt.
Headlines	0.5 pt.rule	0.5 pt.
Box and Barlines†	>0.5 pt.rule	1 pt.
Characters Per Inch	No more than 99	≤ 99

* Headings are to be at least 2px larger than Body Text.
 ** Drug Facts Title to be larger than all other type size used in DPP.
 † FDA Recommendations 2.5px min.

NO COPY

<p>Drug Facts (continued)</p> <p>Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.</p> <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>		<p>Drug Facts (continued)</p> <p>Other information</p> <ul style="list-style-type: none"> phenylketonurics: contains phenylalanine 1.4 mg per tablet safety sealed: do not use if the individual blister unit imprinted with Children's Claritin® is open or torn store between 20° to 25°C (68° to 77°F) <p>Inactive ingredients</p> <p>aspartame, citric acid anhydrous, colloidal silicon dioxide, D&C red No. 27 aluminum lake, FD&C blue No. 2 aluminum lake, flavor, magnesium stearate, mannitol, microcrystalline cellulose, sodium starch glycolate, stearic acid</p> <p>Questions or comments? 1-800-CLARITIN (1-800-252-7484) or www.claritin.com</p> <p><small>Bayer, the Bayer Cross, Claritin, Children's Claritin & Balloon Design and the Claritin Vista are registered trademarks of Bayer. © 2019 Bayer. Dist by: Bayer HealthCare LLC Whippany, NJ 07981 Product of Ireland. 00000000</small></p>									
<p>Directions</p> <p>chew or crush tablets completely before swallowing.</p> <table border="1"> <tr> <td>adults and children 6 years and over</td> <td>chew 2 tablets daily; not more than 2 tablets in 24 hours</td> </tr> <tr> <td>children 2 to under 6 years of age</td> <td>chew 1 tablet daily; not more than 1 tablet in 24 hours</td> </tr> <tr> <td>children under 2 years of age</td> <td>ask a doctor</td> </tr> <tr> <td>consumers with liver or kidney disease</td> <td>ask a doctor</td> </tr> </table>		adults and children 6 years and over	chew 2 tablets daily; not more than 2 tablets in 24 hours	children 2 to under 6 years of age	chew 1 tablet daily; not more than 1 tablet in 24 hours	children under 2 years of age	ask a doctor	consumers with liver or kidney disease	ask a doctor	Clear Area	
adults and children 6 years and over	chew 2 tablets daily; not more than 2 tablets in 24 hours										
children 2 to under 6 years of age	chew 1 tablet daily; not more than 1 tablet in 24 hours										
children under 2 years of age	ask a doctor										
consumers with liver or kidney disease	ask a doctor										

Sample - Not for Sale

Children's Claritin Chewables
loratadine chewable tablets
5 mg/antihistamine
Indoor & Outdoor Allergies
Grape Flavored
2 CHEWABLE TABLETS

*When taken as directed. See Drug Facts Panel.

Drug Facts	
Active ingredient (in each tablet)	Purpose
Loratadine 5 mg.....	Antihistamine
Uses	
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:	
<ul style="list-style-type: none"> runny nose itchy, watery eyes sneezing itching of the nose or throat 	
Warnings	
Do not use if you have ever had an allergic reaction to this product or any of its ingredients.	
Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.	
When using this product do not take more than directed. Taking more than directed may cause drowsiness.	

Actual Size 100%

FDA00150-3

DRUG FACTS INFO (updated 5/18/18)		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt **	9 pt
Drug Facts Continued	8 pt	8 pt
Headings*	8 pt-bold italic*	8 pt
Subheadings	≥6 pt	6 pt
Body Text	≥6 pt	6 pt
Bullets	5 pt	3.5 pt
Leading	≥0.5 pt	6.5 pt
Hairlines	0.5 pt rule	0.5 pt
Box and Barlines†	>0.5 pt rule	1 pt
Characters Per Inch:	No more than 39	≤ 39

* Headings are to be at least 2pts larger than Body Text.
** Drug Facts Title to be larger than all other type size used in DFP
† FDA Recommends 2.5pt rule



Drug Facts
Active ingredient (in each tablet) Purpose
 Loratadine 5 mg.....Antihistamine

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 ■ runny nose ■ itchy, watery eyes
 ■ sneezing ■ itching of the nose or throat

Warnings
Do not use if you have ever had an allergic reaction to this product or any of its ingredients.
Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.
When using this product do not take more than directed. Taking more than directed may cause drowsiness.
Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)
Directions
 ■ chew or crush tablets completely before swallowing.
 adults and children 6 years and over chew 2 tablets daily; not more than 2 tablets in 24 hours
 children 2 to under 6 years of age chew 1 tablet daily; not more than 1 tablet in 24 hours
 children under 2 years of age ask a doctor
 consumers with liver or kidney disease ask a doctor

Other information
 ■ phenylketonurics: contains phenylalanine 1.4 mg per tablet
 ■ safety sealed: do not use if the individual blister unit imprinted with Children's Claritin® is open or torn
 ■ store between 20° to 25°C (68° to 77°F)

Inactive ingredients
 aspartame, citric acid anhydrous, colloidal silicon dioxide, D&C red No. 27 aluminum lake, FD&C blue No. 2 aluminum lake, flavor, magnesium stearate, mannitol, microcrystalline cellulose, sodium starch glycolate, stearic acid

Questions or comments?
 1-800-CLARITIN (1-800-252-7484) or www.claritin.com

Actual Size 100%

FDA00056-D-3

DRUG FACTS INFO (updated 5/18/18)		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt. **	9 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings*	8 pt.-bold italic*	8 pt.
Subheadings	≥6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	3.5 pt.
Leading	≥0.5 pt.	6.5 pt.
Hairlines	0.5 pt. rule	0.5 pt.
Box and Barlines ¹	>0.5 pt. rule	1 pt.
Characters Per Inch:	No more than 39	≤ 39

* Headings are to be at least 2pts larger than Body Text.
 ** Drug Facts Title to be larger than all other type size used in DFP
 1 FDA Recommends 2.5pt rule

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

THERESA M MICHELE
06/17/2019 05:21:35 PM