

naproxen sodium capsules, 220 mg (NSAID) pain reliever/fever reducer

Do not use if printed safety seal under cap is torn or missing.

Drug Facts (continued)

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

Adults and children 12 years and older	<ul style="list-style-type: none"> Take 1 capsule every 8 hours to 12 hours while symptoms last for the first dose you may take 2 capsules within the first hour do not exceed 2 capsules in any 8-hour to 12-hour period do not exceed 3 capsules in a 24-hour period
Children under 12 years	ask a doctor

Other information

- each capsule contains: sodium 20 mg
- store at 20° to 25°C (68° to 77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- read all directions and warnings before use. Keep carton.

Inactive ingredients

FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments?
call toll-free 1-888-235-2466 (Mon - Fri 9AM - 5PM EST)

Manufactured for:
BIONPHARMA
600 Alexander Road,
Princeton, NJ 08540

L0000000
R1019

naproxen sodium capsules, 220 mg (NSAID) pain reliever/fever reducer

12 hour back & muscle pain relief

50 capsules**

NP1 Actual Size

***Compare to the active ingredient in ALEVE® BACK & MUSCLE PAIN

for temporary relief of minor back and muscle aches and pains

NDC 60452-269-16

Drug Facts

Active ingredient (in each capsule) Naproxen sodium 220 mg (naproxen 200 mg) (NSA D)*

Purposes Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - backache
 - minor pain of arthritis
 - headache
 - the common cold
 - temporarily reduces fever
 - muscular aches
 - menstrual cramps
 - toothache

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSA D, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

***This product is not manufactured or distributed by the owners of ALEVE® Back & Muscle pain.



Lot No.:
Exp. Date:

BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique Georgia			
Drug Facts:	8pt	Leading:	7pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	0.6pt
Body Text:	6pt	Hairlines:	0.2pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	

Drug Facts (continued)

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

Adults and children 12 years and older	<ul style="list-style-type: none"> Take 1 capsule every 8 hours to 12 hours while symptoms last for the first dose you may take 2 capsules within the first hour do not exceed 2 capsules in any 8-hour to 12-hour period do not exceed 3 capsules in a 24-hour period
Children under 12 years	<ul style="list-style-type: none"> ask a doctor

Other information

- each capsule contains: sodium 20 mg
- store at 20° to 25°C (68° to 77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- read all directions and warnings before use. Keep carton.

Inactive ingredients

FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments?
call toll-free 1-888-235-2466 (Mon - Fri 9AM - 5PM EST)

Manufactured for:
BIONPHARMA
600 Alexander Road,
Princeton, NJ 08540

L0000000
R1019

Drug Facts

Active ingredient (in each capsule)	Purposes
Naproxen sodium 220 mg (naproxen 200 mg) (NSA D)*	Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug	

Uses

- temporarily relieves minor aches and pains due to:
 - backache
 - minor pain of arthritis
 - headache
 - the common cold
 - temporarily reduces fever
 - muscular aches
 - menstrual cramps
 - toothache

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSA D, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Do not use if printed safety seal under cap is torn or missing.

naproxen sodium capsules, 220 mg (NSAID)
pain reliever/fever reducer

naproxen sodium capsules, 220 mg (NSAID)
pain reliever/fever reducer

12 hour back & muscle pain relief

Actual Size

50 capsules**

**Hybrid-filled capsule


for temporary relief of minor back and muscle aches and pains

NDC 60452-269-16



Lot No.:
Exp. Date:

BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique Georgia			
Drug Facts:	8pt	Leading:	7pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	0.6pt
Body Text:	6pt	Hairlines:	0.2pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	



BION PHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique Georgia			
Drug Facts:	NA	Leading:	5pt
Header:	5pt	Bullets:	3.5pt
Subheader:	4.5pt	Barlines:	NA
Body Text:	4.5pt	Hairlines:	NA
Drug Facts (continued):	NA		
Horizontal Scale: 100%		Average Kerning: 0%	

Front

.125" corner radius

***Compare to the active ingredient in ALEVE® BACK & MUSCLE PAIN

NDC 69452-269-20

naproxen sodium capsules, 220 mg (NSAID)
pain reliever/fever reducer

12 hour back & muscle pain relief

Actual Size

NP1

100 capsules**
**liquid-filled capsule

for temporary relief of minor back and muscle aches and pains

Do not use if printed safety seal under cap is torn or missing.

Drug Facts

Active ingredient (in each capsule)
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)* ...Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug

Purposes

Uses

- temporarily relieves minor aches and pains due to:
 - backache
 - muscular aches
 - minor pain of arthritis
 - menstrual cramps
 - headache
 - toothache
 - the common cold
 - temporarily reduces fever

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Manufactured for: **BIONPHARMA**
600 Alexander Road,
Princeton, NJ 08540

Lot/EXP: L000000 R1019

694521269206

Non Varnish Area

PEEL HERE

2.375"

6.00"

Back

Drug Facts (continued)

Ask a doctor before use if:

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are:

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit or aspirin
- taking any other drug

When using this product:

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if:

- you experience any of the following signs of stomach bleeding:
 - heart burn
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor, because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- if taken with food, this product may take longer to work

Adults and children 12 years and older:

- take 1 capsule every 8 hours to 12 hours while symptoms last
- for the first dose you may take 2 capsules within the first hour
- do not exceed 2 capsules in any 8-hour to 12-hour period
- do not exceed 3 capsules in a 24-hour period

Children under 12 years:

- ask a doctor

Other information

- each capsule contains: sodium 20 mg
- store at 20° to 25°C (68° to 77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- read all directions and warnings before use.

Inactive ingredients: FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments?
call toll-free 1-888-235-2466 (Mon - Fri 9AM - 5PM EST)

2.375"

6.00"

BIONPHARMA	
Labeling Format Information:	
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique Georgia	
Drug Facts:	8pt
Header:	7pt
Subheader:	6pt
Body Text:	6pt
Drug Facts (continued):	7pt
Leading:	5pt
Bullets:	5pt
Barlines:	0.6pt
Hairlines:	0.2pt
Horizontal Scale: 100%	
Average Kerning: 0%	

Front

.125" corner radius

Back

BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique Georgia			
Drug Facts:	8pt	Leading:	5pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	0.6pt
Body Text:	6pt	Hairlines:	0.2pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	



BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique Georgia			
Drug Facts:	8pt	Leading:	7pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	0.6pt
Body Text:	6pt	Hairlines:	0.2pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	



BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique Georgia			
Drug Facts:	8pt	Leading:	7pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	0.6pt
Body Text:	6pt	Hairlines:	0.2pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	

naproxen sodium capsules, 220 mg (NSAID)
pain reliever/fever reducer

Do not use if printed safety seal under cap is torn or missing.

Drug Facts

Active ingredient (in each capsule) Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug

Purposes

- temporarily relieves minor aches and pains due to:
 - menstrual cramps
 - muscular aches
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSA D, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSA Ds, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

naproxen sodium capsules, 220 mg (NSAID)
pain reliever/fever reducer

Actual Size
NP1

80 capsules**
**liquid-filled capsule

12 hour menstrual pain relief

temporarily relieves minor cramps, backache, and headache

***compare to the active ingredient in Menstridol™

NDC 69452-270-78

Drug Facts (continued)

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

under 12 years	ask a doctor
adults and children 12 years and older	take 1 capsule every 8 hours to 12 hours while symptoms last
	for the first dose you may take 2 capsules within the first hour
	do not exceed 2 capsules in any 8-hour to 12-hour period
	do not exceed 3 capsules in a 24-hour period

Other information

- each capsule contains: sodium 20 mg
- store at 20° to 25°C (68° to 77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- read all directions and warnings before use. Keep carton.

Inactive ingredients

FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments?
call toll-free 1-888-235-2466 (Mon - Fri 9AM - 5PM EST)

Manufactured for:
BIONPHARMA
600 Alexander Road,
Princeton, NJ 08540

L0000000
R1019

Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

***This product is not manufactured or distributed by the owners of Menstridol™.



Lot No.:
Exp. Date:

BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique Georgia			
Drug Facts:	8pt	Leading:	7pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	0.6pt
Body Text:	6pt	Hairlines:	0.2pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	



BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique Georgia			
Drug Facts:	8pt	Leading:	7pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	0.6pt
Body Text:	6pt	Hairlines:	0.2pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	



BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique Georgia			
Drug Facts:	8pt	Leading:	7pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	0.6pt
Body Text:	6pt	Hairlines:	0.2pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	

Do not use if printed safety seal under cap is torn or missing.

naproxen sodium capsules, 220 mg (NSAID)
pain reliever/fever reducer

Drug Facts (continued)

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- do not take more than directed
 - the smallest effective dose should be used
 - drink a full glass of water with each dose
 - if taken with food, this product may take longer to work
- | | |
|--|--|
| under 12 years | ask a doctor |
| adults and children 12 years and older | take 1 capsule every 8 hours to 12 hours while symptoms last |
| | for the first dose you may take 2 capsules within the first hour |
| | do not exceed 2 capsules in any 8-hour to 12-hour period |
| | do not exceed 3 capsules in a 24-hour period |

Other information

- each capsule contains: sodium 20 mg
- store at 20° to 25°C (68° to 77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- read all directions and warnings before use. Keep carton.

Inactive ingredients

FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments?

call toll-free 1-888 235-2466 (Mon - Fri 9AM - 5PM EST)

Manufactured for:
BIONPHARMA
600 Alexander Road,
Princeton, NJ 08540

L0000000
R1019

Drug Facts

Active ingredient (in each capsule)
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Purposes

Uses

- temporarily relieves minor aches and pains due to:
 - menstrual cramps
 - muscular aches
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

 If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSA Ds (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSA Ds, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

40 capsules**

liquid-filled capsule

NP1

Actual Size

12 hour menstrual pain relief

temporarily relieves minor cramps, backache, and headache

naproxen sodium capsules, 220 mg (NSAID)
pain reliever/fever reducer

NDC 69482-270-15



Lot No.:
Exp. Date:

naproxen sodium capsules, 220 mg (NSAID)
 pain reliever/fever reducer

Do not use if printed safety seal under cap is torn or missing.

Drug Facts

Active ingredient (in each capsule)	Purposes
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*	Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - menstrual cramps
 - headache
 - minor pain of arthritis
 - the common cold
 - muscular aches
 - backache
 - toothache
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSA D, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSA Ds, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

naproxen sodium capsules, 220 mg (NSAID)
 pain reliever/fever reducer

12 hour menstrual pain relief

temporarily relieves minor cramps, backache, and headache

Actual Size
 NP1

80 capsules**

**liquid-filled capsule

NDC 69452-270-78

Drug Facts (continued)

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

under 12 years	ask a doctor
adults and children 12 years and older	<ul style="list-style-type: none"> take 1 capsule every 8 hours to 12 hours while symptoms last for the first dose you may take 2 capsules within the first hour do not exceed 2 capsules in any 8-hour to 12-hour period do not exceed 3 capsules in a 24-hour period

Other information

- each capsule contains: sodium 20 mg
- store at 20° to 25°C (68° to 77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- read all directions and warnings before use. Keep carton.

Inactive ingredients

FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments?
 call toll-free 1-888-235-2466 (Mon - Fri 9AM - 5PM EST)

Manufactured for:
BIONPHARMA
 600 Alexander Road,
 Princeton, NJ 08540

L0000000
 R1019

Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear


If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.



Lot No.:
 Exp. Date:

BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique Georgia			
Drug Facts:	8pt	Leading:	7pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	0.6pt
Body Text:	6pt	Hairlines:	0.2pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	



BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique Georgia			
Drug Facts:	NA	Leading:	5pt
Header:	5pt	Bullets:	3.5pt
Subheader:	4.5pt	Barlines:	NA
Body Text:	4.5pt	Hairlines:	NA
Drug Facts (continued): NA			
Horizontal Scale: 100%		Average Kerning: 0%	

2.375"

NDC 69452-270-78

naproxen sodium capsules, 220 mg (NSAID)

pain reliever/fever reducer

12 hour menstrual pain relief

temporarily relieves minor cramps, backache, and headache

80 capsules**

**liquid-filled capsule

Do not use if printed safety seal under cap is torn or missing. KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Warnings Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), stork, skin reddening, rash, or blisters. † An allergic reaction occurs. stop use and seek medical help right away. **Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: ● are age 60 or older ● have had stomach ulcers or bleeding problems ● take a blood thinning (anticoagulant) or steroid drug ● take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ● have 3 or more alcoholic drinks every day while using this product ● take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer.**

Ask a doctor or pharmacist before use if you are ● under a doctor's care for any serious condition ● taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin ● taking any other drug. **Stop use and ask a doctor if you experience any of the following signs of stomach bleeding (feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better), have symptoms of heart problems or stroke (chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling), or if pain or fever gets worse. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions Under 12 years: ask a doctor. Adults and children 12 years and older: take 1 capsule with water every 8 hours to 12 hours while symptoms last. For the first dose you may take 2 capsules within the first hour. Do not exceed 3 capsules in a 24-hour period. ask a doctor.

Questions or comments? call toll-free 1-888-235-2466 (Mon - Fri 9AM - 5PM EST)

Non Varnish Area

L000000
R1019
LotEXP

.125" corner radius

6.00"

BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique Georgia			
Drug Facts:	NA	Leading:	7.2pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	NA
Body Text:	6pt	Hairlines:	NA
Drug Facts (continued):	NA		
Horizontal Scale: 100%		Average Kerning: 0%	

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

KAREN M MAHONEY
10/26/2019 09:21:31 PM