

N 0 00000-00000 0

- **Unflavored**
- **Sugar Free**
- **Dissolves in any Beverage**

Drug Facts	Drug Facts (continued)
Active ingredient (in each dose) Polyethylene Glycol 3350, 17 g.....Osmotic Laxative	Purpose If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Use ■ relieves occasional constipation (irregularity) ■ generally produces a bowel movement in 1 to 3 days	Directions ■ do not take more than directed unless advised by your doctor ■ adults and children 17 years of age and older: ■ stir and dissolve one packet of powder (17 g) in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink ■ ensure that the powder is fully dissolved before drinking ■ do not drink if there are any clumps ■ use once a day ■ use no more than 7 days ■ children 16 years of age or under: ask a doctor
Warnings Allergy alert: Do not use if you are allergic to polyethylene glycol Do not use if you have kidney disease, except under the advice and supervision of a doctor Ask a doctor before use if you have ■ nausea, vomiting or abdominal pain ■ a sudden change in bowel habits that lasts over 2 weeks ■ irritable bowel syndrome Ask a doctor or pharmacist before use if you are taking a prescription drug When using this product you may have loose, watery, more frequent stools Stop use and ask a doctor if ■ you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition. ■ you get diarrhea ■ you need to use a laxative for longer than 1 week	Other information ■ store at 20° - 25°C (68° - 77°F) ■ tamper-evident: do not use if foil is open or broken Inactive ingredients none Questions or comments? 1-800-MiraLAX (1-800-647-2529)

Bayer, the Bayer Cross, Mix-In Pax and MiraLAX are registered trademarks of Bayer.
© 2019 Bayer.
Dist by: Bayer HealthCare LLC, Whippany, NJ 07981
00000

Bayer

FOLD AT LINE
TEAR AT ARROW

#1 DOCTOR RECOMMENDED BRAND

MiraLAX[®]

Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax[®] Travel Size

• Relieves Occasional Constipation/Irregularity
• Softens Stool

UNFLAVORED
POWDER
GRIT FREE

1 DOSE

NET WT 0.5 OZ (17 g)

Actual Size 100%

FDA0006

DRUG FACTS INFO		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt.**	9 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings*	8 pt:bold italic*	8 pt.
Subheadings	≥6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	3.5 pt.
Leading	≥0.5 pt.	.5 pt.
Hairlines	0.5 pt rule	.5 pt.
Box and Barlines†	>0.5 pt rule	1 pt.
Characters Per Inch:	No more than 39	≥39

* Headings are to be at least 2pts larger than Body Text.
** Drug Facts Title to be larger than all other type size used in DFP
† FDA Recommends 2.5pt rule

Sample-Not For Sale

- **Unflavored**
- **Sugar Free**
- **Dissolves in any Beverage**

Drug Facts		Drug Facts (continued)
Active ingredient (in each dose) Polyethylene Glycol 3350, 17 g.....Osmotic Laxative	Purpose	If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Use		Directions
<ul style="list-style-type: none"> relieves occasional constipation (irregularity) generally produces a bowel movement in 1 to 3 days 		<ul style="list-style-type: none"> do not take more than directed unless advised by your doctor adults and children 17 years of age and older: <ul style="list-style-type: none"> stir and dissolve one packet of powder (17 g) in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink ensure that the powder is fully dissolved before drinking do not drink if there are any clumps use once a day use no more than 7 days children 16 years of age or under: ask a doctor
Warnings		Other information
<p>Allergy alert: Do not use if you are allergic to polyethylene glycol</p> <p>Do not use if you have kidney disease, except under the advice and supervision of a doctor</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> nausea, vomiting or abdominal pain a sudden change in bowel habits that lasts over 2 weeks irritable bowel syndrome <p>Ask a doctor or pharmacist before use if you are taking a prescription drug</p> <p>When using this product you may have loose, watery, more frequent stools</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition. you get diarrhea you need to use a laxative for longer than 1 week 		<ul style="list-style-type: none"> store at 20° - 25°C (68° - 77°F) tamper-evident: do not use if foil is open or broken
		Inactive ingredients none
		Questions or comments? 1-800-MiraLAX (1-800-647-2529)
		<p>Bayer, the Bayer Cross, Mix-In Pax and MiraLAX are registered trademarks of Bayer. © 2019 Bayer. Dist by: Bayer HealthCare LLC, Whippany, NJ 07981 00000</p> <p style="text-align: right;">Bayer</p>



Actual Size 100%

FDA0004

DRUG FACTS INFO		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt **	9 pt
Drug Facts Continued	8 pt	8 pt
Headings*	8 pt-bold italic*	8 pt
Subheadings	≥6 pt	6 pt
Body Text	≥6 pt	6 pt
Bullets	5 pt	3.5 pt
Leading	≥0.5 pt	.5 pt
Hairlines	0.5 pt rule	.5 pt
Box and Barlines†	>0.5 pt rule	1 pt
Characters Per Inch:	No more than 39	≥39

* Headings are to be at least 2pts larger than Body Text.
** Drug Facts Title to be larger than all other type size used in DFP
† FDA Recommends 2.5pt rule

Not For Individual Sale

- **Unflavored**
- **Sugar Free**
- **Dissolves in Any Beverage**

Drug Facts		Drug Facts (continued)	
Active ingredient (in each dose) Polyethylene Glycol 3350, 17 g.....Osmotic Laxative	Purpose	if pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Use <ul style="list-style-type: none"> relieves occasional constipation (irregularity) generally produces a bowel movement in 1 to 3 days 		Directions <ul style="list-style-type: none"> do not take more than directed unless advised by your doctor adults and children 17 years of age and older: <ul style="list-style-type: none"> stir and dissolve one packet of powder (17 g) in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink ensure that the powder is fully dissolved before drinking do not drink if there are any clumps use once a day use no more than 7 days children 16 years of age or under: ask a doctor 	
Warnings Allergy alert: Do not use if you are allergic to polyethylene glycol. Do not use if you have kidney disease, except under the advice and supervision of a doctor Ask a doctor before use if you have <ul style="list-style-type: none"> nausea, vomiting or abdominal pain a sudden change in bowel habits that lasts over 2 weeks irritable bowel syndrome Ask a doctor or pharmacist before use if you are taking a prescription drug When using this product you may have loose, watery, more frequent stools Stop use and ask a doctor if <ul style="list-style-type: none"> you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition. you get diarrhea you need to use a laxative for longer than 1 week 		Other information <ul style="list-style-type: none"> store at 20° - 25°C (68° - 77°F) tamper-evident: do not use if foil is open or broken Inactive ingredients none Questions or comments? 1-800-MiraLAX (1-800-647-2529) Bayer, the Bayer Cross, Mix-In Pax and MiraLAX are registered trademarks of Bayer. © 2019 Bayer. Dist by: Bayer HealthCare LLC, Whippany, NJ 07981 00000 	



Actual Size 100%

FDA00005

DRUG FACTS INFO		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt. **	9 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings [†]	8 pt.-bold italic [‡]	8 pt.
Subheadings	≥6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	3.5 pt.
Leading	≥0.5 pt.	.5 pt.
Hairlines	0.5 pt. rule	.5 pt.
Box and Barlines [†]	>0.5 pt. rule	1 pt.
Characters Per Inch:	No more than 39	≥39

^{*} Headings are to be at least 2pts larger than Body Text.
^{**} Drug Facts Title to be larger than all other type size used in DFP
[†] FDA Recommends 2.5pt rule

PRE-MEASURED,
SINGLE DOSES

3

MiraLAX[®]
Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax[®]

Drug Facts

Active ingredient (in each dose) Polyethylene Glycol 3350, 17 g
Purpose Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol. Do not use if you have kidney disease, except under the advice and supervision of a doctor.

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

Drug Facts (continued)

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed unless advised by your doctor
- adults and children 17 years of age and older:
 - stir and dissolve one packet of powder (17 g) in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
 - ensure that the powder is fully dissolved before drinking
 - do not drink if there are any clumps
 - use once a day
 - use no more than 7 days
 - children 16 years of age or under: ask a doctor

Other information

- store at 20° - 25°C (68° - 77°F)
- tamper-evident: do not use if foil is open or broken

Inactive ingredients none

Questions or comments?

1-800-MiraLAX (1-800-647-2529)

Bayer, the Bayer Cross, Mix-In Pax and MiraLAX are registered trademarks of Bayer.

© 2019 Bayer. Dist by: Bayer HealthCare LLC, Whippany, NJ 07981

Bayer

00000



Mix-In Pax[®]

MiraLAX[®]
Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Travel Size

3
Single Doses




MiraLAX[®]

Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax[®]

- Relieves Occasional Constipation/Irregularity
- Softens Stool

3
ONCE-DAILY DOSES



Unflavored Powder Dissolves in ANY Beverage!

UNFLAVORED
POWDER
GRIT FREE

3 PACKETS
NET WT 0.5 OZ
(17g) EACH

Mix-In Pax[®]
Travel Size

MiraLAX[®]
Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Actual Size 100%

FDA00011

DRUG FACTS INFO		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt. **	9 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings*	8 pt.-bold italic*	8 pt.
Subheadings	≥6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	5 pt.
Leading	≥0.5 pt.	6.5 pt.
Hairlines	0.5 pt. rule	0.5 pt.
Box and Barlines†	>0.5 pt. rule	2.5 pt.
Characters Per Inch:	No more than 39	≤39

* Headings are to be at least 2pts larger than Body Text.
** Drug Facts Title to be larger than all other type size used in DFP
† FDA Recommends 2.5pt rule



Actual Size 100%

FDA00033

DRUG FACTS INFO	REQUIRED	ACTUAL IN FILE
Category	>8 pt **	8 pt
Drug Facts Title	>8 pt **	8 pt
Drug Facts Continued	8 pt	8 pt
Headings*	8 pt-bold italic*	8 pt
Subheadings*	≥6 pt	6 pt
Body Text	≥6 pt	6 pt
Bullets	5 pt	5 pt
Leading	≥0.5 pt	0.5 pt
Headlines	0.5 pt rule	0.5 pt
Box and Borderline	>0.5 pt rule	2.5 pt
Characters Per Inch:	No more than 38	≤38

* Headings are to be at least 2pt larger than Body Text.
 ** Drug Facts Title to be larger than all other type sizes used in DFP
 † FDA Recommendation: 6-8pt rule



Drug Facts
Active ingredient (in each dose) Polyethylene Glycol 3350, 17 g
Purpose Osmotic Laxative
Use
 ■ relieves occasional constipation (irregularity)
 ■ generally produces a bowel movement in 1 to 3 days
Warnings
Allergy alert: Do not use if you are allergic to polyethylene glycol
 Do not use if you have kidney disease, except under the advice and supervision of a doctor.
Ask a doctor before use if you have
 ■ nausea, vomiting or abdominal pain
 ■ a sudden change in bowel habits that lasts over 2 weeks
 ■ irritable bowel syndrome
Ask a doctor or pharmacist before use if you are taking a prescription drug
When using this product you may have loose, watery, more frequent stools.
Stop use and ask a doctor if
 ■ you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
 ■ you get diarrhea
 ■ you need to use a laxative for longer than 1 week

Drug Facts (continued)
 If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
 ■ do not take more than directed unless advised by your doctor
 ■ adults and children 17 years of age and older:
 ■ stir and dissolve one packet of powder (17 g) in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
 ■ ensure that the powder is fully dissolved before drinking
 ■ do not drink if there are any clumps
 ■ use once a day
 ■ use no more than 7 days
 ■ children 16 years of age or under: ask a doctor
Other information
 ■ store at 20° - 25°C (68° - 77°F)
 ■ tamper-evident: do not use if foil is open or broken
Inactive ingredients none
Questions or comments?
 1-800-MiraLAX (1-800-647-2529)
 Bayer, the Bayer Cross, Mix-In Pax and MiraLAX are registered trademarks of Bayer.
 © 2019 Bayer
 Dist by: Bayer HealthCare LLC, Whippany, NJ 07981 00000

MiraLAX
 Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative
Mix-In Pax
 • Unflavored
 • Sugar Free
 • Dissolves in Any Beverage
 10 PRE-MEASURED, SINGLE DOSES
 NET WT 0.5 OZ (17g) EACH

MiraLAX
 Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative
Mix-In Pax
 • Relieves Occasional Constipation/Irregularity
 • Softens Stool
 10 SINGLE DOSES
 UNFLAVORED POWDER GRIT FREE
 10 PACKETS NET WT 0.5 OZ (17g) EACH
 BAYER
 Unflavored Powder Dissolves in ANY Beverage!

MiraLAX
 Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative
Mix-In Pax
 Travel Size

Lot & Expiry (Non - Varnish Area)



Actual Size 100%

FDA00032

DRUG FACTS INFO		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt. ^{1,2}	9 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings ³	8 pt.-bold italic ³	8 pt.
Subheadings	≥5 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	5 pt.
Leading	≥0.5 pt.	6.5 pt.
Headlines	0.5 pt. rule	0.5 pt.
Box and Barlines ⁴	>0.5 pt. rule	2.5 pt.
Characters Per Inch:	No more than 39	≤39

¹ Headings are to be at least 2pts larger than Body Text.
² Drug Facts Title to be larger than all other type size used in DFP.
³ FDA recommends 2.5pt rule.

MiraLAX[®]
Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax[®]

MiraLAX[®]
Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax[®]

- Unflavored
- Sugar Free
- Dissolves in Any Beverage



20
Single Doses

MiraLAX[®]
Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax[®]

- Relieves Occasional Constipation/Irregularity
- Softens Stool

20
ONCE-DAILY DOSES



UNFLAVORED
POWDER
GRIT FREE

20 PACKETS
NET WT 0.5 OZ
(17g) EACH

#1
DOCTOR
RECOMMENDED
BRAND

MiraLAX[®]
Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax[®]



Travel Size

20
PRE-MEASURED, UNFLAVORED,
SINGLE DOSES

Drug Facts

Active ingredient (in each dose)
Polyethylene Glycol 3350, 17 g

Purpose
Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol.

Do not use if you have kidney disease, except under the advice and supervision of a doctor.

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools.

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed unless advised by your doctor
- adults and children 17 years of age and older:
 - stir and dissolve one packet of powder (17 g) in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
 - ensure that the powder is fully dissolved before drinking
 - do not drink if there are any clumps
 - use once a day
 - use no more than 7 days
 - children 16 years of age or under: ask a doctor

Other information

- store at 20° - 25°C (68° - 77°F)
- tamper-evident: do not use if foil is open or broken

Inactive ingredients none

Questions or comments?
1-800-MiraLAX (1-800-647-2529)

Bayer, the Bayer Cross, Mix-In Pax and MiraLAX are registered trademarks of Bayer.
© 2019 Bayer.
Distributed by Bayer HealthCare LLC, Whippany, NJ 07981
00000

Lot & Expiry
(Non - Varnish Area)
1.75"W X .75"H



Bayer

Actual Size 100%

FDA00025

DRUG FACTS INFO		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>= 8 pt. ^{1*}	9 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings ²	8 pt.-bold italic ²	8 pt.
Subheadings	>= 6 pt.	6 pt.
Body Text	>= 6 pt.	6 pt.
Bullets	5 pt.	5 pt.
Leading	>= 0.5 pt.	6.5 pt.
Headlines	0.5 pt. rule	0.5 pt.
Box and Barlines ³	>= 0.5 pt. rule	2.5 pt.
Character's Per Inch	No more than 39	≤39

¹ Headings are to be at least 2 pts. larger than Body Text.
² Drug Facts Title to be larger than all other type size used in ODP.
³ FDA Recommendations: 2.5 pt. rule.



Drug Facts
Active ingredient (in each dose) Polyethylene Glycol 3350, 17 g
Purpose Osmotic Laxative
Use
 ■ relieves occasional constipation (irregularity)
 ■ generally produces a bowel movement in 1 to 3 days
Warnings
Allergy alert: Do not use if you are allergic to polyethylene glycol.
Do not use if you have kidney disease, except under the advice and supervision of a doctor.
Ask a doctor before use if you have
 ■ nausea, vomiting or abdominal pain
 ■ a sudden change in bowel habits that lasts over 2 weeks
 ■ irritable bowel syndrome
Ask a doctor or pharmacist before use if you are taking a prescription drug.
When using this product you may have loose, watery, more frequent stools.
Stop use and ask a doctor if
 ■ you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
 ■ you get diarrhea
 ■ you need to use a laxative for longer than 1 week

Drug Facts (continued)
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
 ■ do not take more than directed unless advised by your doctor
 ■ adults and children 17 years of age and older:
 ■ stir and dissolve one packet of powder (17 g) in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
 ■ ensure that the powder is fully dissolved before drinking
 ■ do not drink if there are any clumps
 ■ use once a day
 ■ use no more than 7 days
 ■ children 16 years of age or under: ask a doctor
Other information
 ■ store at 20° - 25°C (68° - 77°F)
 ■ tamper-evident: do not use if foil is open or broken
Inactive ingredients none
Questions or comments?
 1-800-MiraLAX (1-800-647-2529)

Lot & Expiry (Non - Varnish Area)
 1.75"W X .75"H



Bayer, the Bayer Cross, Mix-In Pax and MiraLAX are registered trademarks of Bayer.
 © 2019 Bayer
 Distributed by Bayer HealthCare LLC, Whippany, NJ 07981
 000000

Bayer

MiraLAX
 Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax

- Unflavored
- Sugar Free
- Dissolves in Any Beverage



MiraLAX
 Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax

20 Single Doses

MiraLAX
 Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax

- Relieves Occasional Constipation/Irregularity
- Softens Stool

20 ONCE-DAILY DOSES



UNFLAVORED POWDER GRIT FREE

20 PACKETS
 NET WT 0.5 OZ (17g) EACH

#1 DOCTOR RECOMMENDED BRAND

MiraLAX
 Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax



Travel Size

20 PRE-MEASURED, UNFLAVORED, SINGLE DOSES

Actual Size 100%

FDA00024

CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt **	9 pt
Drug Facts Continued	8 pt	8 pt
Headings*	6 pt bold italic*	8 pt
Subheadings	>6 pt	6 pt
Body Text	>6 pt	6 pt
Bullets	5 pt	5 pt
Leading	>0.5 pt	0.5 pt
Headlines	0.5 pt rule	0.5 pt
Box and Barlines†	>0.5 pt rule	2.5 pt
Characters Per Inch:	No more than 35	<39

* Headings are to be at least 20% larger than body text.
 ** Drug Facts Title to be larger than all other type size used in DFP
 † FDA Recommendations 2.5pt rule



MiraLAX[®]
Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax[®]

24
Single Doses

#1
DOCTOR
RECOMMENDED
BRAND

MiraLAX[®]
Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax[®]

- Relieves Occasional Constipation/Irregularity
- Softens Stool

24 ONCE-DAILY DOSES



Unflavored Powder
Dissolves in ANY
Beverage!

UNFLAVORED
POWDER
GRIT FREE

24 PACKETS
NET WT 0.5 OZ.
(17g) EACH

MiraLAX[®]
Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax[®]



Travel Size

CONTAINS 24 ONCE-DAILY DOSES
EACH PACKET CONTAINS ONE DOSE, NET WT 0.5 OZ (17g)

MiraLAX[®]
Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax[®]

Drug Facts

Active ingredient (in each dose) Purpose
Polyethylene Glycol 3350, 17 g.....Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol.

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

■ **do not take more than directed unless advised by your doctor**

■ adults and children 17 years of age and older:

■ stir and dissolve one packet of powder (17 g) in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink

■ ensure that the powder is fully dissolved before drinking

■ do not drink if there are any clumps

■ use once a day

■ use no more than 7 days

■ children 16 years of age or under: ask a doctor

Other information

■ store at 20° - 25° C (68° - 77°F)

■ tamper-evident: do not use if foil is open or broken

Inactive ingredients none

Questions or comments?

1-800-MiraLAX (1-800-647-2529)

Bayer, the Bayer Cross, Mix-In Pax and MiraLAX are registered trademarks of Bayer.
© 2019 Bayer.
Dist by: Bayer HealthCare LLC, Whippany, NJ 07981
00000

Bayer

No Varnish

No Varnish
No Copy
Lot/Exp Area

MiraLAX[®]
Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax[®]

• Unflavored • Sugar Free • Dissolves in Any Beverage



CONTAINS 24 ONCE-DAILY DOSES
EACH PACKET CONTAINS ONE DOSE, NET WT 0.5 OZ (17g)

OPEN HERE

24
ONCE-DAILY DOSES

24 PACKETS
NET WT 0.5 OZ (17g)

Actual Size 100%

FDA00007

DRUG FACTS INFO		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt. **	8 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings*	8 pt bold italic†	8 pt.
Subheadings	>8 pt.	7 pt.
Body Text	>8 pt.	7 pt.
Bullets	5 pt.	5 pt.
Leading	>0.5 pt.	7.5 pt.
Headlines	0.5 pt rule	5 pt.
Box and Barlines†	>0.5 pt rule	2.5 pt.
Characters Per Inch	No more than 30	< 30

* Headings are to be at least 20pt larger than Body Text.
** Drug Facts Title to be larger than all other text used in DFP
† FDA Recommends 2.5pt rule

MiraLAX
Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax

PATIENT SAMPLES NOT FOR SALE

#1 DOCTOR RECOMMENDED BRAND

MiraLAX
Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax

- Relieves Occasional Constipation/Irregularity
- Softens Stool

24 ONCE-DAILY DOSES

UNFLAVORED POWDER GRIT FREE

24 PACKETS NET WT 0.5 OZ (17g) EACH

Unflavored Powder Dissolves in ANY Beverage!

BAYER

MiraLAX
Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax

PATIENT SAMPLES NOT FOR SALE

Travel Size

CONTAINS 24 ONCE-DAILY DOSES
EACH PACKET CONTAINS ONE DOSE, NET WT 0.5 OZ (17g)

MiraLAX
Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax

PATIENT SAMPLES NOT FOR SALE

Drug Facts

Active ingredient (in each dose) Purpose
Polyethylene Glycol 3350, 17 g Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed unless advised by your doctor
- adults and children 17 years of age and older: stir and dissolve one packet of powder (17 g) in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
- ensure that the powder is fully dissolved before drinking
- do not drink if there are any clumps
- use once a day
- use no more than 7 days
- children 16 years of age or under: ask a doctor

Other information

- store at 20° - 25°C (68° - 77°F)
- tamper-evident: do not use if foil is open or broken

Inactive ingredients none

Questions or comments?
1-800-MiraLAX (1-800-647-2529)

Bayer, the Bayer Cross, Mix-In Pax and MiraLAX are registered trademarks of Bayer.
© 2019 Bayer.
Dist. by: Bayer HealthCare LLC, Whippany, NJ 07981
00000

MiraLAX
Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax

PATIENT SAMPLES NOT FOR SALE

Unflavored • Sugar Free • Dissolves in Any Beverage

CONTAINS 24 ONCE-DAILY DOSES
EACH PACKET CONTAINS ONE DOSE, NET WT 0.5 OZ (17g)

24 ONCE-DAILY DOSES

24 PACKETS NET WT 0.5 OZ (17g)

OPEN HERE

No Varnish
No Copy
Lot/Exp Area

Actual Size 100%

CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	≥6 pt **	8 pt
Drug Facts Continued	8 pt	8 pt
Headings†	8 pt bold italic‡	8 pt
Subheadings	≥6 pt	7 pt
Body Text	≥6 pt	7 pt
Bullets	5 pt	5 pt
Leading	≥0.5 pt	7.5 pt
Headlines	0.5 pt rule	5 pt
Box and Barlines†	≥0.5 pt rule	2.5 pt
Characters Per Inch	No more than 35	≤ 35

† Headings are to be at least 6pt larger than Body Text.
‡ Drug Facts Title to be larger than all other text used in DDP
‡ FDA Recommendation: 2.5pt rule

FDA0008

Drug Facts
Active ingredient (in each dose) Polyethylene Glycol 3350, 17 g
Purpose Osmotic Laxative
Use
 ■ relieves occasional constipation (irregularity)
 ■ generally produces a bowel movement in 1 to 3 days
Warnings
Allergy alert: Do not use if you are allergic to polyethylene glycol.
Do not use if you have kidney disease, except under the advice and supervision of a doctor.
Ask a doctor before use if you have:
 ■ nausea, vomiting or abdominal pain
 ■ a sudden change in bowel habits that lasts over 2 weeks
 ■ irritable bowel syndrome
Ask a doctor or pharmacist before use if you are taking a prescription drug.
When using this product you may have loose, watery, more frequent stools.
Stop use and ask a doctor if:
 ■ you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
 ■ you get diarrhea
 ■ you need to use a laxative for longer than 1 week

Drug Facts (continued)
If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
 ■ do not take more than directed unless advised by your doctor
 ■ adults and children 17 years of age and older:
 ■ stir and dissolve one packet of powder (17 g) in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
 ■ ensure that the powder is fully dissolved before drinking
 ■ do not drink if there are any clumps
 ■ use once a day
 ■ use no more than 7 days
 ■ children 18 years of age or under: ask a doctor
Other information
 ■ store at 20°–25°C (68°–77°F)
 ■ tamper-evident: do not use if foil is open or broken
Inactive ingredients none
Questions or comments?
 1-800-MiraLAX (1-800-647-2525)

Bayer, the Bayer Cross, Mix-In Pax and MiraLAX are registered trademarks of Bayer.
 © 2019 Bayer.
 Dist by: Bayer HealthCare LLC, Whippany, NJ 07981
 00000



Bayer

Lot Code/Exp
 1.75" x 3/4"

MiraLAX[®]
 Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax[®]

- Unflavored
- Sugar Free
- Dissolves in Any Beverage




MiraLAX[®]
 Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax[®]

- Relieves Occasional Constipation/Irregularity
- Softens Stool

40 ONCE-DAILY DOSES



Twin Pack
 2 x 20 Doses



MiraLAX[®]
 Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax[®]

- Relieves Occasional Constipation/Irregularity
- Softens Stool



2x20 COUNT CARTONS
 40 PACKETS TOTAL
 NET WT 0.5 OZ (17g) EACH



MiraLAX[®]
 Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax[®]



Travel Size

40 PRE-MEASURED, UNFLAVORED, SINGLE DOSES

Actual Size 100%

FDA00034







DRUG FACTS INFO		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>6 pt **	9 pt
Drug Facts Continued	8 pt	8 pt
Headings*	8 pt-bold italic*	8 pt
Subheadings	>6 pt	6 pt
Body Text	>6 pt	6 pt
Bullets	5 pt	5 pt
Leading	>0.5 pt	6.5 pt
Headlines	0.5 pt rule	0.5 pt
Box and Barlines†	>0.5 pt rule	2.5 pt
Characters Per Inch	No more than 36	<36

* Headings are to be set in size larger than Body Text.
 ** Drug Facts Title to be larger than all other type size used in DFP.
 † FDA recommends 2.5 pt rule.



Actual Size 100%

FDA00009



Actual Size 100%

FDA00010

DRUG FACTS INFO		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt **	9 pt
Drug Facts Continued	8 pt	8 pt
Headings*	8 pt-bold italic*	8 pt
Subheadings	≥6 pt	6 pt
Body Text	≥6 pt	6 pt
Bullets	5 pt	5 pt
Leading	≥0.5 pt	6.5 pt
Hairlines	0.5 pt rule	.5 pt
Box and Barlines†	>0.5 pt rule	2.5 pt
Characters Per Inch:	No more than 39	≤ 39

* Headings are to be at least 2xts larger than Body Text.
 ** Drug Facts Title to be larger than all other type size used in DFP
 † FDA Recommends 2.5pt rule



MiraLAX® increases frequency of bowel movements and softens the stool.

XXXXXXXX

Lot Expiry Area

MiraLAX®


Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative



• Dissolves in Any Beverage

Bayer, the Bayer Cross, and MiraLAX are registered trademarks of Bayer. © 2019 Bayer. Dist by: Bayer HealthCare LLC Whippany, NJ 07981

Bayer



UNFLAVORED
POWDER
GRIT FREE

- Relieves Occasional Constipation/Irregularity
- Softens Stool



14 ONCE-DAILY DOSES

NET WT 8.3 OZ (238 g)

Actual Size 100%

FDA00012

Peel here

Drug Facts

Active ingredient (in each dose)	Purpose
Polyethylene Glycol 3350, 17 g (cap filled to line)	Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

• Sugar Free

XXXXXX

Drug Facts (continued)

- use once a day
- use no more than 7 days
- children 16 years of age or under: ask a doctor

Other information

- store at 20° - 25°C (68° - 77°F)
- tamper-evident: do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken

Inactive ingredients none

Questions or comments?
1-800-MiraLAX (1-800-647-2529)

Drug Facts (continued)

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed unless advised by your doctor
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
 - fill to top of white section in cap which is marked to indicate the correct dose (17g)
 - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
 - ensure that the powder is fully dissolved before drinking
 - do not drink if there are any clumps

Actual Size 100%

FDA00013

DRUG FACTS INFO		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt. **	9 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings*	8 pt.-bold italic*	8 pt.
Subheadings	≥6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	5 pt.
Leading	≥0.5 pt.	6.5 pt.
Hairlines	0.5 pt rule	0.5 pt.
Box and Barlines†	>0.5 pt rule	1.5 and 2.5 pt.
Characters Per Inch:	No more than 39	≤39

* Headings are to be at least 2pts larger than Body Text.
 ** Drug Facts Title to be larger than all other type size used in DFP
 † FDA Recommends 2.5pt rule



• *Dissolves in Any Beverage*

XXXXXXXX

Lot
Expiry Area



UNFLAVORED
POWDER
GRIT FREE

- Relieves Occasional Constipation/Irregularity
- Softens Stool

30 ONCE-DAILY DOSES


NET WT 17.9 OZ (510g)

#1
DOCTOR
RECOMMENDED
BRAND

MiraLAX® increases frequency of bowel movements and softens the stool.

Bayer, the Bayer Cross, and MiraLAX are registered trademarks of Bayer.
© 2019 Bayer.
Dist by: Bayer HealthCare LLC
Whippany, NJ 07981

Bayer



Actual Size 100%

FDA00016

Drug Facts

Active ingredient (in each dose) Purpose

Polyethylene Glycol 3350,
17 g (cap filled to line).....Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed unless advised by your doctor**
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
 - fill to top of white section in cap which is marked to indicate the correct dose (17g)
 - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
 - ensure that the powder is fully dissolved before drinking
- do not drink if there are any clumps
- use once a day
- use no more than 7 days
- children 16 years of age or under: ask a doctor

Other information

- store at 20° - 25°C (68° - 77°F)
- tamper-evident: do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken

Inactive ingredients none

Questions or comments?

1-800-MiraLAX (1-800-647-2529)

• Sugar Free

XXXXXX

Actual Size 100%

FDA00017

DRUG FACTS INFO		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt **	9 pt
Drug Facts Continued	8 pt	8 pt
Headings†	8 pt:bold italic*	8 pt
Subheadings	≥6 pt	6 pt
Body Text	≥6 pt	6 pt
Bullets	5 pt	5 pt
Leadings	≥0.5 pt	6.5 pt
Hairlines	0.5 pt:rule	0.5 pt
Box and Barlines‡	>0.5 pt:rule	1.5 and 2.5 pt
Characters Per Inch:	No more than 39	≤39

* Headings are to be at least 2pts larger than Body Text.
 ** Drug Facts Title to be larger than all other type size used in DFP
 † FDA Recommends 2.5pt:rule



Actual Size 100%

FDA00018

Drug Facts

Active ingredient (in each dose) Purpose
Polyethylene Glycol 3350,
17 g (cap filled to line).....Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed unless advised by your doctor**
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
 - fill to top of white section in cap which is marked to indicate the correct dose (17g)
 - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
 - ensure that the powder is fully dissolved before drinking
- do not drink if there are any clumps
- use once a day
- use no more than 7 days
- children 16 years of age or under: ask a doctor

Other information

- store at 20° - 25°C (68° - 77°F)
- tamper-evident: do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken

Inactive ingredients none

Questions or comments?
1-800-MiraLAX (1-800-647-2529)

• Sugar Free

000000

Actual Size 100%

FDA00026

DRUG FACTS INFO		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt. **	9 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings*	8 pt.-bold italic*	8 pt.
Subheadings	≥6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	5 pt.
Leading	≥0.5 pt.	6.5 pt.
Hairlines	0.5 pt. rule	0.5 pt.
Box and Barlines†	>0.5 pt. rule	1.5 and 2.5 pt.
Characters Per Inch:	No more than 39	≤39


* Headings are to be at least 2pts larger than Body Text.
 ** Drug Facts Title to be larger than all other type size used in DFP
 † FDA Recommends 2.5pt rule



Actual Size 100%

FDA00020

• Sugar Free

Drug Facts		Drug Facts (continued)	
Active ingredient (in each dose)	Purpose	if pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Polyethylene Glycol 3350, 17 g (cap filled to line).....Osmotic Laxative			
Use		Directions	
<ul style="list-style-type: none"> relieves occasional constipation (irregularity) generally produces a bowel movement in 1 to 3 days 		<ul style="list-style-type: none"> do not take more than directed unless advised by your doctor the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap) adults and children 17 years of age and older: <ul style="list-style-type: none"> fill to top of white section in cap which is marked to indicate the correct dose (17g) stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink ensure that the powder is fully dissolved before drinking do not drink if there are any clumps use once a day use no more than 7 days children 16 years of age or under: ask a doctor 	
Warnings		Other information	
Allergy alert: Do not use if you are allergic to polyethylene glycol Do not use if you have kidney disease, except under the advice and supervision of a doctor Ask a doctor before use if you have <ul style="list-style-type: none"> nausea, vomiting or abdominal pain a sudden change in bowel habits that lasts over 2 weeks irritable bowel syndrome 		<ul style="list-style-type: none"> store at 20° - 25°C (68° - 77°F) tamper-evident: do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken 	
Ask a doctor or pharmacist before use if you are taking a prescription drug When using this product you may have loose, watery, more frequent stools Stop use and ask a doctor if <ul style="list-style-type: none"> you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition. you get diarrhea you need to use a laxative for longer than 1 week 		Lot Expiry Area XXXXXX	
			
Bayer, the Bayer Cross, and MiraLAX are registered trademarks of Bayer. © 2019 Bayer. Dist by: Bayer HealthCare LLC, Whippany, NJ 07981			
			

Actual Size 100%

FDA00021

DRUG FACTS INFO		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt. **	9 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings [†]	8 pt.-bold italic [†]	8 pt.
Subheadings	≥6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	5 pt.
Leading	≥0.5 pt.	6.5 pt.
Hairlines	0.5 pt. rule	0.5 pt.
Box and Barlines [†]	>0.5 pt. rule	1.5 and 2.5 pt.
Characters Per Inch:	No more than 39	≤39

[†] Headings are to be at least 20pts larger than Body Text.
^{**} Drug Facts Title to be larger than all other type size used in DFP
[†] FDA Recommends 2.5pt. rule



Actual Size 100%

FDA00027

Drug Facts

Active ingredient (in each dose) Purpose
Polyethylene Glycol 3350,
17 g (cap filled to line).....Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed unless advised by your doctor**
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
 - fill to top of white section in cap which is marked to indicate the correct dose (17g)
 - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
 - ensure that the powder is fully dissolved before drinking
- do not drink if there are any clumps
- use once a day
- use no more than 7 days
- children 16 years of age or under: ask a doctor

Other information

- store at 20° - 25°C (68° - 77°F)
- tamper-evident: do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken

Inactive ingredients none

Questions or comments?
1-800-MiraLAX (1-800-647-2529)

• **Sugar Free**

000000

Actual Size 100%

FDA00026

DRUG FACTS INFO		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt. **	9 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings*	8 pt.-bold italic*	8 pt.
Subheadings	≥6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	5 pt.
Leading	≥0.5 pt.	6.5 pt.
Hairlines	0.5 pt. rule	0.5 pt.
Box and Barlines†	>0.5 pt. rule	1.5 and 2.5 pt.
Characters Per Inch:	No more than 39	≤39

* Headings are to be at least 2pts larger than Body Text.
 ** Drug Facts Title to be larger than all other type size used in DFP
 † FDA Recommends 2.5pt rule





Drug Facts

Active ingredient (in each dose) Purpose

Polyethylene Glycol 3350,
17 g (cap filled to line).....Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed unless advised by your doctor**
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
 - fill to top of white section in cap which is marked to indicate the correct dose (17g)
 - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
 - ensure that the powder is fully dissolved before drinking
 - do not drink if there are any clumps
 - use once a day
 - use no more than 7 days
- children 16 years of age or under: ask a doctor

Other information

- store at 20° - 25°C (68° - 77°F)
- tamper-evident: do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken

Inactive ingredients none

Questions or comments?

1-800-MiraLAX (1-800-647-2529)

• Sugar Free

000000



Drug Facts

Active ingredient (in each dose) Purpose

Polyethylene Glycol 3350,
17 g (cap filled to line).....Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed unless advised by your doctor**
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
 - fill to top of white section in cap which is marked to indicate the correct dose (17g)
 - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
 - ensure that the powder is fully dissolved before drinking
 - do not drink if there are any clumps
 - use once a day
 - use no more than 7 days
- children 16 years of age or under: ask a doctor

Other information

- store at 20° - 25°C (68° - 77°F)
- tamper-evident: do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken

Inactive ingredients none

Questions or comments?

1-800-MiraLAX (1-800-647-2529)

• Sugar Free

000000



2 FREE MIX-IN PAX® Single Dose Packets SEE BACK

#1 DOCTOR RECOMMENDED BRAND

MiraLAX®

Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

UNFLAVORED POWDER GRIT FREE

- Relieves Occasional Constipation/Irregularity
- Softens Stool

1 BOTTLE - NET WT 8.3 OZ (238 g)
2 PACKETS - NET WT 0.5 OZ (17 g) EACH
TOTAL NET WT 9.3 OZ (272 g)

16 ONCE-DAILY DOSES

BAYER

2 FREE MIX-IN PAX® Single Dose Packets Included



LOT/EXP AREA

Drug Facts		Drug Facts (continued)	
Active ingredient (in each dose)	Purpose	if pregnant or breast-feeding, ask a health professional before use.	
Polyethylene Glycol 3350, 17g (cap filled to line)	Osmotic Laxative	Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Use		Directions	
relieves occasional constipation (irregularity)		do not take more than directed unless advised by your doctor	
generally produces a bowel movement in 1 to 3 days		the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)	
Warnings		adults and children 17 years of age and older:	
Allergy alert: Do not use if you are allergic to polyethylene glycol		fill to top of white section in cap which is marked to indicate the correct dose (17g)	
Do not use if you have kidney disease, except under the advice and supervision of a doctor		str and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink	
Ask a doctor before use if you have		ensure that the powder is fully dissolved before drinking	
nausea, vomiting or abdominal pain		do not drink if there are any clumps	
a sudden change in bowel habits that lasts over 2 weeks		use once a day	
irritable bowel syndrome		use no more than 7 days	
Ask a doctor or pharmacist before use if you are taking a prescription drug		children 16 years of age or under: ask a doctor	
When using this product you may have loose, watery, more frequent stools		Other information	
Stop use and ask a doctor if		store at 20° - 25°C (68° - 77°F)	
you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.		tamper-evident: do not use if foil seal under cap, printed with "SEALED FOR YOUR PROTECTION" is missing, open or broken	
you get diarrhea		Inactive ingredients none	
you need to use a laxative for longer than 1 week		Questions or comments? 1-800-MiraLAX (1-800-647-2529)	



0 00000 00000 0

Bayer
00000
Whippany, NJ 07981
Dist by: Bayer HealthCare LLC,
© 2019 Bayer.
MiraLAX are registered trademarks of Bayer.
Bayer, the Bayer Cross, Mix-In-Pax and

Actual Size 100%

FDA00019

DRUG FACTS INFO		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt **	9 pt
Drug Facts Continued	8 pt	8 pt
Headings†	8 pt-bold italic*	8 pt
Subheadings	≥6 pt	6 pt
Body Text	≥6 pt	6 pt
Bullets	5 pt	3.5 pt
Leading	≥0.5 pt	6.5 pt
Headlines	0.5 pt rule	0.5 pt
Box and Barlines†	>0.5 pt rule	1 pt
Characters Per Inch:	No more than 39	≤39

* Headings are to be at least 2pts larger than Body Text.
** Drug Facts Title to be larger than all other type size used in DFP
† FDA Recommends 2.5pt rule





Drug Facts

Active ingredient (in each dose) **Purpose**
Polyethylene Glycol 3350,
17g (cap filled to line) Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed unless advised by your doctor**
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (note section in cap)
- adults and children 17 years of age and older:
 - fill to top of white section in cap which is marked to indicate the correct dose (17g)
 - stir and dissolve in any 4 to 8 ounces of beverage (at hot or room temperature) then drink
 - ensure that the powder is fully dissolved before drinking
 - do not drink if there are any clumps
 - use once a day
 - use no more than 7 days
- children 16 years of age or under: ask a doctor

Other information

- store at 20° - 25°C (68° - 77°F)
- tamper-evident: do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken

Inactive ingredients none

Questions or comments?

1-800-MiraLAX (1-800-647-2529)

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

KAREN M MAHONEY
06/03/2019 02:44:03 PM