

NOT FOR RESALE

Treats **FREQUENT Heartburn!**  24 HR

Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

14 TABLETS

One 14-day course of treatment
May take 1 to 4 days for full effect



Safety Feature — Do not use if printed seal under cap is broken or missing. This label does not contain full product information. See carton for complete information. Read warnings and directions on carton before use. Retain carton for reference.

Drug Facts

Active Ingredient Purpose
(in each tablet)
Omeprazole 20 mg.....Acid reducer

Use ■ Treats frequent heartburn (occurs *Zorinone* days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Other information

■ read the directions and warnings before use ■ keep the carton. ■ contains important information. ■ store at 20°-25°C (68°-77°F) and protect from moisture

Questions? 1-800-719-9260

Manufactured by:
Dowlex Pharmaceuticals Ltd
10 Hekdama Street, Yokneam 2069200, Israel

Made in Israel

092618

: 915D5 FA F10

Location for lot number and expiration date

NON-DFL FORMAT

TYPE SIZES:

HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

SUBHEADINGS: 7 pt Helvetica Neue LT Std 77 Bold Condensed

TEXT: 6 pt Helvetica Neue LT Std 57 Condensed

LEADING: 6 pt

BULLETS: 5 pt Square with 2 M's spacing between statements

TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed

HEAVY LINES: 1.0 pt

HAIRLINES LINES: 0.5 pt

Batch No.

EXP

915714 00 V3 092818

Location for lot number and expiration date

Drug Facts Active ingredient (in each tablet) Omeprazole 20 mg.....Acid reducer	Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect
Warnings Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness These may be signs of a serious condition. See your doctor.	Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain
Purpose Acid reducer	Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you get diarrhea ■ you develop a rash or joint pain
Drug Facts (continued) Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).	Directions ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours
14-Day Course of Treatment ■ swallow 1 tablet with a glass of water before eating in the morning ■ take every day for 14 days	Other information ■ read the directions and warnings before use ■ keep the carton, it contains important information. ■ store at 20-25°C (68-77°F) and protect from moisture
1-Day Course of Treatment ■ take every day for 14 days ■ do not take more than 1 tablet a day	Inactive ingredients carboxylic acid, ferric oxide red, ferric oxide yellow, hypromellose, hydroxypropylcellulose, lactose monohydrate, monoethanolamine, polyethylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate
Repeated 14-Day Courses (if needed) ■ do not use for more than 14 days unless directed by your doctor ■ do not chew or crush tablets. ■ you may repeat a 14-day course every 4 months	Questions or comments? 1-800-719-9260

NOT FOR RESALE

Safety Feature –
Do not use if printed tablet blister unit is open or torn.

Compare to Prilosec OTC®*

Treats **FREQUENT** Heartburn! **24HR**

Omeprazole

Delayed Release Tablets 20 mg
Acid Reducer

14 TABLETS

One 14-day course of treatment
May take 1 to 4 days for full effect



Actual Size

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued)
8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Made in Israel
 Manufactured by:
Dexcel Pharma Technologies Ltd
 10 Handkina Street, Yokneam 2109200, Israel

*This product is not
 manufactured or distributed
 by Pocher & Gamble,
 distributor of Prilosec OTC®.

<p>Drug Facts (continued)</p> <p>Directions If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p> <p>For adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours</p> <p>14-Day Course of Treatment ■ swallow 1 tablet with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ swallow whole. Do not chew or crush tablets.</p> <p>Repeated 14-Day Courses (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor</p> <p>■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.</p> <p>Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F) and protect from moisture</p> <p>Inactive ingredients camabta wax, ferric oxide red, ferric oxide yellow, hypromellose, polyethylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate</p> <p>Questions or comments? 1-800-719-9260</p>	<p>Drug Facts</p> <p>Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Warnings ■ Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain ■ These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain</p> <p>Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain</p> <p>Acid reducer</p>
---	---

Location for lot number and expiration date

91574 FA C18 100218




Omeprazole

Delayed Release Tablets 20 mg

Safety Feature - Do not use if printed tablet blister unit is open or torn.

Compare to Prilosec OTC®*



Treats **FREQUENT** Heartburn!




Omeprazole

Delayed Release Tablets 20 mg

Acid Reducer

14 TABLETS

One 14-day course of treatment
 May take 1 to 4 days for full effect





STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

0000000000

Safety Feature - Do not use if printed seal under cap is broken or missing.

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- swallow whole. Do not chew or crush tablets.

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor

children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Drug Facts (continued)

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

Inactive ingredients camauba wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-719-9260

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

1 BOTTLE INSIDE

Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

NDC 0000-000-00 Compare to Prilosec OTC®*

DIE CUT WINDOW

Treats **FREQUENT Heartburn!** **24 HR**

Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

14 TABLETS

One 14-day course of treatment
May take 1 to 4 days for full effect



Drug Facts

Active ingredient (in each tablet)	Purpose
Omeprazole 20 mg.....	Acid reducer

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

0000000000



0 00000 00000 0

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

915D5 FA C12 092818

Location for lot number and expiration date

STANDARD FORMAT

TYPE SIZES:

TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

(continued) 8 pt Helvetica Neue LT Std 57 Condensed

HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed

TEXT: 6 pt Helvetica Neue LT Std 57 Condensed

LEADING: 6.5 pt

BULLETS: 5 pt Square with 2 M'S spacing between statements

TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed

HEAVY LINES: 1.5 pt

HAIRLINES: 0.5 pt

Hairlines extend to within two spaces of the "Drug Facts" box

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Questions or comments? 1-800-719-9260

Inactive ingredients canabadi wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, polyethylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate

Other information
 ■ read the directions and warnings before use
 ■ keep the carton. It contains important information.
 ■ store at 20-25°C (68-77°F) and protect from moisture

Ask a doctor before use if you have:
 ■ head heartburn over 3 months. This may be a sign of a more serious condition.
 ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss
 ■ nausea or vomiting
 ■ stomach pain

Ask a doctor before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.
Stop use and ask a doctor if:
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you get diarrhea
 ■ you develop a rash or joint pain

Warnings
 Do not use if you have:
 ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
 ■ heartburn with lightheadedness, sweating or dizziness
 ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
Do not use if you have:
 ■ frequent chest pain
 ■ These may be signs of a serious condition. See your doctor.

Use
 ■ Treats frequent heartburn (occurs 2 or more days a week)
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Active ingredient (in each tablet)
 Omeprazole 20 mg.....Acid reducer

Purpose
 Acid reducer

Directions
 ■ for adults 18 years of age and older
 ■ this product is to be used once a day (every 24 hours), every day for 14 days
 ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
 ■ swallow 1 tablet with a glass of water before eating in the morning
 ■ take every day for 14 days
 ■ do not take more than 1 tablet a day

Repeated 14-Day Courses (if needed)
 ■ do not use for more than 14 days unless directed by your doctor
 ■ swallow whole. Do not chew or crush tablets.

Do not take for more than 14 days or more often than every 4 months unless directed by a doctor
 ■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Drug Facts (continued)
 If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Made in Israel
 Manufactured by:
Dexcel Pharma Technologies Ltd
 10 Hakdima Street, Yokneam 2069200, Israel

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Location for lot number and expiration date

91574 FA C18-F 100218



Omeprazole
 Delayed Release Tablets 20 mg

Safety Feature - Do not use if printed tablet blister unit is open or torn.

Compare to Prilosec OTC®
FDA Approved

Treats **FREQUENT** Heartburn!



Omeprazole
 Delayed Release Tablets 20 mg
 Acid Reducer

14 TABLETS

One 14-day course of treatment
 May take 1 to 4 days for full effect



STANDARD FORMAT

TYPE SIZES:
 TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued) 8 pt Helvetica Neue LT Std 57 Condensed
 HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
 TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
 LEADING: 6.5 pt
 BULLETS: 5 pt Square with 2 M'S spacing between statements
 TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
 HEAVY LINES: 1.5 pt
 HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box



Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- swallow whole. Do not chew or crush tablets.

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor

children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Drug Facts (continued)

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

Inactive ingredients carnauba wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-719-9260

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

1 BOTTLE INSIDE

Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

NDC 0000-000-00 Compare to Prilosec OTC®*
FDA Approved

DIE CUT WINDOW

Treats **FREQUENT Heartburn!** **24 HR**

Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

14 TABLETS

One 14-day course of treatment
May take 1 to 4 days for full effect



*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

915D5 FA C12-F 092818

Drug Facts

Active ingredient (in each tablet)	Purpose
Omeprazole 20 mg.....	Acid reducer

Use

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain


These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued) **8 pt Helvetica Neue LT Std 57 Condensed**
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box



- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Tips for Managing Heartburn

- you develop a rash or joint pain
- you get diarrhea
- you need to take more than 1 course of treatment every 4 months
- your heartburn continues or worsens

Stop use and ask a doctor if:

- you need to take this product for more than 14 days
- you develop a rash or joint pain
- you get diarrhea
- you need to take more than 1 course of treatment every 4 months
- your heartburn continues or worsens

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

- stomach pain
- nausea or vomiting
- unexplained weight loss
- frequent wheezing, particularly with heartburn
- had heartburn over 3 months. This may be a sign of a more serious condition.

Ask a doctor before use if you have:

- frequent chest pain
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- heartburn with lightheadedness, sweating or dizziness
- black stools
- trouble or pain swallowing food, vomiting with blood, or bloody or black stools

Do not use if you have:

- Allergy alert: Do not use if you are allergic to omeprazole
- frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Use
Omeprazole 20 mg (in each tablet)

Purpose
Acid reducer

Drug Facts (continued)

Questions or comments? 1-800-719-9260

Inactive ingredients carnauba wax, ferric oxide red, ferric oxide yellow, hypromellose, hydroxypropyl methylcellulose, lactose monohydrate, monoethanolamine, polyethylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate

Other information

- store at 20-25°C (68-77°F) and protect from moisture
- keep the carton. It contains important information.
- read the directions and warnings before use

■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

■ 4 months unless directed by a doctor

■ do not take for more than 14 days or more often than every 4 months

■ you may repeat a 14-day course every 4 months

■ Repeated 14-Day Courses (if needed)

■ do not take more than 14 days unless directed by your doctor

■ do not take more than 1 tablet a day

■ take every day for 14 days

■ swallow 1 tablet with a glass of water before eating in the morning

■ 14-Day Course of Treatment

■ symptoms within 24 hours

■ it may take 1 to 4 days for full effect; some people get complete relief or

■ this product is to be used once a day (every 24 hours), every day for 14 days

■ for adults 18 years of age and older

■ Directions

■ If pregnant or breast-feeding, ask a health professional before use.

■ Keep out of reach of children. In case of overdose, get medical help or

■ contact a Poison Control Center right away (1-800-222-1222).

■ Drug Facts (continued)

Omeprazole
Delayed Release Tablets 20 mg

Made in Israel
Manufactured by:
Cescael Pharma Technologies Ltd
10 Hakdima Street, Yokneam 2089200, Israel
*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

91530 FA C18 100318

Location for lot number and expiration date

Safety Feature - Do not use if printed tablet blister unit is open or torn.



Omeprazole
Delayed Release Tablets 20 mg

Compare to Prilosec OTC®*

Treats **FREQUENT** Heartburn!



Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

28 TABLETS

Two 14-day courses of treatment
May take 1 to 4 days for full effect



STANDARD FORMAT
TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

0000000000



Drug Facts

Active ingredient (in each tablet) Purpose
Omeprazole 20 mg.....Acid reducer

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens

Drug Facts (continued)

- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea ■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- swallow whole. Do not chew or crush tablets.

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**

- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Drug Facts (continued)

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

Inactive ingredients camauba wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-719-9260

2 BOTTLES INSIDE

Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

Safety Feature - Do not use if printed seal under cap is broken or missing.

NDC 0000-000-00

Compare to Prilosec OTC®*

DIE CUT WINDOW

DIE CUT WINDOW

Treats **FREQUENT** Heartburn!



Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

28 TABLETS

Two 14-day courses of treatment
May take 1 to 4 days for full effect



Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

0000000000



*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

915D6 FA C12 092818

Location for lot number and expiration date

STANDARD FORMAT

TYPE SIZES:

- TITLE:** 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
- TITLE (continued):** 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued) 8 pt Helvetica Neue LT Std 57 Condensed
- HEADINGS:** 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
- SUBHEADINGS:** 6 pt Helvetica Neue LT Std 77 Bold Condensed
- TEXT:** 6 pt Helvetica Neue LT Std 57 Condensed
- LEADING:** 6.5 pt
- BULLETS:** 5 pt Square with 2 M'S spacing between statements
- TELEPHONE #:** 6 pt Helvetica Neue LT Std 77 Bold Condensed
- HEAVY LINES:** 1.5 pt
- HAIRLINES:** 0.5 pt
- Hairlines extend to within two spaces of the "Drug Facts" box**



- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

Omeprazole

Delayed Release Tablets 20 mg

Made in Israel
 Manufactured by:
Celexcel Pharma Technologies Ltd
 10 Hakdima Street, Yokneam 2069200, Israel
 *This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

<p>Drug Facts (continued)</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p> <p>Directions</p> <ul style="list-style-type: none"> ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours <p>14-Day Course of Treatment</p> <ul style="list-style-type: none"> ■ swallow 1 tablet with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ swallow whole. Do not chew or crush tablets. <p>Repeated 14-Day Courses (if needed)</p> <ul style="list-style-type: none"> ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor <ul style="list-style-type: none"> ■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition. <p>Other information</p> <ul style="list-style-type: none"> ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F) and protect from moisture. <p>Inactive ingredients carmelum xaa, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monothalamine, polyethylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate</p> <p>Questions or comments? 1-800-719-9260</p>	<p>Drug Facts</p> <p>Omeprazole 20 mg.....Acid reducer</p> <p>Purpose</p> <p>Acid reducer</p> <p>Use</p> <ul style="list-style-type: none"> ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect <p>Warnings</p> <p>Allergy alert: Do not use if you are allergic to omeprazole</p> <p>Do not use if you have:</p> <ul style="list-style-type: none"> ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain <p>These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have:</p> <ul style="list-style-type: none"> ■ had heartburn over 3 months; This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain <p>Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if:</p> <ul style="list-style-type: none"> ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain
---	--

91530 FA C18-F 100318

Location for lot number and expiration date



Omeprazole

Delayed Release Tablets 20 mg

Safety Feature - Do not use if printed tablet blister unit is open or torn.

NDC 0000-000-00

Compare to Prilosec OTC®*
FDA Approved

Treats **FREQUENT** Heartburn!



Omeprazole

Delayed Release Tablets 20 mg

Acid Reducer

28 TABLETS

Two 14-day courses of treatment
 May take 1 to 4 days for full effect



STANDARD FORMAT

TYPE SIZES:

- TITLE:** 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
- TITLE (continued):** 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued) 8 pt Helvetica Neue LT Std 57 Condensed
- HEADINGS:** 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
- SUBHEADINGS:** 6 pt Helvetica Neue LT Std 77 Bold Condensed
- TEXT:** 6 pt Helvetica Neue LT Std 57 Condensed
- LEADING:** 6.5 pt
- BULLETS:** 5 pt Square
- TELEPHONE #:** 6 pt Helvetica Neue LT Std 77 Bold Condensed
- HEAVY LINES:** 1.5 pt
- HAIRLINES:** 0.5 pt
- Hairlines extend to within two spaces of the "Drug Facts" box**

0000000000



0000000000

Drug Facts

Active ingredient (in each tablet) Purpose
Omeprazole 20 mg.....Acid reducer

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness

frequent chest pain

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens

Drug Facts (continued)

- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea ■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- swallow whole. Do not chew or crush tablets.

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Drug Facts (continued)

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture


Inactive ingredients camauba wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-719-9260

2 BOTTLES INSIDE

Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

Safety Feature - Do not use if printed seal under cap is broken or missing.



NDC 0000-000-00

Compare to Prilosec OTC®*
FDA Approved



Treats **FREQUENT** Heartburn! **24 HR**

Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer


28 TABLETS

Two 14-day courses of treatment
May take 1 to 4 days for full effect




- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

0000000000



0000000000




*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Location for lot number and expiration date

915D6 FA C12-F 092818

STANDARD FORMAT

- TYPE SIZES:**
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box



- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Tips for Managing Heartburn

Drug Facts

Active ingredient (in each tablet) Omeprazole 20 mg.....Acid reducer

Purpose Acid reducer

Use Omeprazole 20 mg.....Acid reducer

Warnings Do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief or symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- swallow whole. Do not chew or crush tablets.

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor

Other information

- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.
- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

Inactive ingredients canaxaba wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-719-9260

Omeprazole

Delayed Release Tablets 20 mg

Made in Israel
 Manufactured by:
Deacel Pharma Technologies Ltd
 10 Hakdima Street, Yokneam 2069200, Israel
 *The products are not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

91530 FA C18-B 100318

Location for lot number and expiration date

Safety Feature - Do not use if printed tablet blister unit is open or torn.



Omeprazole

Delayed Release Tablets 20 mg

NDC 0000-000-00

Compare to Prilosec OTC®*

14 FREE

Treats **FREQUENT** Heartburn!



Omeprazole

Delayed Release Tablets 20 mg

Acid Reducer

28 TABLETS

Two 14-day courses of treatment
 May take 1 to 4 days for full effect




STANDARD FORMAT

TYPE SIZES:

- TITLE:** 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
- TITLE (continued):** 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued) 8 pt Helvetica Neue LT Std 57 Condensed
- HEADINGS:** 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
- SUBHEADINGS:** 6 pt Helvetica Neue LT Std 77 Bold Condensed
- TEXT:** 6 pt Helvetica Neue LT Std 57 Condensed
- LEADING:** 6.5 pt
- BULLETS:** 5 pt Square
- TELEPHONE #:** 6 pt Helvetica Neue LT Std 77 Bold Condensed
- HEAVY LINES:** 1.5 pt
- HAIRLINES:** 0.5 pt
- Hairlines extend to within two spaces of the "Drug Facts" box

0000000000



0000000000

Drug Facts

Active ingredient (in each tablet) Purpose
Omeprazole 20 mg.....Acid reducer

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
 - heartburn with **lightheadedness, sweating or dizziness**
 - chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
 - frequent **chest pain**
- These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens

Drug Facts (continued)

- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea ■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- swallow whole. Do not chew or crush tablets.

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor

■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition. ▶

Drug Facts (continued)

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

Inactive ingredients camauba wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-719-9260

2 BOTTLES INSIDE

Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

Safety Feature - Do not use if printed seal under cap is broken or missing.

NDC 0000-000-00

Compare to Prilosec OTC®*

14 FREE

DIE CUT WINDOW

DIE CUT WINDOW

Treats **FREQUENT** Heartburn!

24 HR

Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer


28 TABLETS

Two 14-day courses of treatment
May take 1 to 4 days for full effect




Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking



0000000000

0000000000



*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

915D6 FA C12-B 092818

Location for lot number and expiration date

STANDARD FORMAT

TYPE SIZES:

- TITLE:** 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
- TITLE (continued):** 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
- (continued)** 8 pt Helvetica Neue LT Std 57 Condensed
- HEADINGS:** 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
- SUBHEADINGS:** 6 pt Helvetica Neue LT Std 77 Bold Condensed
- TEXT:** 6 pt Helvetica Neue LT Std 57 Condensed
- LEADING:** 6.5 pt
- BULLETS:** 5 pt Square with 2 M'S spacing between statements
- TELEPHONE #:** 6 pt Helvetica Neue LT Std 77 Bold Condensed
- HEAVY LINES:** 1.5 pt
- HAIRLINES:** 0.5 pt
- Hairlines extend to within two spaces of the "Drug Facts" box**



- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

Drug Facts (continued)

Active ingredient (in each tablet).....Omeprazole 20 mg.....Acid reducer

Purpose
 ■ treats frequent heartburn (occurs *2 or more* days a week)
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Do not use if you have:
 ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
 ■ heartburn with **lightheadedness, sweating or dizziness**
 ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
Ask a doctor before use if you have:
 ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss
 ■ nausea or vomiting
 ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea
 ■ you develop a rash or joint pain

Questions or comments? 1-800-719-9260

Inactive ingredients carnauba wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate

Other information
 ■ keep the carton. It contains important information.
 ■ store at 20-25°C (68-77°F) and protect from moisture

Directions
 ■ for adults 18 years of age and older
 ■ this product is to be used once a day (every 24 hours), every day for 14 days
 ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
 ■ swallow 1 tablet with a glass of water before eating in the morning
 ■ take every day for 14 days
 ■ do not take more than 1 tablet a day

Repeated 14-Day Courses (if needed)
 ■ do not use for more than 14 days unless directed by your doctor
 ■ do not take more than 1 tablet a day
 ■ swallow whole. Do not chew or crush tablets.

4 months unless directed by a doctor
 ■ you may repeat a 14-day course every 4 months
 ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Omeprazole

Delayed Release Tablets 20 mg

Made in Israel
 Manufactured by:
 Dexcel Pharma Technologies Ltd
 10 Hakdima Street, Yokneam 2099200, Israel
 *This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

91530 FA C18-E 100318

Location for lot number and expiration date



Omeprazole

Delayed Release Tablets 20 mg

Safety Feature - Do not use if printed tablet blister unit is open or torn.

NDC 0000-000-000

Compare to Prilosec OTC®*
Bonus! 14 Free

Treats **FREQUENT** Heartburn!



Omeprazole

Delayed Release Tablets 20 mg

Acid Reducer

~~28~~ **14** TABLETS

Two ~~One~~ 14-day courses of treatment
 May take 1 to 4 days for full effect




STANDARD FORMAT

TYPE SIZES:

- TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
- TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued)
- 8 pt Helvetica Neue LT Std 57 Condensed
- HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
- SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
- TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
- LEADING: 6.5 pt
- BULLETS: 5 pt Square
- TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
- HEAVY LINES: 1.5 pt
- HAIRLINES: 0.5 pt
- Hairlines extend to within two spaces of the "Drug Facts" box

00000000



00000000

Drug Facts

Active ingredient (in each tablet) Purpose
Omeprazole 20 mg..... Acid reducer

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens

Drug Facts (continued)

- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea ■ you develop a rash or joint pain

If **pregnant or breast-feeding**, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- swallow whole. Do not chew or crush tablets.

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**

■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition. ▶

Drug Facts (continued)

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

Inactive ingredients: carnauba wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-719-9260

2 BOTTLES INSIDE

Made in Israel
Manufactured by:
Dexel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel



00000000

00000000

Safety Feature - Do not use if printed seal under cap is broken or missing.

NDC 0000-000-00

Compare to Prilosec OTC®*
Bonus! 14 Free



Treats **FREQUENT** Heartburn!



Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

28 14 TABLETS

Two ~~One~~ 14-day courses of treatment
May take 1 to 4 days for full effect




*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

915D6 FA C12-E 100318

Location for lot number and expiration date

STANDARD FORMAT

- TYPE SIZES:**
- TITLE:** 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
- TITLE (continued):** 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
- (continued)** 8 pt Helvetica Neue LT Std 57 Condensed
- HEADINGS:** 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
- SUBHEADINGS:** 6 pt Helvetica Neue LT Std 77 Bold Condensed
- TEXT:** 6 pt Helvetica Neue LT Std 57 Condensed
- LEADING:** 6.5 pt
- BULLETS:** 5 pt Square with 2 M'S spacing between statements
- TELEPHONE #:** 6 pt Helvetica Neue LT Std 77 Bold Condensed
- HEAVY LINES:** 1.5 pt
- HAIRLINES:** 0.5 pt
- Hairlines extend to within two spaces of the "Drug Facts" box**



- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

Omeprazole
Delayed Release Tablets 20 mg

Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakikma Street, Yokneam 2069200, Israel

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Drug Facts (continued)

Active ingredient (in each tablet).....Omeprazole 20 mg.....Acid reducer

Purpose.....Acid reducer

Use
■ treats frequent heartburn (occurs *2 or more* days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:
■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
■ heartburn with **lightheadedness, sweating or dizziness**
■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness

These may be signs of a serious condition. See your doctor.
Ask a doctor before use if you have:
■ frequent **chest pain**
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss
■ nausea or vomiting
■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:
■ your heartburn continues or worsens
■ you need to take this product for more than 14 days
■ you need to take more than 1 course of treatment every 4 months
■ you get diarrhea
■ you develop a rash or joint pain

Questions or comments? 1-800-719-9260

Inactive ingredients carnuba wax, ferric oxide red, ferric oxide yellow, hypromellose, hydroxypropylcellulose, lactose monohydrate, monothramine, polyene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate

Other information
■ keep the carton. It contains important information.
■ store at 20-25°C (68-77°F) and protect from moisture
■ read the directions and warnings before use

Directions
■ for adults 18 years of age and older
■ this product is to be used once a day (every 24 hours), every day for 14 days
■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
■ swallow 1 tablet with a glass of water before eating in the morning
■ take every day for 14 days
■ do not take more than 1 tablet a day

Repeated 14-Day Courses (if needed)
■ you may repeat a 14-day course every 4 months
■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

91530 FA C18-D 100318

Location for lot number and expiration date



Omeprazole
Delayed Release Tablets 20 mg

Safety Feature - Do not use if printed tablet blister unit is open or torn.



NDC 0000-000-00

Compare to Prilosec OTC®*
Bonus Pack! 14 Free

Treats **FREQUENT** Heartburn!



Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

28 ~~14~~ TABLETS



Two ~~One~~ 14-day courses of treatment
May take 1 to 4 days for full effect



STANDARD FORMAT


- TYPE SIZES:**
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

0000000000



0000000000

Safety Feature - Do not use if printed seal under cap is broken or missing.



Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Drug Facts

Active ingredient (in each tablet) Purpose
Omeprazole 20 mg.....Acid reducer

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating, pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens

Drug Facts (continued)

- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea ■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- swallow whole. Do not chew or crush tablets.

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**

- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Drug Facts (continued)

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

Inactive ingredients: carnauba wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-719-9260

2 BOTTLES INSIDE
Made in Israel
Manufactured by:
Dexel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

NDC 0000-000-00

Compare to Prilosec OTC®*

Bonus Pack! 14 Free

DIE CUT WINDOW

DIE CUT WINDOW

Treats **FREQUENT** Heartburn!

24 HR

Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

28 TABLETS



Two ~~One~~ 14-day courses of treatment
May take 1 to 4 days for full effect



*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Location for lot number and expiration date


915D6 FA C12-D 100318

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

Location for lot number and expiration date

- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking




<p>Drug Facts</p> <p>Active ingredient (in each tablet).....Omeprazole 20 mg</p> <p>Purpose.....Acid reducer</p>	<p>Use</p> <ul style="list-style-type: none"> ■ treats frequent heartburn (occurs <i>2 or more</i> days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect 	<p>Warnings</p> <p>Allergy alert: Do not use if you are allergic to omeprazole</p> <p>Do not use if you have:</p> <ul style="list-style-type: none"> ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness <p>Frequent chest pain</p> <p>These may be signs of a serious condition. See your doctor.</p>	<p>Ask a doctor before use if you have:</p> <ul style="list-style-type: none"> ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain <p>Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p>	<p>Stop use and ask a doctor if:</p> <ul style="list-style-type: none"> ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain 	<p>Questions or comments? 1-800-719-9260</p>
<p>Drug Facts (continued)</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p>	<p>Directions</p> <ul style="list-style-type: none"> ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours 	<p>14-Day Course of Treatment</p> <ul style="list-style-type: none"> ■ swallow 1 tablet with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ swallow whole. Do not chew or crush tablets. 	<p>Repeated 14-Day Courses (if needed)</p> <ul style="list-style-type: none"> ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor 	<p>Other information</p> <ul style="list-style-type: none"> ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F) and protect from moisture 	<p>Inactive ingredients camara wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate</p>

91555 FA C18 100318

Omeprazole
Delayed Release Tablets 20 mg

Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakdima Street, Yokneam 2069200, Israel



*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.



Omeprazole

Delayed Release Tablets 20 mg

Safety Feature - Do not use if printed tablet blister unit is open or torn.



NDC 0000-000-00

Compare to Prilosec OTC®*

Treats **FREQUENT** Heartburn! **24 HR**


Omeprazole

Delayed Release Tablets 20 mg

Acid Reducer

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect



STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box

Drug Facts

Active ingredient (in each tablet)

Omeprazole 20 mg.....Acid reducer

Purpose

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath, sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days

Drug Facts (continued)

- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- swallow whole. Do not chew or crush tablets.

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**

■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

Drug Facts (continued)

Inactive ingredients

carnauba wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate

Questions or comments?

1-800-719-9260

3 BOTTLES INSIDE

Made in Israel
Manufactured by:
Dexel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

Safety Feature - Do not use if printed seal under cap is broken or missing.

NDC 0000-000-00

Compare to Prilosec OTC®*



Treats **FREQUENT** Heartburn! **24 HR**

Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect



Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking



*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Location for lot number and expiration date

915D7 FA C12 092818

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

Location for lot number and expiration date

- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Ask a doctor before use if you have:

- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain

Warnings

■ Allergy alert: Do not use if you are allergic to omeprazole

Use

- treats frequent heartburn (occurs *2 or more* days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Active ingredient (in each tablet)

Omeprazole 20 mg, Acid reducer

Purpose

Acid reducer

Drug Facts

Other information

- read the directions and warnings before use
- keep the carton, it contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

Inactive ingredients

yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, glycolate, sodium stearate, sodium lauryl sulfate, sodium starch triethyl citrate

Questions or comments? 1-800-719-9260

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- swallow whole. Do not chew or crush tablets.

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor

4 months unless directed by a doctor

- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Drug Facts (continued)

Use

Active ingredient (in each tablet)

Purpose

Warnings

Do not use if you have:

Ask a doctor before use if you have:

Stop use and ask a doctor if:

Other information

Questions or comments? 1-800-719-9260

Inactive ingredients

Directions

14-Day Course of Treatment

Repeated 14-Day Courses (if needed)

4 months unless directed by a doctor

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Drug Facts (continued)

Omeprazole

Delayed Release Tablets 20 mg

Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel
*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Safety Feature - Do not use if printed tablet blister unit is open or torn.



Omeprazole

Delayed Release Tablets 20 mg

Compare to Prilosec OTC®*
FDA Approved

Treats **FREQUENT** Heartburn!



Omeprazole

Delayed Release Tablets 20 mg

Acid Reducer

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect



STANDARD FORMAT

TYPE SIZES:

TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

(continued) 8 pt Helvetica Neue LT Std 57 Condensed

HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed

TEXT: 6 pt Helvetica Neue LT Std 57 Condensed

LEADING: 6.5 pt

BULLETS: 5 pt Square

TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed

HEAVY LINES: 1.5 pt

HAIRLINES: 0.5 pt

Hairlines extend to within two spaces of the "Drug Facts" box

Drug Facts

Active ingredient (in each tablet)

Omeprazole 20 mg.....Acid reducer

Purpose

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days

Drug Facts (continued)

- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- swallow whole. Do not chew or crush tablets.

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**

■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

Drug Facts (continued)

Inactive ingredients

carnauba wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate

Questions or comments?

1-800-719-9260

3 BOTTLES INSIDE

Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

Safety Feature - Do not use if printed seal under cap is broken or missing.

NDC 0000-000-00

Compare to Prilosec OTC®*

FDA Approved

DIE CUT WINDOW

DIE CUT WINDOW

DIE CUT WINDOW

Treats **FREQUENT** Heartburn!



Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect



Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking



*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Location for lot number and expiration date

915D7 FA C12-F 092818

STANDARD FORMAT

TYPE SIZES:

TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued) 8 pt Helvetica Neue LT Std 57 Condensed

HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed

TEXT: 6 pt Helvetica Neue LT Std 57 Condensed

LEADING: 6.5 pt

BULLETS: 5 pt Square with 2 M'S spacing between statements

TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed

HEAVY LINES: 1.5 pt

HAIRLINES: 0.5 pt

Hairlines extend to within two spaces of the "Drug Facts" box

Location for lot number and expiration date

91555 FA C18-B 100318

- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

Drug Facts	Active ingredient (in each tablet) Omeprazole 20 mgAcid reducer
Use	■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect
Warnings	Allergy alert: Do not use if you are allergic to omeprazole Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain These may be signs of a serious condition. See your doctor. Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.
Stop use and ask a doctor if:	■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain
Drug Facts (continued)	Purpose If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).
Directions	■ for adults 18 years of age and older ■ this product is to be used once a day every 24 hours, every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief or symptoms within 24 hours
14-Day Course of Treatment	■ swallow 1 tablet with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor
Repeated 14-Day Courses (if needed)	■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor
Other information	■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition. ■ read the directions and warnings before use ■ keep the carton, it contains important information ■ store at 20-25°C (68-77°F) and protect from moisture
Inactive ingredients	canabada wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate
Questions or comments? 1-800-719-9260	

Omeprazole

Delayed Release Tablets 20 mg

Made in Israel
Manufactured by:
Cexxel Pharma Technologies Ltd
10 Hekidma Street, Yokneam 2069200, Israel
*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.



Omeprazole

Delayed Release Tablets 20 mg

Safety Feature - Do not use if printed tablet blister unit is open or torn.

NDC 0000-000-00

Compare to Prilosec OTC®*
14 FREE

Treats **FREQUENT** Heartburn!



Omeprazole

Delayed Release Tablets 20 mg

Acid Reducer

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect



STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box

Safety Feature - Do not use if printed seal under cap is broken or missing.

NDC 0000-000-00

Compare to Prilosec OTC®*
14 FREE

DIE CUT WINDOW

DIE CUT WINDOW

DIE CUT WINDOW

Treats **FREQUENT** Heartburn! **24 HR**

Omeprazole

Delayed Release Tablets 20 mg Acid Reducer

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

915D7 FA C12-B 092818

- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

Drug Facts

Active ingredient (in each tablet) Omeprazole 20 mg.....
Purpose Acid reducer

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days

Drug Facts (continued)

- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- swallow whole. Do not chew or crush tablets.

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**

■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

Drug Facts (continued)

Inactive ingredients
carnauba wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate

Questions or comments?
1-800-719-9260

3 BOTTLES INSIDE

Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel



Location for lot number and expiration date

STANDARD FORMAT

TYPE SIZES:

TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued)

8 pt Helvetica Neue LT Std 57 Condensed

HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed

TEXT: 6 pt Helvetica Neue LT Std 57 Condensed

LEADING: 6.5 pt

BULLETS: 5 pt Square with 2 M'S spacing between statements

TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed

HEAVY LINES: 1.5 pt

HAIRLINES: 0.5 pt

Hairlines extend to within two spaces of the "Drug Facts" box

Location for lot number and expiration date

- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

Omeprazole

Delayed Release Tablets 20 mg

Made in Israel
 Manufactured by:
Celexel Pharma Technologies Ltd
 10 Hakdima Street, Yokneam 2069200, Israel
 This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Omeprazole 20 mg.....Acid reducer</p> <p>Purpose Acid reducer</p>	<p>Use ■ treats frequent heartburn (occurs <i>2 or more</i> days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Warnings Allergy alert: Do not use if you are allergic to omeprazole</p> <p>Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness ■ frequent chest pain These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain</p> <p>Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain</p>
<p>Drug Facts (continued)</p> <p>Directions ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours</p> <p>14-Day Course of Treatment ■ swallow 1 tablet with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 tablet a day</p> <p>Repeated 14-Day Courses (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor</p> <p>Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F) and protect from moisture</p> <p>Inactive ingredients canabada wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearoyl fumarate, talc, titanium dioxide, triethyl citrate</p> <p>Questions or comments? 1-800-719-9260</p>	<p>Drug Facts</p> <p>Active ingredient (in each tablet) Omeprazole 20 mg.....Acid reducer</p> <p>Purpose Acid reducer</p>

91555 FA C18-C 100318



Omeprazole

Delayed Release Tablets 20 mg

Safety Feature - Do not use if printed tablet blister unit is open or torn.

NDC 0000-000-00

Compare to Prilosec OTC®*
28 + 14 FREE

Treats **FREQUENT** Heartburn!



Omeprazole

Delayed Release Tablets 20 mg
 Acid Reducer

42 TABLETS

Three 14-day courses of treatment
 May take 1 to 4 days for full effect



STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box

Drug Facts

Active ingredient (in each tablet)

Omeprazole 20 mg.....Acid reducer

Purpose

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
 - heartburn with **lightheadedness, sweating or dizziness**
 - chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
 - frequent **chest pain**
- These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days

Drug Facts (continued)

- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- swallow whole. Do not chew or crush tablets.

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

Drug Facts (continued)

Inactive ingredients

carnauba wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate

Questions or comments?

1-800-719-9260

3 BOTTLES INSIDE

Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

Safety Feature - Do not use if printed seal under cap is broken or missing.

NDC 0000-000-00

Compare to Prilosec OTC®*

28 + 14 FREE

DIE CUT WINDOW

DIE CUT WINDOW

DIE CUT WINDOW

Treats **FREQUENT** Heartburn!



Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect



Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking



*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Location for lot number and expiration date

915D7 FA C12-C 092818

STANDARD FORMAT

TYPE SIZES:

TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

(continued) 8 pt Helvetica Neue LT Std 77 Condensed

HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed

TEXT: 6 pt Helvetica Neue LT Std 57 Condensed

LEADING: 6.5 pt

BULLETS: 5 pt Square with 2 M'S spacing between statements

TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed

HEAVY LINES: 1.5 pt

HAIRLINES: 0.5 pt

Hairlines extend to within two spaces of the "Drug Facts" box

Location for lot number and expiration date

91555 FA C18-E 100218

- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

Drug Facts	Active ingredient (in each tablet) Omeprazole 20 mg Acid reducer
Use	■ treats frequent heartburn (occurs <i>2 or more</i> days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect
Warnings	Allergy alert: Do not use if you are allergic to omeprazole Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain These may be signs of a serious condition. See your doctor. Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain ■ Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.
Stop use and ask a doctor if:	■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you get diarrhea ■ you develop a rash or joint pain
Drug Facts (continued)	Directions ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours
	14-Day Course of Treatment ■ swallow 1 tablet with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 tablet a day
	Repeated 14-Day Courses (if needed) ■ do not use for more than 14 days unless directed by your doctor ■ swallow whole. Do not chew or crush tablets.
	Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F) and protect from moisture
	Inactive ingredients canabada wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate
	Questions or comments? 1-800-719-9260

Omeprazole

Delayed Release Tablets 20 mg

Made in Israel
Manufactured by:
Dexcel Pharma Technologies, Ltd
10 Hakidma Street, Yokneam 2089200, Israel
*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.



Omeprazole

Delayed Release Tablets 20 mg

Safety Feature - Do not use if printed tablet blister unit is open or torn.

NDC 0000-000-00

Compare to Prilosec OTC®*
Bonus! 14 Free

Treats **FREQUENT** Heartburn!



Omeprazole

Delayed Release Tablets 20 mg

Acid Reducer

42 ~~28~~ TABLETS

Three ~~Two~~ 14-day courses of treatment
May take 1 to 4 days for full effect



STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued)
 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box

Drug Facts

Active ingredient (in each tablet)

Omeprazole 20 mg.....Acid reducer

Purpose

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
 - heartburn with **lightheadedness, sweating or dizziness**
 - chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
 - frequent **chest pain**
- These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days

Drug Facts (continued)

- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- swallow whole. Do not chew or crush tablets.

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

Drug Facts (continued)

Inactive ingredients

carnauba wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate

Questions or comments?

1-800-719-9260

3 Z BOTTLES INSIDE

Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

Safety Feature - Do not use if printed seal under cap is broken or missing.

NDC 0000-000-00

Compare to Prilosec OTC®*
Bonus! 14 Free

DIE CUT WINDOW

DIE CUT WINDOW

DIE CUT WINDOW

Treats **FREQUENT** Heartburn!



Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

42 ~~28~~ TABLETS

Three ~~two~~ 14-day courses of treatment
May take 1 to 4 days for full effect



*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

915D7 FA C12-E 100318

Location for lot number and expiration date

STANDARD FORMAT

TYPE SIZES:

TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed

HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed

TEXT: 6 pt Helvetica Neue LT Std 57 Condensed

LEADING: 6.5 pt

BULLETS: 5 pt Square with 2 M'S spacing between statements

TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed


HEAVY LINES: 1.5 pt

HAIRLINES: 0.5 pt

Hairlines extend to within two spaces of the "Drug Facts" box

Location for lot number and expiration date

- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking



Drug Facts

Active ingredient (in each tablet)
Omeprazole 20 mg.....Acid reducer

Purpose
Acid reducer

Use
treats frequent heartburn (occurs *2 or more* days a week)
not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:
trouble or pain swallowing food, vomiting with blood, or bloody or black stools
heartburn with lightheadedness, sweating or dizziness
chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
request chest pain
These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:
had heartburn over 3 months. This may be a sign of a more serious condition.
frequent wheezing, particularly with heartburn
unexplained weight loss
nausea or vomiting
stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:
your heartburn continues or worsens
you need to take this product for more than 14 days
you need to take more than 1 course of treatment every 4 months
you get diarrhea
you develop a rash or joint pain

Drug Facts (continued)

Directions
for adults 18 years of age and older
this product is to be used once a day, every 24 hours, every day for 14 days
it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
swallow 1 tablet with a glass of water before eating in the morning
take every day for 14 days
do not take more than 1 tablet a day

do not use for more than 14 days unless directed by your doctor

do not use for more than 14 days unless directed by your doctor

swallow whole. Do not chew or crush tablets.

Repeated 14-Day Courses (if needed)
you may repeat a 14-day course every 4 months
do not take for more than 14 days or more often than every 4 months unless directed by a doctor

children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information
read the directions and warnings before use
keep the carton. It contains important information.
store at 20-25°C (68-77°F) and protect from moisture

Inactive ingredients camuaba wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-719-9260

91555 FA C18-D 100318

Omeprazole

Delayed Release Tablets 20 mg


Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakdima Street, Yokneam 2089200, Israel
*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Safety Feature - Do not use if printed tablet blister unit is open or torn.



Omeprazole

Delayed Release Tablets 20 mg



NDC 0000-000-00

Compare to Prilosec OTC®*
Bonus Pack! 14 Free

Treats **FREQUENT** Heartburn!



Omeprazole

Delayed Release Tablets 20 mg

Acid Reducer

42 ~~28~~ TABLETS

Three ~~Two~~ 14-day courses of treatment
May take 1 to 4 days for full effect



STANDARD FORMAT

TYPE SIZES:

TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed

HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed

TEXT: 6 pt Helvetica Neue LT Std 57 Condensed

LEADING: 6.5 pt

BULLETS: 5 pt Square

TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed

HEAVY LINES: 1.5 pt

HAIRLINES: 0.5 pt

Hairlines extend to within two spaces of the "Drug Facts" box

Drug Facts

Active ingredient (in each tablet) Purpose
Omeprazole 20 mg.....Acid reducer

Use
■ treats frequent heartburn (occurs **2 or more** days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to omeprazole
Do not use if you have:
■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
■ heartburn with **lightheadedness, sweating or dizziness**
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
■ frequent **chest pain**
These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss
■ nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:
■ your heartburn continues or worsens
■ you need to take this product for more than 14 days

Drug Facts (continued)

■ you need to take more than 1 course of treatment every 4 months
■ you get diarrhea
■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
■ for adults 18 years of age and older
■ this product is to be used once a day (every 24 hours), every day for 14 days
■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
■ swallow 1 tablet with a glass of water before eating in the morning
■ take every day for 14 days
■ do not take more than 1 tablet a day
■ do not use for more than 14 days unless directed by your doctor
■ swallow whole. Do not chew or crush tablets.

Repeated 14-Day Courses (if needed)
■ you may repeat a 14-day course every 4 months
■ **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information
■ read the directions and warnings before use
■ keep the carton. It contains important information.
■ store at 20-25°C (68-77°F) and protect from moisture

Drug Facts (continued)

Inactive ingredients
carnauba wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate

Questions or comments?
1-800-719-9260

3 BOTTLES INSIDE
Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

Safety Feature - Do not use if printed seal under cap is broken or missing.

NDC 0000-000-00

Compare to Prilosec OTC®*

Bonus Pack! 14 Free

DIE CUT WINDOW

DIE CUT WINDOW

DIE CUT WINDOW

Treats **FREQUENT** Heartburn! **24 HR**

Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

42 28 TABLETS

Three ~~Two~~ 14-day courses of treatment
May take 1 to 4 days for full effect



Tips for Managing Heartburn
■ Do not lie flat or bend over after eating
■ Do not wear tight-fitting clothing around the stomach
■ Do not eat before bedtime
■ Raise the head of your bed
■ Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
■ Eat slowly and avoid big meals
■ If overweight, lose weight
■ Quit smoking



*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Location for lot number and expiration date

915D7 FA C12-D 100318

STANDARD FORMAT
TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

Location for lot number and expiration date

- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

Omeprazole

Delayed Release Tablets 20 mg

Made in Israel
 Manufactured by:
Dexcel Pharma Technologies Ltd
 10 Hakdima Street, Yotvata 2069200, Israel
 This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Omeprazole 20 mg</p> <p>Purpose Acid reducer</p> <p>Use Treats frequent heartburn (occurs 2 or more days a week) not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Warnings Allergy alert: Do not use if you are allergic to omeprazole</p> <p>Do not use if you have: black stools trouble or pain swallowing food, vomiting with blood, or bloody or heartburn with lightheadedness, sweating or dizziness chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness</p> <p>requent chest pain These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have: had heartburn over 3 months. This may be a sign of a more serious condition. frequent wheezing, particularly with heartburn unexplained weight loss nausea or vomiting stomach pain</p> <p>Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if: your heartburn continues or worsens you need to take this product for more than 14 days you need to take more than 1 course of treatment every 4 months you get diarrhea you develop a rash or joint pain</p>	<p>Drug Facts (continued)</p> <p>Directions for adults 18 years of age and older this product is to be used once a day (every 24 hours), every day for 14 days it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours</p> <p>14-Day Course of Treatment swallow 1 tablet with a glass of water before eating in the morning take every day for 14 days do not take more than 1 tablet a day</p> <p>Repeated 14-Day Courses (if needed) you may repeat a 14-day course every 4 months do not take for more than 14 days or more often than every 4 months unless directed by a doctor</p> <p>Other information Keep the carton. It contains important information. store at 20-25°C (68-77°F) and protect from moisture</p> <p>Inactive ingredients canabada wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, polyethylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearoyl fumarate, talc, titanium dioxide, triethyl citrate</p> <p>Questions or comments? 1-800-719-9260</p>
---	---

91555 FA C18-G 100318



Omeprazole

Delayed Release Tablets 20 mg

Safety Feature - Do not use if printed tablet blister unit is open or torn.

NDC 0000-000-00

Compare to Prilosec OTC*

Triple Pack
Three 14-day courses of treatment

Treats **FREQUENT Heartburn!**



Omeprazole

Delayed Release Tablets 20 mg
Acid Reducer

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect



STANDARD FORMAT

TYPE SIZES:
 TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued) 8 pt Helvetica Neue LT Std 57 Condensed
 HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
 TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
 LEADING: 6.5 pt
 BULLETS: 5 pt Square
 TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
 HEAVY LINES: 1.5 pt
 HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box

0000000000

0000000000

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Omeprazole 20 mg.....Acid reducer</p> <p>Purpose Acid reducer</p> <p>Use</p> <ul style="list-style-type: none"> ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect <p>Warnings</p> <p>Allergy alert: Do not use if you are allergic to omeprazole</p> <p>Do not use if you have:</p> <ul style="list-style-type: none"> ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath, sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain <p>These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have:</p> <ul style="list-style-type: none"> ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain <p>Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if:</p> <ul style="list-style-type: none"> ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days 	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p> <p>Directions</p> <ul style="list-style-type: none"> ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours <p>14-Day Course of Treatment</p> <ul style="list-style-type: none"> ■ swallow 1 tablet with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ swallow whole. Do not chew or crush tablets. <p>Repeated 14-Day Courses (if needed)</p> <ul style="list-style-type: none"> ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor <p>■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.</p> <p>Other information</p> <ul style="list-style-type: none"> ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F) and protect from moisture
---	---

Drug Facts (continued)

Inactive ingredients
carnauba wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate

Questions or comments?
1-800-719-9260

3 BOTTLES INSIDE
Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

Safety Feature - Do not use if printed seal under cap is broken or missing.

NDC 0000-000-00

Compare to Prilosec OTC®*

Triple Pack Three 14-day courses of treatment

DIE CUT WINDOW **DIE CUT WINDOW** **DIE CUT WINDOW**

Treats **FREQUENT** Heartburn! **24 HR**

Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect



Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

0 000000 000000 0

915D7 FA C12-G 092818

Location for lot number and expiration date

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

NDC 0000-000-00

Compare to Prilosec OTC®*

3 PACK

Three 14-day courses of treatment

Treats **FREQUENT** Heartburn!



Omeprazole

Delayed Release Tablets 20 mg

Acid Reducer

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect



91555 FA C6-A 092818

Made in Israel
Manufactured by:
Cescael Pharma Technologies Ltd
10 Haskonia Street, Yokneam 208200, Israel


*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Tips for Managing Heartburn

- Do not eat or drink before bedtime
- Do not wear tight-fitting clothing around the stomach
- Do not lie flat or bend over after eating
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking
- Raise the head of your bed

Drug Facts (continued) Active ingredient (in each tablet) Omeprazole 20 mg	Use ■ Treats frequent heartburn (occurs 2 or more days a week) ■ Not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect ■ For adults 18 years of age and older	Warnings ■ Allergy alert: Do not use if you are allergic to omeprazole ■ Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness ■ frequent chest pain ■ These may be signs of a serious condition. See your doctor. Ask a doctor before use if you have: ■ had heartburn over 2 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ trouble or vomiting ■ stomach pain Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs. Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you get diarrhea	Other information ■ Keep the bottle. It contains important information. ■ Store at 20-25°C (68-77°F) and protect from moisture ■ Inactive ingredients: carnauba wax, ferric oxide red, ferric oxide yellow, polyethylene glycol, polyethylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate	Questions or comments? 1-800-719-9260
Drug Facts 14-Day Course of Treatment ■ Swallow 1 tablet with 8 glasses of water before eating in the morning, take every day for 14 days ■ Do not take more than 1 tablet a day ■ Do not use for more than 14 days unless directed by your doctor ■ Repeat 14-day courses (if needed) ■ You may repeat a 14-day course every 4 months ■ Do not take for more than 14 days or more often than every 4 months unless directed by a doctor ■ Children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.	Directions ■ If you take 1 to 4 days for full effect, some people get complete relief of symptoms within 24 hours ■ For adults 18 years of age and older ■ This product is to be used once a day (every 24 hours), every day for 14 days	Use ■ Treats frequent heartburn (occurs 2 or more days a week) ■ Not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect ■ For adults 18 years of age and older	Other information ■ Keep the bottle. It contains important information. ■ Store at 20-25°C (68-77°F) and protect from moisture ■ Inactive ingredients: carnauba wax, ferric oxide red, ferric oxide yellow, polyethylene glycol, polyethylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate	Questions or comments? 1-800-719-9260

Safety Feature -
Do not use if printed tablet breaks apart or open or torn.



Location for lot number and expiration date

PRINT at 65% for 11-inch x 17-inch size paper

STANDARD FORMAT
TYPE SIZES: TITLE: 10 pt Helvetica Neue LT Std 77 Bold Condensed Oblique TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued) 8 pt Helvetica Neue LT Std 57 Condensed HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique SUBHEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed TEXT: 6 pt Helvetica Neue LT Std 57 Condensed LEADING: 6.7 pt BULLETS: 5 pt Square TELEPHONE #: 9 pt Helvetica Neue LT Std 77 Bold Condensed HEAVY LINES: 2.0 pt HAIRLINES: 0.5 pt Hairlines extend to within two spaces of the "Drug Facts" box

0000000000





Safety Feature - Do not use if printed seal under cap is broken or missing.

NDC 0000-000-00

Compare to Prilosec OTC®*

3 PACK

Three 14-day courses of treatment

DIE CUT WINDOW

DIE CUT WINDOW

DIE CUT WINDOW

Treats FREQUENT Heartburn!
24 HR

Omeprazole

Delayed Release Tablets 20 mg Acid Reducer

42 TABLETS


Three 14-day courses of treatment
May take 1 to 4 days for full effect



Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

0000000000





*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.



915D7 FA C12-A 092818

→ Location for lot number and expiration date

Drug Facts

Active ingredient (in each tablet)
Omeprazole 20 mg.....

Purpose
Acid reducer

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days

Drug Facts (continued)

- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- swallow whole. Do not chew or crush tablets.

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

Drug Facts (continued)

Inactive ingredients

camauaba wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate

Questions or comments?
1-800-719-9260

3 BOTTLES INSIDE

Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

STANDARD FORMAT

TYPE SIZES:

TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

(continued) 8 pt Helvetica Neue LT Std 57 Condensed

HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed

TEXT: 6 pt Helvetica Neue LT Std 57 Condensed

LEADING: 6.5 pt

BULLETS: 5 pt Square with 2 M'S spacing between statements

TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed

HEAVY LINES: 1.5 pt

HAIRLINES: 0.5 pt

Hairlines extend to within two spaces of the "Drug Facts" box

NDC 0000-000-00

Compare to Prilosec OTC®*

Value Pack!
3-14 Tablet Cartons

Treats **FREQUENT** Heartburn!



Omeprazole

Delayed Release Tablets 20 mg

Acid Reducer

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect



01656 FA C6-J 092818


Made in Israel
Manufactured by:
Cexcel Pharma Technologies Ltd
10 Haddana Street, Yokneam 208200, Israel

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking
- Raise the head of your bed

Drug Facts Active ingredient (in each tablet) Omeprazole 20 mg Purpose Acid reducer	Use Treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect ■ Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or burning with sweating, or lightheadedness ■ spreading to arms, neck or shoulders, or lightheadedness ■ frequent chest pain These may be signs of a serious condition. See your doctor. Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs. Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take the product for more than 14 days ■ you get dizziness
Warnings Allergy alert: Do not use if you are allergic to omeprazole Do not use if you have: ■ 14-day course of treatment ■ swallow 1 tablet with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ swallow whole. Do not chew or crush tablets. Repeat 14-Day Course (if needed) ■ You may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor These may be signs of a serious condition. See your doctor. ■ children under 18 years of age; ask a doctor; heartburn in children may sometimes be caused by a serious condition.	Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information ■ store at 20-25°C (68-77°F) and protect from moisture Inactive ingredients carnaub wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monobasic potassium phosphate, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium lauryl sulfate, titanium dioxide, triethyl citrate Questions or comments? 1-800-719-9260
Directions ■ Take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours ■ swallow 1 tablet with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ swallow whole. Do not chew or crush tablets. Repeat 14-Day Course (if needed) ■ You may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor These may be signs of a serious condition. See your doctor. ■ children under 18 years of age; ask a doctor; heartburn in children may sometimes be caused by a serious condition.	Drug Facts (continued) If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Safety Feature - Do not use if blister unit is open or torn.



Location for lot number and expiration date

PRINT at 65% for 11-inch x 17-inch size paper

STANDARD FORMAT
TYPE SIZES:
TITLE: 10 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued)
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
HEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 9.7 pt
BULLETS: 5 pt Square
TELEPHONE #: 9 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 2.0 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

Drug Facts

Active ingredient (in each tablet)
Omeprazole 20 mg.....Acid reducer

Purpose

Use
■ treats frequent heartburn (occurs **2 or more** days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
 - heartburn with **lightheadedness, sweating or dizziness**
 - chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
 - frequent **chest pain**
- These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days

Drug Facts (continued)

- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- swallow whole. Do not chew or crush tablets.

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

Drug Facts (continued)

Inactive ingredients

carnauba wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate

Questions or comments?

1-800-719-9260

3 BOTTLES INSIDE

Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

Safety Feature - Do not use if printed seal under cap is broken or missing.

NDC 0000-000-00

Compare to Prilosec OTC®*

Value Pack! 3-14 Tablet Bottles

DIE CUT WINDOW

DIE CUT WINDOW

DIE CUT WINDOW

Treats **FREQUENT** Heartburn! **24 HR**

Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect



- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking



*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Location for lot number and expiration date

915D7 FA C12-J 092818

STANDARD FORMAT

- TYPE SIZES:**
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

Location for
lot number and
expiration date

0927 18

Made in Israel

: 401D5 FA F4

Manufactured by:
Dexcel Pharma Technologies Ltd
10 Haldrama Street, Yokneam 2069200, Israel



WILDBERRY
MINT
Coated Tablet

14 TABLETS

One 14-day course of treatment
May take 1 to 4 days for full effect

NOT FOR RESALE

Treats FREQUENT Heartburn! 24 HR

Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

SWALLOW - DO NOT CHEW

Safety Feature – Do not use if printed seal under cap is broken or missing. This label does not contain full product information. See carton for complete information. Read warnings and directions on carton before use. Retain carton for reference.

Drug Facts

Active ingredient (in each tablet) Purpose
Omeprazole 20 mg.... Acid reducer

Use ■ treats frequent heartburn (occurs 2 or more days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect.

Other Information

■ read the directions and warnings before use. ■ keep the carton. It contains important information.
■ store at 20-25°C (68-77°F) and protect from moisture.

Questions? 1-800-719-9280

NON-DFL FORMAT

TYPE SIZES:

HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

SUBHEADINGS: 7 pt Helvetica Neue LT Std 77 Bold Condensed

TEXT: 6 pt Helvetica Neue LT Std 57 Condensed

LEADING: 6 pt

BULLETS: 5 pt Square with 2 M's spacing between statements

TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed

HEAVY LINES: 1.0 pt

HAIRLINES LINES: 0.5 pt

Batch No.

EXP

40174 00 VS 092718

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Omeprazole 20 mg. Acid reducer.</p> <p>Purpose Acid reducer.</p>	<p>Use Treats frequent heartburn (occurs <i>2 or more</i> days a week). Not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect.</p> <p>Warnings Allergy alert: Do not use if you are allergic to omeprazole.</p> <p>Do not use if you have: ● trouble or pain swallowing food; vomiting with blood, or bloody or black stools ● heartburn with lightheadedness, sweating or dizziness ● chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness</p> <p>These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have: ● frequent wheezing, particularly with heartburn ● had heartburn over 3 months. This may be a sign of a more serious condition.</p> <p>Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if: ● your heartburn continues or worsens ● you need to take this product for more than 14 days ● you need to take more than 1 course of treatment every 4 months ● you get diarrhea ● you develop a rash or joint pain</p>
<p>Drug Facts (continued)</p> <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p> <p>Directions ● for adults 18 years of age and older ● this product is to be used once a day (every 24 hours), every day for 14 days ● if you take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours</p> <p>14-Day Course of Treatment ● swallow 1 tablet with a glass of water before eating in the morning ● take every day for 14 days ● do not take more than 1 tablet a day unless directed by your doctor.</p> <p>Repeated 14-Day Courses (if needed) ● do not take more than 14 days or more often than every 4 months ● you may repeat a 14-day course every 4 months ● do not take for more than 14 days or more often than every 4 months unless directed by a doctor</p> <p>Other information ● children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.</p> <p>Read the directions and warnings before use ● keep the carton. It contains important information. ● store at 20-25°C (68-77°F) and protect from moisture</p> <p>Inactive ingredients blue #2/indigo carmine aluminum lake, flavor, hypromellose, hypromellose acetate succinate, lactose monohydrate, mannitol, modified starch, monothenediolamine, polyethylene glycol 3350, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, titanium dioxide, triacetin, triethyl citrate</p> <p>Questions or comments? 1-800-719-9260</p>	<p>Drug Facts</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p> <p>Directions ● for adults 18 years of age and older ● this product is to be used once a day (every 24 hours), every day for 14 days ● if you take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours</p> <p>14-Day Course of Treatment ● swallow 1 tablet with a glass of water before eating in the morning ● take every day for 14 days ● do not take more than 1 tablet a day unless directed by your doctor.</p> <p>Repeated 14-Day Courses (if needed) ● do not take for more than 14 days or more often than every 4 months ● you may repeat a 14-day course every 4 months ● do not take for more than 14 days or more often than every 4 months unless directed by a doctor</p> <p>Other information ● children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.</p> <p>Read the directions and warnings before use ● keep the carton. It contains important information. ● store at 20-25°C (68-77°F) and protect from moisture</p> <p>Inactive ingredients blue #2/indigo carmine aluminum lake, flavor, hypromellose, hypromellose acetate succinate, lactose monohydrate, mannitol, modified starch, monothenediolamine, polyethylene glycol 3350, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, titanium dioxide, triacetin, triethyl citrate</p> <p>Questions or comments? 1-800-719-9260</p>

Location for lot number and expiration date

NOT FOR RESALE

Safety Feature – Do not use if printed tablet blister unit is open or torn.

Compare to Prilosec OTC®*

Treats **FREQUENT** Heartburn!



Omeprazole

Delayed Release Tablets 20 mg
Acid Reducer



Wildberry Mint Coated Tablet

SWALLOW – DO NOT CHEW

14 TABLETS

One 14-day course of treatment
May take 1 to 4 days for full effect



Actual Size

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued)
 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Questions or comments? 1-800-719-9260

Inactive ingredients benzyl alcohol, carmine, carnauba wax, FD&C sodium stearyl fumarate, sucralose, talc, titanium dioxide, triacetin, triethyl citrate, polyethylene glycol 3350, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, succinate, lactose monohydrate, mannitol, modified starch, monostearatamide, blue #2/indigo carmine aluminum lake, flavor, hypromellose, hydroxyethyl acetate

Other information
 ■ read the directions and warnings before use
 ■ keep the carton. It contains important information.
 ■ store at 20-25°C (68-77°F) and protect from moisture

14-Day Course of Treatment
 ■ swallow 1 tablet with a glass of water before eating in the morning
 ■ take every day for 14 days
 ■ do not take more than 1 tablet a day
 ■ do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)
 ■ swallow whole. Do not chew, crush, or suck tablets.
 ■ do not use for more than 14 days unless directed by your doctor
 ■ you may repeat a 14-day course every 4 months
 ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

Directions
 ■ for adults 18 years of age and older
 ■ this product is to be used once a day (every 24 hours), every day for 14 days
 ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

Warnings
 ■ Do not use if you are allergic to omeprazole
 ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
 ■ heartburn with lightheadedness, sweating or dizziness
 ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Use
 ■ treats frequent heartburn (occurs 2 or more days a week)
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Drug Facts (continued)
Purpose Acid reducer
Active ingredient (in each tablet) Omeprazole 20 mg
Warnings Allergy alert: Do not use if you are allergic to omeprazole
Do not use if you have:
 ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
 ■ heartburn with lightheadedness, sweating or dizziness
 ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Ask a doctor before use if you have:
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss
 ■ nausea or vomiting
 ■ stomach pain
Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea
 ■ you develop a rash or joint pain

Made in Israel
 Manufactured by:
Cephalon Pharma Technologies Ltd
 10 Hakdima Street, Yokneam 2069200, Israel

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®

40174 FA C6 100218

Location for lot number and expiration date



NDC 0000-000-00

Omeprazole

Delayed Release Tablets 20 mg

Safety Feature - Do not use if printed tablet blister unit is open or torn.

Compare to Prilosec OTC®*

Treats **FREQUENT** Heartburn!



Omeprazole

Delayed Release Tablets 20 mg

Acid Reducer



SWALLOW - DO NOT CHEW

14 TABLETS

One 14-day course of treatment
 May take 1 to 4 days for full effect



WILDBERRY MINT
Coated Tablet

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box



Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- **swallow whole. Do not chew, crush, or suck tablets.**

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**

- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Drug Facts (continued)

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

Inactive ingredients benzyl alcohol, carmine, carnauba wax, FD&C blue #2/indigo carmine aluminum lake, flavor, hypromellose, hypromellose acetate succinate, lactose monohydrate, menthol, modified starch, monoethanolamine, polyethylene glycol 3350, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments? 1-800-719-9260

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight ■ Quit smoking

1 BOTTLE INSIDE

Made in Israel
 Manufactured by:
Dexcel Pharma Technologies Ltd
 10 Hakidma Street, Yokneam 2069200, Israel

NDC 0000-000-00

Compare to Prilosec OTC®*

DIE CUT WINDOW

Treats **FREQUENT** Heartburn! **24 HR**

Omeprazole
 Delayed Release Tablets 20 mg
 Acid Reducer

SWALLOW - DO NOT CHEW

14 TABLETS

One 14-day course of treatment
 May take 1 to 4 days for full effect

WILDBERRY MINT
 Coated Tablet



*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

401D5 FA C6 092718

Drug Facts

Active ingredient (in each tablet)
 Omeprazole 20 mg.....Acid reducer

Use
 ■ treats frequent heartburn (occurs **2 or more** days a week)
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:
 ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
 ■ heartburn with **lightheadedness, sweating or dizziness**
 ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
 ■ frequent **chest pain**
 These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss
 ■ nausea or vomiting ■ stomach pain

Location for lot number and expiration date

STANDARD FORMAT

TYPE SIZES:

- TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique**
- TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued)**
- 8 pt Helvetica Neue LT Std 57 Condensed**
- HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique**
- SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed**
- TEXT: 6 pt Helvetica Neue LT Std 57 Condensed**
- LEADING: 6.5 pt**
- BULLETS: 5 pt Square with 2 M'S spacing between statements**
- TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed**
- HEAVY LINES: 1.5 pt**
- HAIRLINES: 0.5 pt**
- Hairlines extend to within two spaces of the "Drug Facts" box**

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Questions or comments? 1-800-719-9260

Inactive ingredients benzyl alcohol, carmine, camellia wax, FD&C sodium stearyl fumarate, fumarate, succinylsorbate, talc, titanium dioxide, tracetin, triethyl citrate, polyethylene glycol 3350, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sucralose, tacrolimus monohydrate, menthyl, modified starch, monothenediamine, blue #2/indigo carmine aluminum lake, flavor, hypromellose, hypromellose acetate

Other information
 ■ Read the directions and warnings before use
 ■ Keep the carton. It contains important information.
 ■ Store at 20-25°C (68-77°F) and protect from moisture

Directions
 ■ For adults 18 years of age and older
 ■ This product is to be used once a day (every 24 hours), every day for 14 days
 ■ It may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
 ■ Swallow 1 tablet with a glass of water before eating in the morning
 ■ Take every day for 14 days
 ■ Do not take more than 1 tablet a day
 ■ Do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)
 ■ Swallow whole. Do not chew, crush, or suck tablets.
 ■ Do not use for more than 14 days unless directed by your doctor

Do not take for more than 14 days or more often than every 4 months unless directed by a doctor
 ■ Children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Drug Facts (continued)
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Purpose
 Acid reducer

Use
 ■ Treats frequent heartburn (occurs 2 or more days a week)
 ■ Not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:
 ■ Trouble or pain swallowing food, vomiting with blood, or bloody or black stools
 ■ Heartburn or high blood pressure, sweating or dizziness
 ■ Chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Ask a doctor before use if you have:
 ■ Frequent chest pain
 ■ These may be signs of a serious condition. See your doctor.

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.
 ■ Stomach pain
 ■ Nausea or vomiting
 ■ Unexplained weight loss
 ■ Frequent heartburn, particularly with heartburn

Stop use and ask a doctor if:
 ■ Your heartburn continues or worsens
 ■ You need to take this product for more than 14 days
 ■ You need to take more than 1 course of treatment every 4 months
 ■ You develop a rash or joint pain

Made in Israel
 Manufactured by:
Celexel Pharma Technologies Ltd
 10 Hahadna Street, Yokneam 2069200, Israel

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

40174 FA C6-F 100218

Location for lot number and expiration date



Omeprazole
 Delayed Release Tablets 20 mg

Safety Feature - Do not use if printed tablet blister unit is open or torn.

NDC 0000-000-00

Compare to Prilosec OTC®
FDA Approved

Treats **FREQUENT** Heartburn!



Omeprazole
 Delayed Release Tablets 20 mg
 Acid Reducer

SWALLOW - DO NOT CHEW

14 TABLETS

One 14-day course of treatment
 May take 1 to 4 days for full effect



WILDBERRY MINT
 Coated Tablet



STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box



Safety Feature - Do not use if printed seal under cap is broken or missing.

NDC 0000-000-00

Compare to Prilosec OTC®*
FDA Approved

DIE CUT WINDOW

Treats FREQUENT Heartburn! 24 HR

Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

14 TABLETS

One 14-day course of treatment
May take 1 to 4 days for full effect



WILDBERRY MINT
Coated Tablet



*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

401D5 FA C6-F 092718

Drug Facts	Purpose
Active ingredient (in each tablet) Omeprazole 20 mg.....	Acid reducer

Use
 ■ treats frequent heartburn (occurs 2 or more days a week)
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:
 ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
 ■ heartburn with **lightheadedness, sweating or dizziness**
 ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
 ■ frequent **chest pain**
 These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss
 ■ nausea or vomiting ■ stomach pain

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea
 ■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
 ■ for adults 18 years of age and older
 ■ this product is to be used once a day (every 24 hours), every day for 14 days
 ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
 ■ swallow 1 tablet with a glass of water before eating in the morning
 ■ take every day for 14 days
 ■ do not take more than 1 tablet a day
 ■ do not use for more than 14 days unless directed by your doctor
 ■ **swallow whole. Do not chew, crush, or suck tablets.**

Repeated 14-Day Courses (if needed)
 ■ you may repeat a 14-day course every 4 months
 ■ **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**

■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Drug Facts (continued)

Other information
 ■ read the directions and warnings before use
 ■ keep the carton. It contains important information.
 ■ store at 20-25°C (68-77°F) and protect from moisture

Inactive ingredients benzyl alcohol, carmine, carnauba wax, FD&C blue #2/indigo carmine aluminum lake, flavor, hypromellose, hypromellose acetate succinate, lactose monohydrate, menthol, modified starch, monoethanolamine, polyethylene glycol 3350, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments? 1-800-719-9260


Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight ■ Quit smoking

1 BOTTLE INSIDE
 Made in Israel
 Manufactured by:
Dexcel Pharma Technologies Ltd
 10 Hakidma Street, Yokneam 2069200, Israel

STANDARD FORMAT

- TYPE SIZES:**
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box



- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

Questions or comments? 1-800-719-9260

Inactive ingredients: benzyl alcohol, carmine, carnuba wax, FD&C blue #2/indigo carmine aluminum lake, flavor, hypromellose, hydroxypropyl methylcellulose, lactose monohydrate, menthol, modified starch, monoethanolamine, polyethylene glycol 3350, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, titanium dioxide, tracefin, triethyl citrate

Other information

- store at 20-25°C (68-77°F) and protect from moisture
- keep the carton, it contains important information
- read the directions and warnings before use

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- swallow whole. Do not chew, crush, or suck tablets.
- chest pain or shoulder pain with shortness of breath; sweating; pain
- heartburn with lightheadedness, sweating or dizziness
- trouble or pain swallowing food, vomiting with blood, or bloody or black stools

Do not use if you have:

- frequent chest pain
- these may be signs of a serious condition. See your doctor.

Warnings

- Allergy alert: Do not use if you are allergic to omeprazole
- Do not use if you have:

Use

- treats frequent heartburn (occurs *2 or more* days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Drug Facts

Active ingredient (in each tablet).....acid reducer

Purpose

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- swallow whole. Do not chew, crush, or suck tablets.
- chest pain or shoulder pain with shortness of breath; sweating; pain
- heartburn with lightheadedness, sweating or dizziness
- trouble or pain swallowing food, vomiting with blood, or bloody or black stools

Do not use if you have:

- frequent chest pain
- these may be signs of a serious condition. See your doctor.

Warnings

- Allergy alert: Do not use if you are allergic to omeprazole
- Do not use if you have:

Use

- treats frequent heartburn (occurs *2 or more* days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Drug Facts

Active ingredient (in each tablet).....acid reducer

Purpose

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- swallow whole. Do not chew, crush, or suck tablets.
- chest pain or shoulder pain with shortness of breath; sweating; pain
- heartburn with lightheadedness, sweating or dizziness
- trouble or pain swallowing food, vomiting with blood, or bloody or black stools

Do not use if you have:

- frequent chest pain
- these may be signs of a serious condition. See your doctor.

Warnings

- Allergy alert: Do not use if you are allergic to omeprazole
- Do not use if you have:

Use

- treats frequent heartburn (occurs *2 or more* days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Drug Facts

Active ingredient (in each tablet).....acid reducer

Purpose

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- swallow whole. Do not chew, crush, or suck tablets.
- chest pain or shoulder pain with shortness of breath; sweating; pain
- heartburn with lightheadedness, sweating or dizziness
- trouble or pain swallowing food, vomiting with blood, or bloody or black stools

Do not use if you have:

- frequent chest pain
- these may be signs of a serious condition. See your doctor.

Warnings

- Allergy alert: Do not use if you are allergic to omeprazole
- Do not use if you have:

Use

- treats frequent heartburn (occurs *2 or more* days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Omeprazole


Delayed Release Tablets 20 mg

Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakdima Street, Yokneam 2069200, Israel
*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

40130 FA C6 100318

Location for lot number and expiration date

Safety Feature - Do not use if printed tablet blister unit is open or torn.



Omeprazole

Delayed Release Tablets 20 mg

NDC 0000-000-00-00

Compare to Prilosec OTC®*

Treats **FREQUENT** Heartburn!



Omeprazole

Delayed Release Tablets 20 mg

Acid Reducer

SWALLOW - DO NOT CHEW

28 TABLETS

Two 14-day courses of treatment
May take 1 to 4 days for full effect




WILDBERRY MINT
Coated Tablet

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

000000000



000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

Drug Facts

Active ingredient (in each tablet) Purpose
Omeprazole 20 mg.....Acid reducer

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating, pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens

Drug Facts (continued)

- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea ■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- swallow whole. Do not chew, crush, or suck tablets.

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor

■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition. ▶

Drug Facts (continued)

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

Inactive ingredients: benzyl alcohol, carmine, carnauba wax, FD&C blue #2/indigo carmine aluminum lake, flavor, hypromellose, hypromellose acetate succinate, lactose monohydrate, menthol, modified starch, monoethanolamine, polyethylene glycol 3350, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments? 1-800-719-9260

2 BOTTLES INSIDE

Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

NDC 0000-000-00

Compare to Prilosec OTC®*

DIE CUT WINDOW

DIE CUT WINDOW

Treats **FREQUENT** Heartburn!

24 HR

Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

SWALLOW - DO NOT CHEW

28 TABLETS

Two 14-day courses of treatment
May take 1 to 4 days for full effect



WILDBERRY MINT
Coated Tablet

- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

Safety Feature - Do not use if printed seal under cap is broken or missing.



*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.


401D6 FA C6 092718

Location for lot number and expiration date

STANDARD FORMAT

TYPE SIZES:

- TITLE:** 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
- TITLE (continued):** 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
- (continued)** 8 pt Helvetica Neue LT Std 57 Condensed
- HEADINGS:** 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
- SUBHEADINGS:** 6 pt Helvetica Neue LT Std 77 Bold Condensed
- TEXT:** 6 pt Helvetica Neue LT Std 57 Condensed
- LEADING:** 6.5 pt
- BULLETS:** 5 pt Square with 2 M'S spacing between statements
- TELEPHONE #:** 6 pt Helvetica Neue LT Std 77 Bold Condensed
- HEAVY LINES:** 1.5 pt
- HAIRLINES:** 0.5 pt
- Hairlines extend to within two spaces of the "Drug Facts" box**



- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

Questions or comments? 1-800-719-9260

Inactive ingredients benzyl alcohol, carnine, carnauba wax, FD&C blue #2/indigo carmine aluminum lake, flavor, hypromellose, hypromellose acetate succinate, lactose monohydrate, menthol, modified starch, monoethanolamine, polyethylene glycol 3350, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, titanium dioxide, tracetin, triethyl citrate

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take for more than 14 days unless directed by your doctor
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- swallow whole. Do not chew, crush, or suck tablets.
- do not use for more than 14 days unless directed by your doctor
- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor

children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain

These may be signs of a serious condition. See your doctor.

Warnings

Allergy alert: Do not use if you are allergic to omeprazole

Use

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Drug Facts

Active ingredient (in each tablet) Omeprazole 20 mg.....Acid reducer

Purpose Acid reducer

40130 FA C6-F 100318


Location for lot number and expiration date

Omeprazole

Delayed Release Tablets 20 mg

Made in Israel
Manufactured by:
Dececal Pharma Technologies Ltd
10 Hakdima Street, Yokneam 2069200, Israel

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.



Omeprazole

Delayed Release Tablets 20 mg

Safety Feature - Do not use if printed tablet blister unit is open or torn.

NDC 0000-000-00

Compare to Prilosec OTC®*
FDA Approved

Treats **FREQUENT** Heartburn!



Omeprazole

Delayed Release Tablets 20 mg

Acid Reducer

SWALLOW - DO NOT CHEW

28 TABLETS

Two 14-day courses of treatment
May take 1 to 4 days for full effect




WILDBERRY MINT
Coated Tablet


STANDARD FORMAT

- TYPE SIZES:**
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

000000000



000000000



000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000

000000000



000000000

000000000

000000000

000000000

Safety Feature - Do not use if printed seal under cap is broken or missing.



NDC 0000-000-00

Compare to Prilosec OTC[®]*

FDA Approved

DIE CUT WINDOW

DIE CUT WINDOW

Treats FREQUENT Heartburn!

24 HR

Omeprazole Delayed Release Tablets 20 mg Acid Reducer

SWALLOW - DO NOT CHEW

28 TABLETS

Two 14-day courses of treatment May take 1 to 4 days for full effect



WILDBERRY MINT Coated Tablet

- Tips for Managing Heartburn
Do not lie flat or bend over after eating
Do not wear tight-fitting clothing around the stomach
Do not eat before bedtime
Raise the head of your bed
Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
Eat slowly and avoid big meals
If overweight, lose weight
Quit smoking

Drug Facts (continued)

- Other information
read the directions and warnings before use
Keep the carton. It contains important information.
store at 20-25°C (68-77°F) and protect from moisture

Inactive ingredients benzyl alcohol, carmine, carnauba wax, FD&C blue #2/indigo carmine aluminum lake, flavor, hypromellose, hypromellose acetate succinate, lactose monohydrate, menthol, modified starch, monoethanolamine, polyethylene glycol 3350, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments? 1-800-719-9260

2 BOTTLES INSIDE
Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

Drug Facts

Active ingredient (in each tablet) Purpose
Omeprazole 20 mg.....Acid reducer

- Use
treats frequent heartburn (occurs 2 or more days a week)
not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to omeprazole

- Do not use if you have:
trouble or pain swallowing food, vomiting with blood, or bloody or black stools
heartburn with lightheadedness, sweating or dizziness
chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
frequent chest pain
These may be signs of a serious condition. See your doctor.

- Ask a doctor before use if you have:
had heartburn over 3 months. This may be a sign of a more serious condition.
frequent wheezing, particularly with heartburn
unexplained weight loss
nausea or vomiting
stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:
your heartburn continues or worsens

Drug Facts (continued)

- you need to take this product for more than 14 days
you need to take more than 1 course of treatment every 4 months
you get diarrhea
you develop a rash or joint pain

if pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
for adults 18 years of age and older
this product is to be used once a day (every 24 hours), every day for 14 days
it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
swallow 1 tablet with a glass of water before eating in the morning
take every day for 14 days
do not take more than 1 tablet a day
do not use for more than 14 days unless directed by your doctor
swallow whole. Do not chew, crush, or suck tablets.

Repeated 14-Day Courses (if needed)
you may repeat a 14-day course every 4 months
do not take for more than 14 days or more often than every 4 months unless directed by a doctor
children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

STANDARD FORMAT

- TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box



*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC[®].

401D6 FA C6-F 092718

Location for lot number and expiration date



- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

Questions or comments? 1-800-719-9260

Inactive ingredients benzyl alcohol, carmine, carnuba wax, FD&C blue #2/indigo carmine aluminum lake, flavor, hypromellose, hypromellose acetate succinate, lactose monohydrate, menthol, modified starch, monoethanolamine, polyethylene glycol 3350, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, titanium dioxide, tracetin, triethyl citrate

Other information

- store at 20-25°C (68-77°F) and protect from moisture
- keep the carton. It contains important information.
- read the directions and warnings before use

■ sometimes be caused by a serious condition.

■ children under 18 years of age: ask a doctor. Heartburn in children may

4 months unless directed by a doctor

■ do not take for more than 14 days or more often than every

Repeated 14-Day Courses (if needed)

■ swallow whole. Do not chew, crush, or suck tablets.

■ do not use for more than 14 days unless directed by your doctor

■ do not take more than 1 tablet a day

■ take every day for 14 days

■ swallow 1 tablet with a glass of water before eating in the morning

14-Day Course of Treatment

■ symptoms within 24 hours

■ it may take 1 to 4 days for full effect; some people get complete relief of

■ this product is to be used once a day (every 24 hours), every day for 14 days

■ for adults 18 years of age and older

Directions

■ Keep out of reach of children. In case of overdose, get medical help or

contact a Poison Control Center right away (1-800-222-1222).

Drug Facts (continued)

Use

Omeprazole 20 mg.....acid reducer

Purpose

■ not intended for immediate relief of heartburn; this drug may take 1 to

4 days for full effect

Warnings

■ Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:

■ trouble or pain swallowing food, vomiting with blood, or bloody or

black stools

■ heartburn with lightheadedness, sweating or dizziness

■ chest pain or shoulder pain with shortness of breath; sweating; pain

spreading to arms, neck or shoulders; or lightheadedness

■ frequent chest pain

■ These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

■ had heartburn over 3 months. This may be a sign of a more serious

condition.

■ frequent wheezing, particularly with heartburn

■ unexplained weight loss

■ nausea or vomiting

■ stomach pain

Stop use and ask a doctor if:

■ your heartburn continues or worsens

■ you need to take this product for more than 14 days

■ you need to take more than 1 course of treatment every 4 months

■ you get diarrhea

■ you develop a rash or joint pain

Location for lot number and expiration date

40130 FA C6-B 100318

Omeprazole
Delayed Release Tablets 20 mg

Made in Israel
Manufactured by:
Cexcel Pharma Technologies Ltd
10 Hakdima Street, Yokneam 2099200, Israel
This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC.

Safety Feature - Do not use if printed tablet blister unit is open or torn.



Omeprazole
Delayed Release Tablets 20 mg

Compare to Prilosec OTC®*
14 FREE

Treats **FREQUENT** Heartburn!



Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

SWALLOW - DO NOT CHEW

28 TABLETS

Two 14-day courses of treatment
May take 1 to 4 days for full effect



WILDBERRY MINT
Coated Tablet

STANDARD FORMAT

TYPE SIZES:

TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed

HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed

TEXT: 6 pt Helvetica Neue LT Std 57 Condensed

LEADING: 6.5 pt

BULLETS: 5 pt Square


TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed

HEAVY LINES: 1.5 pt

HAIRLINES: 0.5 pt


Hairlines extend to within two spaces of the "Drug Facts" box

00000000



00000000

00000000



00000000

00000000

Drug Facts

Active ingredient (in each tablet) Purpose
Omeprazole 20 mg.....Acid reducer

Use
■ treats frequent heartburn (occurs **2 or more** days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:
■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
■ heartburn with **lightheadedness, sweating or dizziness**
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
■ frequent **chest pain**
These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss
■ nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:
■ your heartburn continues or worsens

Drug Facts (continued)

■ you need to take this product for more than 14 days
■ you need to take more than 1 course of treatment every 4 months
■ you get diarrhea ■ you develop a rash or joint pain
If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
■ for adults 18 years of age and older
■ this product is to be used once a day (every 24 hours), every day for 14 days
■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
■ swallow 1 tablet with a glass of water before eating in the morning
■ take every day for 14 days
■ do not take more than 1 tablet a day
■ do not use for more than 14 days unless directed by your doctor
■ **swallow whole. Do not chew, crush, or suck tablets.**

Repeated 14-Day Courses (if needed)
■ you may repeat a 14-day course every 4 months
■ **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**

■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Drug Facts (continued)

Other information

■ read the directions and warnings before use
■ keep the carton. It contains important information.
■ store at 20-25°C (68-77°F) and protect from moisture

Inactive ingredients benzyl alcohol, carmine, carnauba wax, FD&C blue #2/indigo carmine aluminum lake, flavor, hypromellose, hypromellose acetate succinate, lactose monohydrate, menthol, modified starch, monoethanolamine, polyethylene glycol 3350, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments? 1-800-719-9260

2 BOTTLES INSIDE
Made in Israel
Manufactured by:
Dexel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

Safety Feature - Do not use if printed seal under cap is broken or missing.

NDC 0000-000-00

Compare to Prilosec OTC®*

14 FREE



Treats **FREQUENT** Heartburn!



Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

SWALLOW - DO NOT CHEW

28 TABLETS

Two 14-day courses of treatment
May take 1 to 4 days for full effect



WILDBERRY MINT
Coated Tablet

Tips for Managing Heartburn
■ Do not lie flat or bend over after eating
■ Do not wear tight-fitting clothing around the stomach
■ Do not eat before bedtime
■ Raise the head of your bed
■ Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
■ Eat slowly and avoid big meals
■ If overweight, lose weight
■ Quit smoking



*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Location for lot number and expiration date

401D6 FA C6-B 092718

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box



- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- For adults 18 years of age and older
- This product is to be used once a day (every 24 hours), every day for 14 days
- It may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- do not chew, crush, or suck tablets.

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor

children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

Inactive ingredients benzyl alcohol, carmine, carnuba wax, FD&C blue #2, indigo carmine aluminum lake, flavor, hypromellose, hydroxypropyl methylcellulose, lactose monohydrate, menthol, modified starch, starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments? 1-800-719-9260

Drug Facts

Active ingredient (in each tablet) Omeprazole 20 mg.....Acid reducer

Purpose Acid reducer

Use

- treats frequent heartburn (occurs *2 or more* days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

40130 FA C6-E 100318

Location for lot number and expiration date

Omeprazole

Delayed Release Tablets 20 mg

Made in Israel
 Manufactured by:
Cesceel Pharma Technologies Ltd
 10 Hakdama Street, Yokneam 2089200, Israel
 *This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.



Omeprazole

Delayed Release Tablets 20 mg

Safety Feature - Do not use if printed tablet blister unit is open or torn.

NDC 0000-000-00-00

Compare to Prilosec OTC®*
Bonus! 14 Free

Treats **FREQUENT** Heartburn!



Omeprazole

Delayed Release Tablets 20 mg

Acid Reducer

SWALLOW - DO NOT CHEW

28 ~~14~~ TABLETS

Two ~~One~~ 14-day courses of treatment
 May take 1 to 4 days for full effect



WILDBERRY MINT
 Coated Tablet

STANDARD FORMAT

TYPE SIZES:
 TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued) 8 pt Helvetica Neue LT Std 57 Condensed
 HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
 TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
 LEADING: 6.5 pt
 BULLETS: 5 pt Square
 TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
 HEAVY LINES: 1.5 pt
 HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box

000000000



000000000

Drug Facts

Active ingredient (in each tablet) Purpose
Omeprazole 20 mg.....Acid reducer

Use
■ treats frequent heartburn (occurs **2 or more** days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:
■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
■ heartburn with **lightheadedness, sweating or dizziness**
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
■ frequent **chest pain**
These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss
■ nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:
■ your heartburn continues or worsens

Drug Facts (continued)

■ you need to take this product for more than 14 days
■ you need to take more than 1 course of treatment every 4 months
■ you get diarrhea ■ you develop a rash or joint pain
If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
■ for adults 18 years of age and older
■ this product is to be used once a day (every 24 hours), every day for 14 days
■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
■ swallow 1 tablet with a glass of water before eating in the morning
■ take every day for 14 days
■ do not take more than 1 tablet a day
■ do not use for more than 14 days unless directed by your doctor
■ **swallow whole. Do not chew, crush, or suck tablets.**
Repeated 14-Day Courses (if needed)
■ you may repeat a 14-day course every 4 months
■ **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Drug Facts (continued)


Other information
■ read the directions and warnings before use
■ keep the carton. It contains important information.
■ store at 20-25°C (68-77°F) and protect from moisture

Inactive ingredients benzyl alcohol, carmine, carnauba wax, FD&C blue #2/indigo carmine aluminum lake, flavor, hypromellose, hypromellose acetate succinate, lactose monohydrate, menthol, modified starch, monoethanolamine, polyethylene glycol 3350, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments? 1-800-719-9260


2 BOTTLES INSIDE
Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

000000000



000000000

Safety Feature - Do not use if printed seal under cap is broken or missing.



NDC 0000-000-00

Compare to Prilosec OTC®*
Bonus! 14 Free



Treats **FREQUENT** Heartburn!



Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer
SWALLOW - DO NOT CHEW



WILDBERRY MINT
Coated Tablet

28 TABLETS

Two ~~one~~ 14-day courses of treatment
May take 1 to 4 days for full effect




*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Location for lot number and expiration date

401D6 FA C6-E 100318

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box



- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

Omeprazole
Delayed Release Tablets 20 mg

Made in Israel
Manufactured by:
Chexcel Pharma Technologies Ltd
10 Hakdima Street, Yokneam 2089200, Israel

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Drug Facts (continued)

Active ingredient (in each tablet) Omeprazole 20 mg.....Acid reducer

Purpose Acid reducer

Use
 ■ treats frequent heartburn (occurs 2 or more days a week)
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:
 ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
 ■ heartburn with lightheadedness, sweating or dizziness
 ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

These may be signs of a serious condition. See your doctor.
 ■ frequent chest pain
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.

Ask a doctor before use if you have:
 ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss
 ■ nausea or vomiting
 ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea
 ■ you develop a rash or joint pain

Questions or comments? 1-800-719-9260
 titanium dioxide, triacetin, triethyl citrate
 starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, monoethanolamine, polyethylene glycol 3350, sodium lauryl sulfate, sodium acetate succinate, lactose monohydrate, menthol, modified starch, blue #2/indigo carmine aluminum lake, flavor, hypromellose, hypromellose acetate succinate, benzyl alcohol, carnine, carnauba wax, FD&C

Inactive ingredients

Other information
 ■ keep the carton. It contains important information.
 ■ store at 20-25°C (68-77°F) and protect from moisture
 ■ read the directions and warnings before use

Directions
 ■ for adults 18 years of age and older
 ■ this product is to be used once a day (every 24 hours), every day for 14 days
 ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
 ■ swallow 1 tablet with a glass of water before eating in the morning
 ■ take every day for 14 days

Repeated 14-Day Courses (if needed)
 ■ do not use for more than 14 days unless directed by your doctor
 ■ do not take for more than 14 days unless directed by your doctor
 ■ swallow whole. Do not chew, crush, or suck tablets.

4 months unless directed by a doctor
 ■ you may repeat a 14-day course every 4 months
 ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

40130 FA C6-D 100318

Location for lot number and expiration date



Omeprazole
Delayed Release Tablets 20 mg

Safety Feature - Do not use if printed tablet blister unit is open or torn.

NDC 0000-000-00

Compare to Prilosec OTC®*
Bonus Pack! 14 Free

Treats **FREQUENT** Heartburn!



Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

SWALLOW - DO NOT CHEW

28 ~~14~~ TABLETS

Two ~~One~~ 14-day courses of treatment
May take 1 to 4 days for full effect



WILDBERRY MINT
Coated Tablet

STANDARD FORMAT

TYPE SIZES:

TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued) 8 pt Helvetica Neue LT Std 57 Condensed

HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed

TEXT: 6 pt Helvetica Neue LT Std 57 Condensed

LEADING: 6.5 pt

BULLETS: 5 pt Square

TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed

HEAVY LINES: 1.5 pt

HAIRLINES: 0.5 pt

Hairlines extend to within two spaces of the "Drug Facts" box

00000000



00000000

Drug Facts

Active ingredient (in each tablet) Purpose
Omeprazole 20 mg.....Acid reducer

Use
■ treats frequent heartburn (occurs **2 or more** days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:
■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
■ heartburn with **lightheadedness, sweating or dizziness**
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
■ frequent **chest pain**
These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss
■ nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:
■ your heartburn continues or worsens

Drug Facts (continued)

■ you need to take this product for more than 14 days
■ you need to take more than 1 course of treatment every 4 months
■ you get diarrhea ■ you develop a rash or joint pain
If **pregnant or breast-feeding**, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
■ for adults 18 years of age and older
■ this product is to be used once a day (every 24 hours), every day for 14 days
■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
■ swallow 1 tablet with a glass of water before eating in the morning
■ take every day for 14 days
■ do not take more than 1 tablet a day
■ do not use for more than 14 days unless directed by your doctor
■ **swallow whole. Do not chew, crush, or suck tablets.**
Repeated 14-Day Courses (if needed)
■ you may repeat a 14-day course every 4 months
■ **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Drug Facts (continued)


Other information
■ read the directions and warnings before use
■ keep the carton. It contains important information.
■ store at 20-25°C (68-77°F) and protect from moisture

Inactive ingredients benzyl alcohol, carmine, carnauba wax, FD&C blue #2/indigo carmine aluminum lake, flavor, hypromellose, hypromellose acetate succinate, lactose monohydrate, menthol, modified starch, monoethanolamine, polyethylene glycol 3350, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments? 1-800-719-9260


2 BOTTLES INSIDE
Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

00000000



00000000

Safety Feature - Do not use if printed seal under cap is broken or missing.



NDC 0000-000-00

Compare to Prilosec OTC®*

Bonus Pack! 14 Free



Treats **FREQUENT** Heartburn!



Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer
SWALLOW - DO NOT CHEW



WILDBERRY MINT
Coated Tablet

28 14 TABLETS

Two ~~one~~ 14-day courses of treatment
May take 1 to 4 days for full effect



*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Location for lot number and expiration date

401D6 FA C6-D 100318

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

Location for lot number and expiration date

40155 FA C6 100318

- Tips for Managing Heartburn**
- Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking



<p>Drug Facts</p> <p>Active ingredient (in each tablet).....Omeprazole 20 mg.....Acid reducer</p> <p>Purpose</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p> <p>Drug Facts (continued)</p>	<p>Use</p> <ul style="list-style-type: none"> Treats frequent heartburn (occurs 2 or more days a week) not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect <p>Warnings</p> <p>Allergy alert: Do not use if you are allergic to omeprazole</p> <p>Do not use if you have:</p> <ul style="list-style-type: none"> trouble or pain swallowing food, vomiting with blood, or bloody or black stools heartburn with lightheadedness, sweating or dizziness chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness <p>frequent chest pain</p> <p>These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have:</p> <ul style="list-style-type: none"> had heartburn over 3 months. This may be a sign of a more serious condition. frequent wheezing, particularly with heartburn unexplained weight loss nausea or vomiting stomach pain <p>Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if:</p> <ul style="list-style-type: none"> your heartburn continues or worsens you need to take this product for more than 14 days you need to take more than 1 course of treatment every 4 months you get diarrhea you develop a rash or joint pain
<p>Directions</p> <ul style="list-style-type: none"> for adults 18 years of age and older this product is to be used once a day (every 24 hours), every day for 14 days it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours <p>14-Day Course of Treatment</p> <ul style="list-style-type: none"> swallow 1 tablet with a glass of water before eating in the morning take every day for 14 days do not take more than 1 tablet a day do not use for more than 14 days unless directed by your doctor swallow whole. Do not chew, crush, or suck tablets. <p>Repeated 14-Day Courses (if needed)</p> <ul style="list-style-type: none"> you may repeat a 14-day course every 4 months do not take for more than 14 days or more often than every 4 months unless directed by a doctor <p>children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.</p> <p>Other information</p> <ul style="list-style-type: none"> read the directions and warnings before use keep the carton. It contains important information. store at 20-25°C (68-77°F) and protect from moisture <p>Inactive ingredients benzyl alcohol, camille, carnuba wax, FD&C blue #2/indigo carmine aluminum lake, flavor, hypromellose, hypromellose acetate succinate, lactose monohydrate, mentho, modified starch, monoethanolamine, polyethylene glycol 3350, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, titanium dioxide, triacetin, triethyl citrate</p> <p>Questions or comments? 1-800-719-9260</p>	<p>Directions</p> <ul style="list-style-type: none"> for adults 18 years of age and older this product is to be used once a day (every 24 hours), every day for 14 days it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours <p>14-Day Course of Treatment</p> <ul style="list-style-type: none"> swallow 1 tablet with a glass of water before eating in the morning take every day for 14 days do not take more than 1 tablet a day do not use for more than 14 days unless directed by your doctor swallow whole. Do not chew, crush, or suck tablets. <p>Repeated 14-Day Courses (if needed)</p> <ul style="list-style-type: none"> you may repeat a 14-day course every 4 months do not take for more than 14 days or more often than every 4 months unless directed by a doctor <p>children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.</p> <p>Other information</p> <ul style="list-style-type: none"> read the directions and warnings before use keep the carton. It contains important information. store at 20-25°C (68-77°F) and protect from moisture <p>Inactive ingredients benzyl alcohol, camille, carnuba wax, FD&C blue #2/indigo carmine aluminum lake, flavor, hypromellose, hypromellose acetate succinate, lactose monohydrate, mentho, modified starch, monoethanolamine, polyethylene glycol 3350, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, titanium dioxide, triacetin, triethyl citrate</p> <p>Questions or comments? 1-800-719-9260</p>

Omeprazole

Delayed Release Tablets 20 mg

Made in Israel
 Manufactured by:
 Dexcel Pharma Technologies Ltd
 10 Hakdima Street, Yokneam 2069200, Israel
 *This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.



Omeprazole

Delayed Release Tablets 20 mg

Safety Feature - Do not use if printed tablet blister unit is open or torn.



NDC 0000-000-00

Compare to Prilosec OTC®*

Treats **FREQUENT** Heartburn!



Omeprazole

Delayed Release Tablets 20 mg

Acid Reducer

SWALLOW - DO NOT CHEW

42 TABLETS

Three 14-day courses of treatment
 May take 1 to 4 days for full effect




WILDBERRY MINT
 Coated Tablet

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box

0000000000



0000000000

Drug Facts

Active ingredient (in each tablet)

Omeprazole 20 mg.....Acid reducer

Purpose

Acid reducer

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
 - heartburn with **lightheadedness, sweating or dizziness**
 - chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
 - frequent **chest pain**
- These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days

Drug Facts (continued)

- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- **swallow whole. Do not chew, crush, or suck tablets.**

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**

■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

Drug Facts (continued)

Inactive ingredients

benzyl alcohol, carmine, carnauba wax, FD&C blue #2/indigo carmine aluminum lake, flavor, hypromellose, hypromellose acetate succinate, lactose monohydrate, menthol, modified starch, monoethanolamine, polyethylene glycol 3350, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

1-800-719-9260

3 BOTTLES INSIDE

Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

Safety Feature - Do not use if printed seal under cap is broken or missing.

NDC 0000-000-00

Compare to Prilosec OTC®*



Treats **FREQUENT** Heartburn!



Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

SWALLOW - DO NOT CHEW


42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect



WILDBERRY MINT
Coated Tablet

- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking



0000000000

0000000000



*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

401D7 FA C6 092718

Location for lot number and expiration date

STANDARD FORMAT

TYPE SIZES:

TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued) 8 pt Helvetica Neue LT Std 57 Condensed

HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed

TEXT: 6 pt Helvetica Neue LT Std 57 Condensed

LEADING: 6.5 pt

BULLETS: 5 pt Square with 2 M'S spacing between statements

TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed

HEAVY LINES: 1.5 pt

HAIRLINES: 0.5 pt

Hairlines extend to within two spaces of the "Drug Facts" box

000000000

Drug Facts

Active ingredient (in each tablet) Omeprazole 20 mg.....**Purpose** Acid reducer

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days

Drug Facts (continued)

- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- **swallow whole. Do not chew, crush, or suck tablets.**

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**

- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

Drug Facts (continued)

Inactive ingredients

benzyl alcohol, carmine, carnauba wax, FD&C blue #2/indigo carmine aluminum lake, flavor, hypromellose, hypromellose acetate succinate, lactose monohydrate, menthol, modified starch, monoethanolamine, polyethylene glycol 3350, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?
1-800-719-9260

3 BOTTLES INSIDE

Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

Safety Feature - Do not use if printed seal under cap is broken or missing.

NDC 0000-000-00

Compare to Prilosec OTC[®]*
FDA Approved

DIE CUT WINDOW

DIE CUT WINDOW

DIE CUT WINDOW

Treats **FREQUENT** Heartburn!

Omeprazole

Delayed Release Tablets 20 mg
Acid Reducer
SWALLOW - DO NOT CHEW

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect

WILDBERRY
MINT
Coated Tablet

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC[®].

000000000

000000000

000000000

STANDARD FORMAT

TYPE SIZES:

TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

(continued) 8 pt Helvetica Neue LT Std 57 Condensed

HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed

TEXT: 6 pt Helvetica Neue LT Std 57 Condensed

LEADING: 6.5 pt

BULLETS: 5 pt Square with 2 M'S spacing between statements

TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed

HEAVY LINES: 1.5 pt

HAIRLINES: 0.5 pt

Hairlines extend to within two spaces of the "Drug Facts" box

Location for lot number and expiration date

Location for lot number and expiration date

- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking



<p>Drug Facts</p> <p>Active ingredient (in each tablet).....Omeprazole 20 mg</p> <p>Purpose.....Acid reducer</p>	<p>Use</p> <ul style="list-style-type: none"> ■ treats frequent heartburn (occurs <i>2 or more</i> days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect 	<p>Warnings</p> <p>Do not use if you have:</p> <ul style="list-style-type: none"> ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness ■ frequent chest pain <p>These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have:</p> <ul style="list-style-type: none"> ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain <p>Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if:</p> <ul style="list-style-type: none"> ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain
<p>Drug Facts (continued)</p> <p>Directions</p> <ul style="list-style-type: none"> ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours <p>14-Day Course of Treatment</p> <ul style="list-style-type: none"> ■ swallow 1 tablet with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor. <p>Swallow whole. Do not chew, crush, or suck tablets.</p> <p>Repeated 14-Day Courses (if needed)</p> <ul style="list-style-type: none"> ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor <p>Other information</p> <ul style="list-style-type: none"> ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F) and protect from moisture <p>Inactive ingredients benzyl alcohol, carmine, carnuba wax, FD&C blue #2/indigo carmine aluminum lake, flavor, hypromellose acetate succinate, lactose monohydrate, mannitol, modified starch, starch glycolate, sodium stearate, sodium lauryl sulfate, sodium monohydroxide, sodium stearyl fumarate, sucralose, talc, titanium dioxide, triacetin, triethyl citrate</p> <p>Questions or comments? 1-800-719-9260</p>	<p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p>	

40155 FA C6-B 100318

Omeprazole
Delayed Release Tablets 20 mg

Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakdima Street, Yokneam 2069200, Israel

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.



Omeprazole
Delayed Release Tablets 20 mg

Safety Feature - Do not use if printed tablet blister unit is open or torn.



NDC 0000-000-00

Compare to Prilosec OTC®*
14 FREE

Treats **FREQUENT** Heartburn! **24 HR**

Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

SWALLOW - DO NOT CHEW

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect



WILDBERRY MINT
Coated Tablet

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued)
8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

Drug Facts

Active ingredient (in each tablet)

Omeprazole 20 mg.....Acid reducer

Purpose

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
 - heartburn with **lightheadedness, sweating or dizziness**
 - chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
 - frequent **chest pain**
- These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days

Drug Facts (continued)

- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- **swallow whole. Do not chew, crush, or suck tablets.**

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

Drug Facts (continued)

Inactive ingredients

benzyl alcohol, carmine, carnauba wax, FD&C blue #2/indigo carmine aluminum lake, flavor, hypromellose, hypromellose acetate succinate, lactose monohydrate, menthol, modified starch, monoethanolamine, polyethylene glycol 3350, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

1-800-719-9260

3 BOTTLES INSIDE

Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

Safety Feature - Do not use if printed seal under cap is broken or missing.

NDC 0000-000-00

Compare to Prilosec OTC®*

14 FREE

DIE CUT WINDOW

DIE CUT WINDOW

DIE CUT WINDOW

Treats **FREQUENT** Heartburn! **24 HR**

Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

SWALLOW - DO NOT CHEW

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect



WILDBERRY MINT
Coated Tablet

- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking



*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Location for lot number and expiration date

401D7 FA C6-B 092718


STANDARD FORMAT

- TYPE SIZES:**
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

Location for lot number and expiration date

40155 FA C6-C 100318

- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking



Drug Facts

Active ingredient (in each tablet) Omeprazole 20 mg

Purpose Acid reducer

Use
 ■ treats frequent heartburn (occurs 2 or more days a week)
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:
 ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
 ■ heartburn with lightheadedness, sweating or dizziness
 ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
 ■ frequent chest pain
 These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss
 ■ nausea or vomiting
 ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea
 ■ you develop a rash or joint pain

Questions or comments? 1-800-719-9260

Inactive ingredients benzyl alcohol, camphor, carnauba wax, FD&C blue #2/indigo carmine aluminum lake, flavor, hypromellose, hypromellose acetate succinate, lactose monohydrate, menthyl, modified starch, starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, titanium dioxide, triacetin, triethyl citrate

Other information
 ■ read the directions and warnings before use
 ■ keep the carton. It contains important information.
 ■ store at 20-25°C (68-77°F) and protect from moisture

Directions
 ■ for adults 18 years of age and older
 ■ this product is to be used once a day (every 24 hours), every day for 14 days
 ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
 ■ swallow 1 tablet with a glass of water before eating in the morning
 ■ take every day for 14 days
 ■ do not take more than 1 tablet a day

Repeated 14-Day Courses (if needed)
 ■ do not use for more than 14 days unless directed by your doctor
 ■ swallow whole. Do not chew, crush, or suck tablets.

4 months unless directed by a doctor
 ■ you may repeat a 14-day course every 4 months
 ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Drug Facts (continued)
 If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Omeprazole

Delayed Release Tablets 20 mg

Made in Israel
 Manufactured by:
Dexcel Pharma Technologies Ltd
 10 Hakikima Street, Yokneam 2069200, Israel
 *This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.



Omeprazole

Delayed Release Tablets 20 mg

Safety Feature - Do not use if printed tablet blister unit is open or torn.

NDC 0000-000-00

Compare to Prilosec OTC®*
28 + 14 FREE

Treats **FREQUENT** Heartburn!



Omeprazole

Delayed Release Tablets 20 mg

Acid Reducer

SWALLOW - DO NOT CHEW

42 TABLETS

Three 14-day courses of treatment
 May take 1 to 4 days for full effect



WILDBERRY MINT
 Coated Tablet

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box

Drug Facts

Active ingredient (in each tablet)

Omeprazole 20 mg.....Acid reducer

Purpose

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days

Drug Facts (continued)

- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- **swallow whole. Do not chew, crush, or suck tablets.**

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**

■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

Drug Facts (continued)

Inactive ingredients

benzyl alcohol, carmine, carnauba wax, FD&C blue #2/indigo carmine aluminum lake, flavor, hypromellose, hypromellose acetate succinate, lactose monohydrate, menthol, modified starch, monoethanolamine, polyethylene glycol 3350, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

1-800-719-9260

3 BOTTLES INSIDE

Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

Safety Feature - Do not use if printed seal under cap is broken or missing.

NDC 0000-000-00

Compare to Prilosec OTC®*

28 + 14 FREE

DIE CUT WINDOW

DIE CUT WINDOW

DIE CUT WINDOW

Treats **FREQUENT** Heartburn!



Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

SWALLOW - DO NOT CHEW

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect



WILDBERRY MINT
Coated Tablet



*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

401D7 FA C6-C 092718

Location for lot number and expiration date

STANDARD FORMAT

TYPE SIZES:

TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box

Location for lot number and expiration date

40155 FA C6-E 100318

- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking



Drug Facts

Active ingredient (in each tablet)
Omeprazole 20 mg.....Acid reducer

Purpose
Acid reducer

Use
■ treats frequent heartburn (occurs **2 or more** days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to omeprazole
Do not use if you have:
■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
■ heartburn with **lightheadedness, sweating or dizziness**
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
■ frequent **chest pain**
These may be signs of a serious condition. See your doctor.
Ask a doctor before use if you have:
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss
■ nausea or vomiting
■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:
■ your heartburn continues or worsens
■ you need to take this product for more than 14 days
■ you get diarrhea
■ you develop a rash or joint pain

Inactive ingredients
blue #2/indigo carmine aluminum lake, flavor, hypromellose, hypromellose acetate succinate, lactose monohydrate, menthol, modified starch, monoethanolamine, polyethylene glycol 3350, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, titanium dioxide, triacetin, triethyl citrate

Other information
■ read the directions and warnings before use
■ keep the carton. It contains important information.
■ store at 20-25°C (68-77°F) and protect from moisture

Directions
■ for adults 18 years of age and older
■ this product is to be used once a day (every 24 hours), every day for 14 days
■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
■ swallow 1 tablet with a glass of water before eating in the morning
■ take every day for 14 days
■ do not take more than 1 tablet a day
■ do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)
■ swallow **whole. Do not chew, crush, or suck tablets.**
■ you may repeat a 14-day course every 4 months
■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Questions or comments? 1-800-719-9260

Omeprazole

Delayed Release Tablets 20 mg

Made in Israel
Manufactured by:
Cesca Pharma Technologies Ltd
10 Hakdima Street, Yokneam 2069200, Israel
*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.



Omeprazole

Delayed Release Tablets 20 mg

Safety Feature - Do not use if printed tablet blister unit is open or torn.

NDC 0000-000-00

Compare to Prilosec OTC®*
Bonus! 14 Free

Treats **FREQUENT** Heartburn! **24 HR**

Omeprazole

Delayed Release Tablets 20 mg

Acid Reducer

SWALLOW - DO NOT CHEW

42 ~~28~~ TABLETS

Three ~~Two~~ 14-day courses of treatment
May take 1 to 4 days for full effect



WILDBERRY MINT
Coated Tablet

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

Drug Facts

Active ingredient (in each tablet)

Omeprazole 20 mg.....Acid reducer

Purpose

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days

Drug Facts (continued)

- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- **swallow whole. Do not chew, crush, or suck tablets.**

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

Drug Facts (continued)

Inactive ingredients

benzyl alcohol, carmine, carnauba wax, FD&C blue #2/indigo carmine aluminum lake, flavor, hypromellose, hypromellose acetate succinate, lactose monohydrate, menthol, modified starch, monoethanolamine, polyethylene glycol 3350, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

1-800-719-9260

3 BOTTLES INSIDE

Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

Safety Feature - Do not use if printed seal under cap is broken or missing.

NDC 0000-000-00

Compare to Prilosec OTC®*
Bonus! 14 Free

DIE CUT WINDOW

DIE CUT WINDOW

DIE CUT WINDOW

Treats **FREQUENT** Heartburn!



Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

SWALLOW - DO NOT CHEW

42 ~~28~~ TABLETS

Three ~~two~~ 14-day courses of treatment
May take 1 to 4 days for full effect



WILDBERRY MINT
Coated Tablet



*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

401D7 FA C6-E 100318

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Location for lot number and expiration date


STANDARD FORMAT

TYPE SIZES:

- TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique**
- TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued) 8 pt Helvetica Neue LT Std 57 Condensed**
- HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique**
- SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed**
- TEXT: 6 pt Helvetica Neue LT Std 57 Condensed**
- LEADING: 6.5 pt**
- BULLETS: 5 pt Square with 2 M'S spacing between statements**
- TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed**
- HEAVY LINES: 1.5 pt**
- HAIRLINES: 0.5 pt**
- Hairlines extend to within two spaces of the "Drug Facts" box**

Location for lot number and expiration date

- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Raise the head of your bed
 - Do not eat before bedtime
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking



Drug Facts

Active Ingredient (in each tablet) Omeprazole 20 mg.....acid reducer

Purpose If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Use ■ treats frequent heartburn (occurs *2 or more* days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings **Allergy alert:** Do not use if you are allergic to omeprazole

Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with **lightheadedness, sweating or dizziness** ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness

Ask a doctor before use if you have: ■ frequent chest pain ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain

Inactive ingredients benzyl alcohol, carnitine, carnauba wax, FD&C blue #2/indigo carmine aluminum lake, flavor, hypromellose, hypromellose acetate succinate, lactose monohydrate, menthol, modified starch, monoethanolamine, polyethylene glycol 3350, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, titanium dioxide, tritethyn, triethyl citrate

Other information ■ read the directions and warnings before use ■ keep the carton, it contains important information. ■ store at 20-25°C (68-77°F) and protect from moisture

Directions ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment ■ swallow 1 tablet with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ swallow whole. **Do not chew, crush, or suck tablets.**

Repeated 14-Day Courses (if needed) ■ you may repeat a 14-day course every 4 months ■ **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**

■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Questions or comments? 1-800-719-9260

40155 FA C6-D 100318

Omeprazole
Delayed Release Tablets 20 mg

Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakdama Street, Yokneam 2069200, Israel

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.



Omeprazole

Delayed Release Tablets 20 mg

Safety Feature - Do not use if printed tablet blister unit is open or torn.

NDC 0000-000-00

Compare to Prilosec OTC®*

Bonus Pack! 14 Free

Treats **FREQUENT** Heartburn!

24 HR

Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

SWALLOW - DO NOT CHEW

42 ~~28~~ TABLETS

Three ~~two~~ 14-day courses of treatment
May take 1 to 4 days for full effect

WILDBERRY MINT
Coated Tablet

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

Drug Facts

Active ingredient (in each tablet) Purpose
Omeprazole 20 mg.....Acid reducer

Use
■ treats frequent heartburn (occurs **2 or more** days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to omeprazole
Do not use if you have:
■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
■ heartburn with **lightheadedness, sweating or dizziness**
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
■ frequent **chest pain**
These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss
■ nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:
■ your heartburn continues or worsens
■ you need to take this product for more than 14 days

Drug Facts (continued)

- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
■ for adults 18 years of age and older
■ this product is to be used once a day (every 24 hours), every day for 14 days
■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
■ swallow 1 tablet with a glass of water before eating in the morning
■ take every day for 14 days
■ do not take more than 1 tablet a day
■ do not use for more than 14 days unless directed by your doctor
■ **swallow whole. Do not chew, crush, or suck tablets.**

Repeated 14-Day Courses (if needed)
■ you may repeat a 14-day course every 4 months
■ **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information
■ read the directions and warnings before use
■ keep the carton. It contains important information.
■ store at 20-25°C (68-77°F) and protect from moisture

Drug Facts (continued)

Inactive ingredients
benzyl alcohol, carmine, carnauba wax, FD&C blue #2/indigo carmine aluminum lake, flavor, hypromellose, hypromellose acetate succinate, lactose monohydrate, menthol, modified starch, monoethanolamine, polyethylene glycol 3350, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?
1-800-719-9260

3 BOTTLES INSIDE
Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

Safety Feature - Do not use if printed seal under cap is broken or missing.

NDC 0000-000-00

Compare to Prilosec OTC®*

Bonus Pack! 14 Free

DIE CUT WINDOW

DIE CUT WINDOW

DIE CUT WINDOW

Treats **FREQUENT** Heartburn!



Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

SWALLOW - DO NOT CHEW

42 ~~28~~ TABLETS

Three ~~two~~ 14-day courses of treatment
May take 1 to 4 days for full effect



WILDBERRY MINT
Coated Tablet



*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

401D7 FA C6-D 100318

Location for lot number and expiration date

STANDARD FORMAT

- TYPE SIZES:**
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

Location for lot number and expiration date

40155 FA C6-G 100318

- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

<p>Drug Facts</p> <p>Active Ingredient (in each tablet) Omeprazole 20 mg Acid reducer</p> <p>Purpose Acid reducer</p>	<p>Use</p> <ul style="list-style-type: none"> treats frequent heartburn (occurs <i>2 or more</i> days a week) not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect 	<p>Warnings</p> <p>Do not use if you have:</p> <ul style="list-style-type: none"> trouble or pain swallowing food, vomiting with blood, or bloody or black stools heartburn with lightheadedness, sweating or dizziness chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness frequent chest pain <p>These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have:</p> <ul style="list-style-type: none"> had heartburn over 3 months. This may be a sign of a more serious condition. frequent wheezing, particularly with heartburn unexplained weight loss nausea or vomiting stomach pain <p>Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p>	<p>Do not use if you have:</p> <ul style="list-style-type: none"> allergy alert: Do not use if you are allergic to omeprazole
<p>Drug Facts (continued)</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p> <p>Directions</p> <ul style="list-style-type: none"> for adults 18 years of age and older this product is to be used once a day (every 24 hours), every day for 14 days it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours <p>14-Day Course of Treatment</p> <ul style="list-style-type: none"> swallow 1 tablet with a glass of water before eating in the morning take every day for 14 days do not take more than 1 tablet a day do not use for more than 14 days unless directed by your doctor swallow whole. Do not chew, crush, or suck tablets. <p>Repeated 14-Day Courses (if needed)</p> <ul style="list-style-type: none"> you may repeat a 14-day course every 4 months do not take for more than 14 days or more often than every 4 months unless directed by a doctor <p>children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.</p>	<p>Other Information</p> <ul style="list-style-type: none"> read the directions and warnings before use keep the carton. It contains important information. store at 20-25°C (68-77°F) and protect from moisture 	<p>Inactive Ingredients benzyl alcohol, camphor, carnauba wax, FD&C blue #2, indigo carmine, aluminum lake, flavor, hypromellose, hypromellose acetate succinate, lactose monohydrate, menthol, modified starch, monohydroxyethylamine, polyethylene glycol 3350, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, titanium dioxide, triacetin, triethyl citrate</p>	<p>Questions or comments? 1-800-719-9260</p>

Omeprazole

Delayed Release Tablets 20 mg

Made in Israel
Manufactured by:
Ceskel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel
*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.



Omeprazole

Delayed Release Tablets 20 mg

Safety Feature - Do not use if printed tablet blister unit is open or torn.

NDC 0000-000-00

Compare to Prilosec OTC®*

Triple Pack
Three 14-day courses of treatment

Treats **FREQUENT** Heartburn!



Omeprazole

Delayed Release Tablets 20 mg

Acid Reducer

SWALLOW - DO NOT CHEW

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect



WILDBERRY MINT
Coated Tablet

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued): 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

000000000

000000000

Drug Facts

Active ingredient (in each tablet) Purpose
Omeprazole 20 mg.....Acid reducer

Use
■ treats frequent heartburn (occurs **2 or more** days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to omeprazole
Do not use if you have:
■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
■ heartburn with **lightheadedness, sweating or dizziness**
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
■ frequent **chest pain**
These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss
■ nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:
■ your heartburn continues or worsens
■ you need to take this product for more than 14 days

Drug Facts (continued)

■ you need to take more than 1 course of treatment every 4 months
■ you get diarrhea
■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
■ for adults 18 years of age and older
■ this product is to be used once a day (every 24 hours), every day for 14 days
■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
■ swallow 1 tablet with a glass of water before eating in the morning
■ take every day for 14 days
■ do not take more than 1 tablet a day
■ do not use for more than 14 days unless directed by your doctor
■ **swallow whole. Do not chew, crush, or suck tablets.**

Repeated 14-Day Courses (if needed)
■ you may repeat a 14-day course every 4 months
■ **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information
■ read the directions and warnings before use
■ keep the carton. It contains important information.
■ store at 20-25°C (68-77°F) and protect from moisture

Drug Facts (continued)

Inactive ingredients
benzyl alcohol, carmine, carnauba wax, FD&C blue #2/indigo carmine aluminum lake, flavor, hypromellose, hypromellose acetate succinate, lactose monohydrate, menthol, modified starch, monoethanolamine, polyethylene glycol 3350, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?
1-800-719-9260

3 BOTTLES INSIDE
Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

Safety Feature - Do not use if printed seal under cap is broken or missing.

NDC 0000-000-00 Compare to Prilosec OTC®*

Triple Pack Three 14-day courses of treatment

DIE CUT WINDOW

DIE CUT WINDOW

DIE CUT WINDOW

Treats **FREQUENT** Heartburn! **24 HR**

Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer
SWALLOW - DO NOT CHEW



WILDBERRY MINT
Coated Tablet

42 TABLETS
Three 14-day courses of treatment
May take 1 to 4 days for full effect



*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

401D7 FA C6-G 092718

Location for lot number and expiration date

STANDARD FORMAT
TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

NDC 0000-000-00

Compare to Prilosec OTC®*

3 PACK

Three 14-day courses of treatment

Treats **FREQUENT** Heartburn!



Omeprazole

Delayed Release Tablets 20 mg

Acid Reducer

SWALLOW - DO NOT CHEW

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect



WILDBERRY MINT
Coated Tablet

40165 FA C6-A 092718


Made in Israel
Manufactured by:
DeKael Pharma Technologies Ltd
10 Hakoma Street, Yokneam 206200, Israel

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Do not eat before bed
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Drug Facts Active ingredient (in each tablet) Omeprazole 20 mg Purpose Acid reducer	Use ■ treats frequent heartburn (occurs <i>2 or more</i> days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect Warnings ■ Allergy alert Do not use if you are allergic to omeprazole Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness ■ frequent chest pain These may be signs of a serious condition. See your doctor. Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs. Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you develop a rash or joint pain	Directions ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours 14-Day Course of Treatment ■ swallow 1 tablet with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ swallow whole. Do not chew, crush, or suck tablets. Repeated 14-Day Courses (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition. Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ avoid heat and moisture ■ store at 20°-25°C (68°-77°F) and protect from moisture. Inactive ingredients FD&C blue #2/indigo carmine aluminum lake, ferrocyanine, carmine, camouba wax, hypromellose acetate succinate, lactose monohydrate, mannitol, modified starch, monobenzene, polyethylene glycol 3350, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, titanium dioxide, triacetin, triethyl citrate	Questions or comments? 1-800-719-9280
--	---	--	--

Safety Feature -
Do not use if
printed tablet
batter unit is
open or torn.



Location for lot number and expiration date

PRINT at 65% for 11-inch x 17-inch size paper

STANDARD FORMAT

TYPE SIZES:
TITLE: 10 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued): 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 9-7 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 2.0 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

00000000



Drug Facts

Active ingredient (in each tablet) Omeprazole 20 mg.....**Purpose** Acid reducer

Use
■ treats frequent heartburn (occurs **2 or more** days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to omeprazole
Do not use if you have:
■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
■ heartburn with **lightheadedness, sweating or dizziness**
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
■ frequent **chest pain**
These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss
■ nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:
■ your heartburn continues or worsens
■ you need to take this product for more than 14 days

Drug Facts (continued)

■ you need to take more than 1 course of treatment every 4 months
■ you get diarrhea
■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
■ for adults 18 years of age and older
■ this product is to be used once a day (every 24 hours), every day for 14 days
■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
■ swallow 1 tablet with a glass of water before eating in the morning
■ take every day for 14 days
■ do not take more than 1 tablet a day
■ do not use for more than 14 days unless directed by your doctor
■ **swallow whole. Do not chew, crush, or suck tablets.**

Repeated 14-Day Courses (if needed)
■ you may repeat a 14-day course every 4 months
■ **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information
■ read the directions and warnings before use
■ keep the carton. It contains important information.
■ store at 20-25°C (68-77°F) and protect from moisture

Drug Facts (continued)

Inactive ingredients
benzyl alcohol, carmine, carnauba wax, FD&C blue #2/indigo carmine aluminum lake, flavor, hypromellose, hypromellose acetate succinate, lactose monohydrate, menthol, modified starch, monoethanolamine, polyethylene glycol 3350, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?
1-800-719-9260

3 BOTTLES INSIDE
Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel



00000000

Safety Feature - Do not use if printed seal under cap is broken or missing.



NDC 0000-000-00

Compare to Prilosec OTC®*

3 PACK Three 14-day courses of treatment



Treats **FREQUENT** Heartburn!

Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

SWALLOW - DO NOT CHEW

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect



WILDBERRY MINT
Coated Tablet

Tips for Managing Heartburn
■ Do not lie flat or bend over after eating
■ Do not wear tight-fitting clothing around the stomach
■ Do not eat before bedtime
■ Raise the head of your bed
■ Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
■ Eat slowly and avoid big meals
■ If overweight, lose weight
■ Quit smoking



*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Location for lot number and expiration date

401D7 FA C6-A 092718

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

NDC 0000-000-00

Treats **FREQUENT** Heartburn!



Compare to Prilosec OTC®*
Value Pack!
3-14 Tablet Cartons

Omeprazole

Delayed Release Tablets 20 mg

Acid Reducer

SWALLOW - DO NOT CHEW

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect



WILDBERRY MINT
Coated Tablet

40165 FA C6-J 092718


Made in Israel
Manufactured by:
DeKael Pharma Technologies Ltd
10 Hakoma Street, Yokneam 206200, Israel

Tips for Managing Heartburn

- Do not eat before bedtime
- Do not wear tight-fitting clothing around the stomach
- Do not lie flat or bend over after eating
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- Raise the head of your bed
- Quit smoking
- If overweight, lose weight

Drug Facts Active ingredient (in each tablet) Omeprazole 20 mg Purpose Acid reducer	Use ■ treats frequent heartburn (occurs <i>2 or more</i> days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect Warnings ■ Do not use if you are allergic to omeprazole Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with tightness or soreness of throat, sweating, pain spreading to arms, neck or shoulders, or lightheadedness ■ frequent chest pain These may be signs of a serious condition. See your doctor. Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs. Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you develop a rash or joint pain	Directions ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours 14-Day Course of Treatment ■ swallow 1 tablet with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ swallow whole. Do not chew, crush, or suck tablets. Repeated 14-Day Courses (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition. Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ Ask a doctor or pharmacist before use if you are taking a prescription drug. Inactive ingredients benzyl alcohol, carmine, carnauba wax, FD&C blue #2/indigo carmine aluminum lake, ferrocyanine, hypromellose acetate succinate, lactose monohydrate, mannitol, modified starch, monobenzene, polyethylene glycol 3350, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, titanium dioxide, triacetin, triethyl citrate
Drug Facts (continued) Questions or comments? 1-800-719-9280		

Safety Feature -
Do not use if
printed tablet
blister unit is
open or torn.



Location for lot number and expiration date

PRINT at 65% for 11-inch x 17-inch size paper

STANDARD FORMAT
TYPE SIZES:
TITLE: 10 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued): 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 9-7 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 2.0 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

Drug Facts

Active ingredient (in each tablet)

Omeprazole 20 mg.....Acid reducer

Purpose

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
 - heartburn with **lightheadedness, sweating or dizziness**
 - chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
 - frequent **chest pain**
- These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days

Drug Facts (continued)

- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- **swallow whole. Do not chew, crush, or suck tablets.**

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**

■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

Drug Facts (continued)

Inactive ingredients

benzyl alcohol, carmine, carnauba wax, FD&C blue #2/indigo carmine aluminum lake, flavor, hypromellose, hypromellose acetate succinate, lactose monohydrate, menthol, modified starch, monoethanolamine, polyethylene glycol 3350, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, sucralose, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

1-800-719-9260

3 BOTTLES INSIDE

Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

Safety Feature - Do not use if printed seal under cap is broken or missing.

NDC 0000-000-00

Compare to Prilosec OTC®*

Value Pack! 3-14 Tablet Bottles

DIE CUT WINDOW

DIE CUT WINDOW

DIE CUT WINDOW

Treats **FREQUENT** Heartburn!



Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

SWALLOW - DO NOT CHEW

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect



WILDBERRY MINT
Coated Tablet



*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

401D7 FA C6-J 092718

Location for lot number and expiration date

STANDARD FORMAT

TYPE SIZES:

TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed

HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed

TEXT: 6 pt Helvetica Neue LT Std 57 Condensed

LEADING: 6.5 pt

BULLETS: 5 pt Square with 2 M'S spacing between statements

TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed

HEAVY LINES: 1.5 pt

HAIRLINES: 0.5 pt

Hairlines extend to within two spaces of the "Drug Facts" box

FRONT PANEL

peel off coupon

Instant Redeemable Coupon - Expires XX-XX-XX

SAVE \$X NOW!

On This Purchase

Compare to Prilosec OTC®*

NDC 0000-000-00

Treats **FREQUENT** Heartburn!



Omeprazole

Delayed Release Tablets 20 mg

Acid Reducer

14 TABLETS

One 14-day course of treatment
May take 1 to 4 days for full effect



91574 FA CP6 092418

BACK PANEL

SAVE \$X NOW!

MANUFACTURER'S COUPON EXPIRATION DATE XX/XX/XXXX

SAVE \$X^{XX}

on your purchase of
14 COUNT SIZE

"Store Brand" Omeprazole
Delayed Release Tablets 20 mg

CONSUMER: Only one coupon is redeemable per purchase. **RETAILER:** This coupon will be redeemable for face value on specified products (plus a handling fee) if submitted in compliance with the terms of this offer. Cash value 1/100th of one cent. Limit one coupon per customer. **Do Not Double.** Coupon is non-reproducible. Coupons cannot be combined. Mail properly handled coupons to: L. Perrigo Company (#1565), NCH Marketing, PO Box 880001, El Paso, TX 88588-0001

COUPON NOT SUBJECT TO DOUBLING

Customer pays any sales tax.



Omeprazole

Delayed Release Tablets 20 mg

14 TABLETS

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Reference ID: 4415974

FRONT PANEL

peel off coupon

Instant Redeemable Coupon - Expires XX-XX-XX

SAVE \$X NOW!
On This Purchase

Compare to
Prilosec OTC®*

NDC 0000-000-00

Treats **FREQUENT** Heartburn!



24 HR

Omeprazole

Delayed Release Tablets 20 mg
Acid Reducer

14 TABLETS



One 14-day course of treatment

Reference ID: 4415974

915D5 FA CP6 092418

BACK PANEL

SAVE \$X NOW!

Omeprazole

Delayed Release
Tablets 20 mg

14 TABLETS

*This product is not
manufactured or distributed by
Procter & Gamble, distributor
of Prilosec OTC®.

MANUFACTURER'S COUPON EXPIRATION DATE XX/XX/XXXX

SAVE \$X^{XX}

on your purchase of
14 COUNT SIZE

"Store Brand" Omeprazole
Delayed Release Tablets 20 mg

**COUPON NOT SUBJECT
TO DOUBLING**

Customer pays any sales tax.

CONSUMER: Only one coupon is redeemable per purchase. **RETAILER:** This coupon will be redeemable for face value on specified products (plus a handling fee) if submitted in compliance with the terms of this offer. Cash value 1/100th of one cent. Limit one coupon per customer. **Do Not Double.** Coupon is non-reproducible. Coupons cannot be combined. Mail properly handled coupons to: L. Perrigo Company (#1565), NCH Marketing, PO Box 880001, El Paso, TX 88588-0001



FRONT PANEL

peel off coupon

Instant Redeemable Coupon - Expires XX-XX-XX

SAVE \$X NOW!

On This Purchase

Compare to Prilosec OTC®*

NDC 0000-000-00

Treats **FREQUENT** Heartburn!



Omeprazole

Delayed Release Tablets 20 mg Acid Reducer

28 TABLETS

Two 14-day courses of treatment
May take 1 to 4 days for full effect



91530 FA CP6 092418

BACK PANEL

SAVE \$X NOW!

MANUFACTURER'S COUPON EXPIRATION DATE XX/XX/XXXX

SAVE \$X^{XX}

on your purchase of
28 COUNT SIZE
"Store Brand" Omeprazole
Delayed Release Tablets 20 mg

**COUPON NOT SUBJECT
TO DOUBLING**

Customer pays any sales tax.

CONSUMER: Only one coupon is redeemable per purchase. RETAILER: This coupon will be redeemable for face value on specified products (plus a handling fee) if submitted in compliance with the terms of this offer. Cash value 1/100th of one cent. Limit one coupon per customer. Do Not Double. Coupon is non-reproducible. Coupons cannot be combined. Mail properly handled coupons to: L. Perrigo Company (#1565), NCH Marketing, PO Box 880001, El Paso, TX 88588-0001



Omeprazole

Delayed Release Tablets 20 mg

28 TABLETS

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Reference ID: 4415974

FRONT PANEL

peel off coupon

Instant Redeemable Coupon - Expires XX-XX-XX

SAVE \$X NOW!
On This Purchase

Compare to Prilosec OTC®*

NDC 0000-000-00

Treats **FREQUENT** Heartburn!

 **24 HR**

Omeprazole

Delayed Release Tablets 20 mg
Acid Reducer

28 TABLETS

Two 14-day courses of treatment
May take 1 to 4 days for full effect



915D6 FA CP6 092418

BACK PANEL

SAVE \$X NOW!

MANUFACTURER'S COUPON EXPIRATION DATE XX/XX/XXXX

SAVE \$X^{XX}

on your purchase of
28 COUNT SIZE
"Store Brand" Omeprazole

Delayed Release Tablets 20 mg

**COUPON NOT SUBJECT
TO DOUBLING**

Customer pays any sales tax.

Omeprazole

Delayed Release Tablets 20 mg

28 TABLETS

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

CONSUMER: Only one coupon is redeemable per purchase. RETAILER: This coupon will be redeemable for face value on specified products (plus a handling fee) if submitted in compliance with the terms of this offer. Cash value 1/100th of one cent. Limit one coupon per customer. Do Not Double. Coupon is non-reproducible. Coupons cannot be combined. Mail properly handled coupons to: L. Perrigo Company (#1565), NCH Marketing, PO Box 880001, El Paso, TX 88588-0001



Reference-ID: 4415974

FRONT PANEL

peel off coupon

Instant Redeemable Coupon - Expires XX-XX-XX

SAVE \$X NOW!

On This Purchase

Compare to Prilosec OTC®*

NDC 0000-000-00

Treats **FREQUENT** Heartburn!

 **24 HR**

Omeprazole

Delayed Release Tablets 20 mg

Acid Reducer

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect



91555 FA CP6 092418

BACK PANEL

SAVE \$X NOW!

MANUFACTURER'S COUPON EXPIRATION DATE XX/XX/XXXX

SAVE \$X^{XX}

on your purchase of
42 COUNT SIZE
"Store Brand" Omeprazole
Delayed Release Tablets 20 mg

CONSUMER: Only one coupon is redeemable per purchase. **RETAILER:** This coupon will be redeemable for face value on specified products (plus a handling fee) if submitted in compliance with the terms of this offer. Cash value 1/100th of one cent. Limit one coupon per customer. **Do Not Double.** Coupon is non-reproducible. Coupons cannot be combined. Mail properly handled coupons to:

L. Perrigo Company (#1565),
NCH Marketing, PO Box 880001,
El Paso, TX 88588-0001

COUPON NOT SUBJECT TO DOUBLING

Customer pays any sales tax.



Omeprazole

Delayed Release Tablets 20 mg

42 TABLETS

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Reference ID: 4415974

FRONT PANEL

Read off coupon

Instant Redeemable Coupon - Expires XX-XX-XX

SAVE \$X NOW!

On This Purchase

Compare to Prilosec OTC®*

NDC 0000-000-00



Treats **FREQUENT** Heartburn!



Omeprazole

Delayed Release Tablets 20 mg
Acid Reducer

42 TABLETS




Three 14-day courses of treatment
May take 1 to 4 days for full effect



915D7 FA CP6 092418

BACK PANEL

SAVE \$X NOW!



MANUFACTURER'S COUPON EXPIRATION DATE XX/XX/XXXX

SAVE \$X^{XX}

on your purchase of
42 COUNT SIZE
"Store Brand" Omeprazole
Delayed Release Tablets 20 mg

**COUPON NOT SUBJECT
TO DOUBLING**
Customer pays any sales tax.

CONSUMER: Only one coupon is redeemable per purchase. **RETAILER:** This coupon will be redeemable for face value on specified products (plus a handling fee) if submitted in compliance with the terms of this offer. Cash value 1/100th of one cent. Limit one coupon per customer. Do Not Double. Coupon is non-reproducible. Coupons cannot be combined. Mail properly handled coupons to: L. Perrigo Company (#1565), NCH Marketing, P.O. Box 680001, El Paso, TX 88568-0001



Reference ID: 4415974

Omeprazole

Delayed Release Tablets 20 mg

42 TABLETS

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

FRONT PANEL

Peel off coupon

Instant Redeemable Coupon - Expires XX-XX-XX

SAVE \$X NOW!

On This Purchase

Compare to Prilosec OTC®*

NDC 0000-000-00

Treats **FREQUENT** Heartburn!

 **24 HR**

Omeprazole

Delayed Release Tablets 20 mg

Acid Reducer

SWALLOW - DO NOT CHEW

14 TABLETS

One 14-day course of treatment
May take 1 to 4 days for full effect



**WILDBERRY
MINT**
Coated Tablet



40174 FA CP6 092418

BACK PANEL

SAVE \$X NOW!

MANUFACTURER'S COUPON EXPIRATION DATE XX/XX/XXXX

SAVE \$X^{XX}

on your purchase of
14 COUNT SIZE
"Store Brand" Omeprazole
Delayed Release Tablets 20 mg

**COUPON NOT SUBJECT
TO DOUBLING**

Customer pays any sales tax.

CONSUMER: Only one coupon is redeemable per purchase. RETAILER: This coupon will be redeemable for face value on specified products (plus a handling fee) if submitted in compliance with the terms of this offer. Cash value 1/100th of one cent. Limit one coupon per customer. Do Not Double. Coupon is non-reproducible. Coupons cannot be combined. Mail properly handled coupons to: L. Perrigo Company (#1565), NCH Marketing, PO Box 880001, El Paso, TX 88588-0001



Omeprazole

Delayed Release Tablets 20 mg

14 TABLETS

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Reference ID: 4415974

FRONT PANEL

peel off coupon

Instant Redeemable Coupon - Expires XX-XX-XX

SAVE \$X NOW!

On This Purchase

Compare to
Prilosec OTC®*

NDC 0000-000-00

Treats **FREQUENT** Heartburn!



Omeprazole

Delayed Release Tablets 20 mg Acid Reducer

SWALLOW - DO NOT CHEW

14 TABLETS



**WILDBERRY
MINT
Coated Tablet**



One 14-day course of treatment

Reference ID: 4415974

401D5 FA CP6 092418

BACK PANEL

SAVE \$X NOW!

Omeprazole

Delayed Release
Tablets 20 mg

14 TABLETS

*This product is not
manufactured or distributed by
Procter & Gamble, distributor
of Prilosec OTC®.

MANUFACTURER'S COUPON EXPIRATION DATE XX/XX/XXXX

SAVE \$X^{XX}

on your purchase of
14 COUNT SIZE

"Store Brand" Omeprazole
Delayed Release Tablets 20 mg

**COUPON NOT SUBJECT
TO DOUBLING**

Customer pays any sales tax.

CONSUMER: Only one coupon is redeemable per purchase. **RETAILER:** This coupon will be redeemable for face value on specified products (plus a handling fee) if submitted in compliance with the terms of this offer. Cash value 1/100th of one cent. Limit one coupon per customer. **Do Not Double.** Coupon is non-reproducible. Coupons cannot be combined. Mail properly handled coupons to: L. Perrigo Company (#1565), NCH Marketing, PO Box 680001, El Paso, TX 88568-0001



FRONT PANEL

peel off coupon

Instant Redeemable Coupon - Expires XX-XX-XX

SAVE \$X NOW!
On This Purchase

Compare to Prilosec OTC®*

NDC 0000-000-00

Treats **FREQUENT** Heartburn!



Omeprazole

Delayed Release Tablets 20 mg

Acid Reducer

SWALLOW - DO NOT CHEW

28 TABLETS

Two 14-day courses of treatment
May take 1 to 4 days for full effect



**WILDBERRY
MINT**
Coated Tablet



40130 FA CP6 092418

BACK PANEL

SAVE \$X NOW!


MANUFACTURER'S COUPON EXPIRATION DATE XX/XX/XXXX

SAVE \$X^{XX} on your purchase of
28 COUNT SIZE
"Store Brand" Omeprazole
Delayed Release Tablets 20 mg

CONSUMER: Only one coupon is redeemable per purchase. **RETAILER:** This coupon will be redeemable for face value on specified products (plus a handling fee) if submitted in compliance with the terms of this offer. Cash value 1/100th of one cent. Limit one coupon per customer. **Do Not Double.** Coupon is non-reproducible. Coupons cannot be combined. Mail properly handled coupons to: L. Perrigo Company (#1565), NCH Marketing, PO Box 880001, El Paso, TX 88588-0001

COUPON NOT SUBJECT TO DOUBLING

Customer pays any sales tax.



Omeprazole

Delayed Release Tablets 20 mg

28 TABLETS

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Reference ID: 4415974

FRONT PANEL

199¢ Off
Coupon

Instant Redeemable Coupon - Expires XX-XX-XX

SAVE \$X NOW!

On This Purchase

Compare to Prilosec OTC®*

NDC 0000-000-00

Treats **FREQUENT** Heartburn!

 **24 HR**

Omeprazole

Delayed Release Tablets 20 mg Acid Reducer

SWALLOW - DO NOT CHEW

28 TABLETS

Two 14-day courses of treatment
May take 1 to 4 days for full effect



**WILDBERRY
MINT**
Coated Tablet



401D6 FA CP6 092418

BACK PANEL

SAVE \$X NOW!

MANUFACTURER'S COUPON EXPIRATION DATE XX/XX/XXXX

SAVE \$X^{XX}

on your purchase of
28 COUNT SIZE

"Store Brand" Omeprazole

Delayed Release Tablets 20 mg

**COUPON NOT SUBJECT
TO DOUBLING**

Customer pays any sales tax.

CONSUMER: Only one coupon is redeemable per purchase. **RETAILER:** This coupon will be redeemable for face value on specified products (plus a handling fee) if submitted in compliance with the terms of this offer. Cash value 1/100th of one cent. Limit one coupon per customer. **Do Not Double.** Coupon is non-reproducible. Coupons cannot be combined. Mail properly handled coupons to: L. Perrigo Company (#1565), NCH Marketing, PO Box 880001, El Paso, TX 88588-0001.



0 00000 00000 0

Reference ID: 4415974

Omeprazole

Delayed Release Tablets 20 mg

28 TABLETS

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

FRONT PANEL

peel off coupon

Instant Redeemable Coupon - Expires XX-XX-XX

SAVE \$X NOW!

On This Purchase

Compare to Prilosec OTC®*

NDC 0000-000-00

Treats **FREQUENT** Heartburn!



Omeprazole

Delayed Release Tablets 20 mg

Acid Reducer

SWALLOW - DO NOT CHEW

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect



WILDBERRY MINT
Coated Tablet



40155 FA CP6 092418

BACK PANEL

SAVE \$X NOW!

MANUFACTURER'S COUPON EXPIRATION DATE XX/XX/XXXX

SAVE \$X^{XX}

on your purchase of
42 COUNT SIZE
"Store Brand" Omeprazole
Delayed Release Tablets 20 mg

CONSUMER: Only one coupon is redeemable per purchase. **RETAILER:** This coupon will be redeemable for face value on specified products (plus a handling fee) if submitted in compliance with the terms of this offer. Cash value 1/100th of one cent. Limit one coupon per customer. **Do Not Double.** Coupon is non-reproducible. Coupons cannot be combined. Mail properly handled coupons to: L. Perrigo Company (#1565), NCH Marketing, PO Box 880001, El Paso, TX 88588-0001

COUPON NOT SUBJECT TO DOUBLING
Customer pays any sales tax.



0 00000 00000 0

Omeprazole

Delayed Release Tablets 20 mg

42 TABLETS

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Reference ID: 4415974

FRONT PANEL

Peel off coupon

Instant Redeemable Coupon - Expires XX-XX-XX

SAVE \$X NOW!
On This Purchase

Compare to Prilosec OTC®*

NDC 0000-000-00

Treats **FREQUENT** Heartburn!

 **24 HR**

Omeprazole
Delayed Release Tablets 20 mg
Acid Reducer

SWALLOW - DO NOT CHEW

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect



WILDBERRY MINT
Coated Tablet



401D7 FA CP6 092418

BACK PANEL

SAVE \$X NOW!

MANUFACTURER'S COUPON EXPIRATION DATE XX/XX/XXXX

SAVE \$X^{XX}


on your purchase of
42 COUNT SIZE

"Store Brand" Omeprazole
Delayed Release Tablets 20 mg

**COUPON NOT SUBJECT
TO DOUBLING**

Customer pays any sales tax.

CONSUMER: Only one coupon is redeemable per purchase. **RETAILER:** This coupon will be redeemable for face value on specified products (plus a handling fee) if submitted in compliance with the terms of this offer. Cash value 1/100th of one cent. Limit one coupon per customer. Do Not Double. Coupon is non-reproducible. Coupons cannot be combined. Mail properly handled coupons to: L. Perrigo Company (#1565), NCH Marketing, PO Box 880001, El Paso, TX 79868-0001



0 00000 00000 0

Reference ID: 4415974

Omeprazole

Delayed Release Tablets 20 mg

42 TABLETS

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

KAREN M MAHONEY
04/08/2019 12:42:31 PM