2 1 Count Packets

Advil®

ALLERGY & CONGESTION RELIEF

Chlorpheniramine Maleate 4 mg Antihistamine
Ibuprofen 200 mg Pain Reliever/Fever Reducer (NSAID)
Phenylephrine HCl 10 mg Nasal Decongestant

Advil® ALLERGY & CONGESTION RELIEF

2 Packets of 1 Coated Tablet

F.P.O.

3 0573 0196 02 4

Do Not Use if pouch is broken or torn.

Advil®

ALLERGY & CONGESTION RELIEF

Chlorpheniramine Maleate 4 mg Antihistamine
Ibuprofen 200 mg Pain Reliever/Fever Reducer (NSAID)
Phenylephrine HCl 10 mg Nasal Decongestant

- ✓ Sneezing
- ✓ Nasal Congestion
- ✓ Sinus Pressure
- ✓ Nasal Swelling
- ✓ Runny Nose
- ✓ Headache
- ✓ Fever
- ✓ Body Aches



ONE TABLET DOSE!

2 Packets of 1 Coated Tablet

Advil® ALLERGY & CONGESTION RELIEF

For most recent product information, visit **www.advil.com**

Questions or comments?
Call weekdays from **9 AM to 5 PM EST** toll free at **1-800-88-ADVIL**

Pfizer, Madison, NJ 07940 USA © 2018 Pfizer Inc.

PAA110481.FDA01

Reference ID: 4381479

PAA110481.FDA01

2 1 Count Packets

Advil®

MULTI-SYMPTOM COLD & FLU

Ibuprofen 200 mg Pain Reliever/Fever Reducer **(NSAID)**
Phenylephrine HCl 10 mg Nasal Decongestant
Chlorpheniramine Maleate 4 mg Antihistamine

Advil® MULTI-SYMPTOM COLD & FLU

2 Packets of 1 Coated Tablet

F.P.O.

3 0573 0190 02 2

Do Not Use if pouch is broken or torn.

Advil®

MULTI-SYMPTOM COLD & FLU

Ibuprofen 200 mg Pain Reliever/Fever Reducer **(NSAID)**
Phenylephrine HCl 10 mg Nasal Decongestant
Chlorpheniramine Maleate 4 mg Antihistamine

- ✓ Headache
- ✓ Fever
- ✓ Body Aches
- ✓ Sinus Pressure
- ✓ Nasal Congestion
- ✓ Runny Nose
- ✓ Sneezing
- ✓ Nasal Swelling

ONE TABLET DOSE!

2 Packets of 1 Coated Tablet

Advil® MULTI-SYMPTOM COLD & FLU

For most recent product information, visit **www.advil.com**

Questions or comments?
Call weekdays from **9 AM to 5 PM EST** toll free at **1-800-88-ADVIL**

Pfizer, Madison, NJ 07940 USA © 2018 Pfizer Inc.

PAA110483.FDA01

Reference ID: 4381479

PAA110483.FDA01

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

LIFT HERE

Drug Facts

Active ingredients (in each tablet) — Purposes

Chlorpheniramine maleate 4 mg ..... Antihistamine
Ibuprofen 200 mg (NSAID)* ..... Pain reliever/Fever reducer
Phenylephrine HCl 10 mg ..... Nasal decongestant
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves these symptoms associated with hay fever or other upper respiratory allergies, and the common cold:
  - runny nose
  - itchy, watery eyes
  - itching of the nose or throat
  - sneezing
  - nasal congestion
  - sinus pressure
  - headache
  - minor body aches and pains
  - fever
- reduces swelling of the nasal passages
- temporarily restores freer breathing through the nose

PAA110482.FDA01

PANEL 1

Drug Facts (continued)

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.
Symptoms may include:
- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

Drug Facts (continued)

- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use
- in children under 12 years of age because this product contains too much medication for children under this age
- if you have ever had an allergic reaction to any other pain reliever/fever reducer

PANEL 2

Drug Facts (continued)

- right before or after heart surgery
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if
- you have a breathing problem such as emphysema or chronic bronchitis
- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn

Drug Facts (continued)

- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, glaucoma, have trouble urinating due to an enlarged prostate gland, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are
- under a doctor's care for any serious condition
- taking sedatives or tranquilizers
- taking any other product that contains phenylephrine, chlorpheniramine or any other nasal decongestant or antihistamine
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

PANEL 3

Drug Facts (continued)

When using this product
- take with food or milk if stomach upset occurs
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- may cause excitability especially in children

Stop use and ask a doctor if
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:

Drug Facts (continued)

- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- nasal congestion lasts for more than 7 days
- redness or swelling is present in the painful area
- you get nervous, dizzy, or sleepless
- symptoms continue or get worse
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child

PANEL 4

Drug Facts (continued)

or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- adults and children 12 years of age and over:
  - take 1 tablet every 4 hours while symptoms persist
  - do not take more than 6 tablets in any 24-hour period, unless directed by a doctor
- children under 12 years of age: do not use because this product contains too much medication for children under this age

Other information

- read all warnings and directions before use. Keep carton.

Drug Facts (continued)

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

acesulfame potassium, artificial flavors, carnauba wax, colloidal silicon dioxide, corn starch, croscarmellose sodium, glycerin, glyceryl behenate, hypromellose, lactic acid, lecithin, maltodextrin, medium-chain triglycerides, microcrystalline cellulose, pharmaceutical ink, polydextrose, polyvinyl alcohol, pregelatinized starch, propyl gallate, silicon dioxide, sucralose, synthetic iron oxide, talc, titanium dioxide, triacetin, xanthan gum

Questions or comments?

Call weekdays from **9 AM to 5 PM EST** toll free at **1-800-88-ADVIL**

BASE PANEL

PAA110482.FDA01

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Reference ID: 4381479

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

LIFT HERE

Drug Facts

Active ingredients (in each tablet) — Purposes

Chlorpheniramine maleate 4 mg ..... Antihistamine
Ibuprofen 200 mg (NSAID)* ..... Pain reliever/Fever reducer
Phenylephrine HCl 10 mg ..... Nasal decongestant
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves these symptoms associated with the common cold or flu:
  - headache
  - fever
  - minor body aches and pains
  - nasal congestion
  - runny nose
  - sinus pressure
  - sneezing
- reduces swelling of the nasal passages
- temporarily restores freer breathing through the nose

PAA110484.FDA01

PANEL 1

Drug Facts (continued)

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
- are age 60 or older
- have had stomach ulcers or bleeding problems

Drug Facts (continued)

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use
- in children under 12 years of age
- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

PANEL 2

Drug Facts (continued)

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if
- you have a breathing problem such as emphysema or chronic bronchitis
- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn

Drug Facts (continued)

- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, glaucoma, have trouble urinating due to an enlarged prostate gland, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are
- under a doctor's care for any serious condition
- taking sedatives or tranquilizers
- taking any other product that contains phenylephrine, chlorpheniramine or any other nasal decongestant or antihistamine
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

PANEL 3

Drug Facts (continued)

When using this product
- take with food or milk if stomach upset occurs
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- may cause excitability especially in children

Stop use and ask a doctor if
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:

Drug Facts (continued)

- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- nasal congestion lasts for more than 7 days
- redness or swelling is present in the painful area
- you get nervous, dizzy, or sleepless
- symptoms continue or get worse
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely

PANEL 4

Drug Facts (continued)

directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
- **do not take more than directed**
- adults and children 12 years of age and over:
  - take 1 tablet every 4 hours while symptoms persist
  - do not take more than 6 tablets in any 24-hour period, unless directed by a doctor
- children under 12 years of age: do not use

Other information
- read all warnings and directions before use. Keep carton.

Drug Facts (continued)

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients
acesulfame potassium, artificial flavors, carnauba wax, colloidal silicon dioxide, corn starch, croscarmellose sodium, glycerin, glyceryl behenate, hypromellose, lactic acid, lecithin, maltodextrin, medium-chain triglycerides, microcrystalline cellulose, pharmaceutical ink, polydextrose, polyvinyl alcohol, pregelatinized starch, propyl gallate, silicon dioxide, sucralose, synthetic iron oxide, talc, titanium dioxide, triacetin, xanthan gum

Questions or comments?
Call weekdays from **9 AM to 5 PM EST** toll free at **1-800-88-ADVIL**

BASE PANEL

PAA110484.FDA01

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Reference ID: 4381479

PAA112204.FDA01

FRONT

MANUFACTURER'S COUPON | EXPIRES: XX/XX/XXXX

SAVE $2.00

on Any **Advil® Cold, Sinus or Allergy** Product

BACK

MANUFACTURER'S COUPON | EXPIRES: XX/XX/XXXX

SAVE $2.00

on Any **Advil®** Cold, Sinus or Allergy Product

030573-037439

FPO

CONSUMER: LIMIT ONE COUPON PER PURCHASE of products and quantities stated. LIMIT OF 4 LIKE COUPONS in same shopping trip. Coupons not authorized if purchasing products for resale. Void if transferred, sold, auctioned, reproduced or altered from original. Any other use constitutes fraud. You may pay sales tax. Do not send to Pfizer Consumer Healthcare. DEALER: Sending to Pfizer Consumer Healthcare 616, NCH Marketing Services, P.O. Box 880001, El Paso, TX 88588-0001 signifies compliance with the "Pfizer Consumer Healthcare Redemption Policy", copy available by writing to the above address. Cash value is 1/100th of 1 cent. © 2018 Pfizer Inc.

PAA112204.FDA01

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

THERESA M MICHELE
01/25/2019 04:16:23 PM