


CORTISPORIN-TC- colistin sulfate, neomycin sulfate, thonzonium bromide and hydrocortisone acetate suspension
Endo Pharmaceuticals, Inc.

Cortisporin®TC Otic with Neomycin and Hydrocortisone (colistin sulfate—neomycin sulfate—thonzonium bromide—hydrocortisone acetate otic suspension)

DESCRIPTION

Cortisporin®TC Otic with Neomycin and Hydrocortisone (colistin sulfate—neomycin sulfate—thonzonium bromide—hydrocortisone acetate otic suspension) is a sterile antibacterial and anti-inflammatory aqueous suspension containing in each mL: Colistin base activity, 3 mg (as the sulfate); Neomycin base activity, 3.3 mg (as the sulfate); Hydrocortisone acetate, 10 mg (1%); Thonzonium bromide, 0.5 mg (0.05%); Polysorbate 80, acetic acid, and sodium acetate in a buffered aqueous vehicle. Thimerosal (mercury derivative), 0.002%, is added as a preservative. It is a nonviscous liquid, buffered at pH 5, for instillation into the canal of the external ear or direct application to the affected aural skin.

The structural formulas of colistin sulfate (mixture of Colistin A & B), neomycin sulfate (mixture of neomycin A, B & C), hydrocortisone acetate ((11β)-21-(acetyloxy)-11,17-dihydroxypregn methyl]-2 pyrimidinylamino] ethyl]-N,N-dimethyl-1-hexadecanaminium, bromide) are represented below:




Thonzonium Bromide




Dbu is L - α, γ - diaminobutyric acid; R is 5 - methylheptyl in Colistin A and 5-methylhexyl in Colistin B)


Colistin sulfate




Neomycin A



Neomycin B Sulfate



Hydrocortisone Acetate



Neomycin C Sulfate

CLINICAL PHARMACOLOGY

Colistin sulfate is a polypeptide antibiotic which penetrates into and disrupts the bacterial cell membrane. Neomycin sulfate is an aminoglycoside antibiotic which inhibits protein synthesis, disrupting the normal cycle of ribosomal function. Hydrocortisone acetate is a corticosteroid hormone which is thought to act by regulating the rate of protein synthesis; it controls

inflammation, edema, pruritus and other dermal reactions. Corticosteroids suppress the inflammatory response to a variety of agents and they may delay healing. Since corticoids may inhibit the body's defense mechanism against infection, a concomitant antimicrobial drug may be used when this inhibition is considered to be clinically significant in a particular case.

The relative potency of corticosteroids depends on the molecular structure, concentration, and release from the vehicle.

Thonzonium bromide is a surface-active agent that promotes tissue contact by dispersion and penetration of the cellular debris and exudate.

Microbiology

Together, colistin sulfate and neomycin sulfate have bactericidal activity against most strains of the following microorganisms, both *in vitro* and in clinical infections as described in the [INDICATIONS AND USAGE](#) section.

Aerobic gram-positive microorganisms:

Staphylococcus aureus.

Aerobic gram-negative microorganisms:

Enterobacter aerogenes

Escherichia coli

Klebsiella pneumoniae

Pseudomonas aeruginosa.

Susceptibility Tests:

It is not recommended that colistin sulfate or neomycin sulfate be routinely tested and reported by clinical microbiology laboratories.

INDICATIONS AND USAGE

Cortisporin[®]TC Otic is indicated for the treatment of superficial bacterial infections of the external auditory canal, caused by organisms susceptible to the action of the antibiotics; and for the treatment of infections of mastoidectomy and fenestration cavities, caused by organisms susceptible to the antibiotics.

CONTRAINDICATIONS

This product is contraindicated in those individuals who have shown hypersensitivity to any of its components.

This product should not be used if the external auditory canal disorder is suspected or known to be due to cutaneous viral infection (e.g., herpes simplex virus or varicella zoster virus).

WARNINGS

Neomycin can induce permanent sensorineural hearing loss due to cochlear damage, mainly destruction of hair cells in the organ of Corti. The risk is greater with prolonged use. Therapy should be limited to 10 consecutive days. (See [PRECAUTIONS-General](#).) Patients being treated with eardrops containing neomycin should be under close clinical observation. Cortisporin® TC Otic should be used cautiously in any patient with a perforated tympanic membrane.

Neomycin sulfate may cause cutaneous sensitization. A precise incidence of hypersensitivity reactions (primarily skin rash) due to topical neomycin is not known. Discontinue promptly if sensitivity or irritation occurs.

When using neomycin-containing products to control secondary infection in the chronic dermatoses, such as chronic otitis externa or stasis dermatitis, it should be borne in mind that the skin in these conditions is more liable than is normal skin to become sensitized to many substances, including neomycin. The manifestation of sensitization to neomycin is usually a low-grade reddening with swelling, dry scaling, and itching; it may be manifest simply as a failure to heal. Periodic examination for such signs is advisable, and the patient should be told to discontinue the product if they are observed. These symptoms regress quickly on withdrawing the medication. Neomycin containing applications should be avoided for the patient thereafter.

PRECAUTIONS

General

As with any other antibiotic preparation, prolonged treatment may result in overgrowth of nonsusceptible organisms and fungi. If the infection is not improved after one week, cultures should be repeated to verify the identity of the organism and to determine whether therapy should be changed.

Treatment should not be continued for longer than ten days.

Allergic cross-reactions may occur which could prevent the use of any or all of the aminoglycoside antibiotics for the treatment of future infections.

Information for Patients

Avoid contaminating the dropper with material from the ear, fingers, or other source. This caution is necessary if the sterility of the drops is to be preserved.

If sensitization or irritation occurs, discontinue use immediately and contact your physician.

Do not use in the eyes.

If you prefer to warm the medication before using it, do not heat the suspension above body temperature in order to avoid loss of potency.

SHAKE WELL BEFORE USING.

Laboratory Tests

Systemic effects of excessive levels of hydrocortisone may include a reduction in the number of circulating eosinophils and a decrease in urinary excretion of 17-hydroxycorticosteroids.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term animal carcinogenicity studies have not been performed with colistin or neomycin, or Cortisporin[®]TC Otic. An increased incidence of chromosome aberrations in human lymphocytes has been reported following *in vitro* exposure to colistin or neomycin.

Fertility studies have not been performed with neomycin, but reports from the scientific literature suggest that it may decrease spermatogenesis in rats. No adverse effects on fertility were observed in male or female rats given intramuscular doses of colistimethate sodium, the methanesulfonate salt of colistin, up to 20 mg/kg (equivalent to 9.3 mg/kg of colistin base). This is approximately 30 times the clinical daily dose based on body surface area, assuming 100% absorption from the ear; however, significant systemic levels of colistin or neomycin would not be anticipated in humans when Cortisporin[®]TC Otic is used as directed.

Long term studies in rodents showed no evidence of carcinogenicity attributable to oral administration of corticosteroids. Mutagenicity studies with hydrocortisone were negative. Studies have not been performed to evaluate the effect on fertility of topical corticosteroids.

Pregnancy

Teratogenic Effects

Pregnancy Category C

There are no adequate and well controlled studies of Cortisporin[®]TC Otic in pregnant women. It is not known whether Cortisporin[®]TC Otic can cause fetal harm when administered to a pregnant woman.

Colistimethate sodium, the methanesulfonate salt of colistin, was not teratogenic in rats or rabbits given intramuscular doses up to 20 mg/kg (equivalent to 9.3 mg/kg of colistin base, approximately 30 times (rats) or 55 times (rabbits) the clinical daily dose based on body surface area and assuming 100% absorption from the ear). Increased resorptions were observed in rabbits at 20 mg/kg, but not 10 mg/kg (equivalent to 4.15 mg/kg of colistin base). Decreased pup survival at weaning was observed in rats at 20 mg/kg, a maternally toxic dose of colistin, but not 10 mg/kg. Colistin has not been shown to have any adverse effects on the developing embryo or fetus at doses relevant to the amount that will be delivered ototopically at the recommended clinical doses.

Although aminoglycosides can cause congenital deafness in humans if administered during pregnancy, significant systemic levels of neomycin would not be anticipated when Cortisporin[®]TC Otic is used as directed.

Corticosteroids are generally teratogenic in laboratory animals when administered systemically at relatively low dosage levels. The more potent corticosteroids have been shown to be teratogenic after dermal application in laboratory animals.

Cortisporin[®]TC Otic should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Nursing Mothers

Hydrocortisone and colistin sulfate appear in human milk following oral administration of the drugs. Since systemic absorption of these drugs may occur when they are used topically, caution should be exercised when Cortisporin[®]TC Otic Suspension is used by a nursing woman.

Pediatric Use

See [DOSAGE AND ADMINISTRATION](#).

The safety and effectiveness of Cortisporin[®]TC Otic in infants below one year of age have not been established. The efficacy of Cortisporin[®]TC Otic in pediatric patients one year or older in the treatment of superficial bacterial infections of the external auditory canal and for the treatment of infections of mastoidectomy and fenestration cavities has been demonstrated in a controlled clinical trial.

Geriatric Use

No overall differences in safety or effectiveness have been observed between elderly and younger patients.

ADVERSE REACTIONS

Neomycin occasionally causes skin sensitization.

Ototoxicity (see [WARNINGS](#) section) and nephrotoxicity have also been reported.

Adverse reactions have occurred with topical use of antibiotic combinations. Exact incidence figures are not available since no denominator of treated patients is available. The reaction occurring most often is allergic sensitization. In one clinical study, using a 20% neomycin patch, neomycin-induced allergic skin reactions occurred in two of 2,175 (0.09%) individuals in the general population. In another study the incidence was found to be approximately 1%.

The following local adverse events have been reported with topical corticosteroids, especially under occlusive dressings: burning, itching, irritation, dryness, folliculitis, hypertrichosis, acneiform eruptions, hypopigmentation, perioral dermatitis, allergic contact dermatitis, maceration of the skin, secondary infection, skin atrophy, striae, and miliaria.

For medical advice about adverse reactions contact your medical professional. To report SUSPECTED ADVERSE REACTIONS, contact Endo Pharmaceuticals Inc. at 1-800-462-3636 or MEDWATCH at 1-800-FDA-1088 or <http://www.fda.gov/medwatch/>.

DOSAGE AND ADMINISTRATION

Therapy with this product should be limited to 10 days. (See [WARNINGS](#).)

The external auditory canal should be thoroughly cleansed and dried with a sterile cotton applicator.

When using the calibrated dropper:

For adults, 5 drops of the suspension should be instilled into the affected ear 3 or 4 times daily. For pediatric patients, 4 drops are suggested because of the smaller capacity of the ear canal.

The patient should lie with the affected ear upward and then the drops should be instilled. This position should be maintained for 5 minutes to facilitate penetration of the drops into the ear canal. Repeat, if necessary, for the opposite ear.

If preferred, a cotton wick may be inserted into the canal and then the cotton may be saturated with the suspension. This wick should be kept moist by adding further solution every 4 hours. The wick should be replaced at least once every 24 hours.

HOW SUPPLIED

Cortisporin[®]TC Otic is supplied as:

NDC 63481-529-10..... 10 mL bottle with dropper

Each mL contains: Colistin sulfate equivalent to 3 mg of colistin base activity, Neomycin sulfate equivalent to 3.3 mg neomycin base activity, Hydrocortisone acetate 10 mg (1%), Thonzonium bromide 0.5 mg (0.05%), and Polysorbate 80 in an aqueous vehicle buffered with acetic acid and sodium acetate. Thimerosal (mercury derivative) 0.002% is added as a preservative.

A sterilized dropper-cap assembly for use on the bottle of suspension is included in the package.

Shake well before using.

Store at 20°–25°C (68°–77°F).

Rx only.

Distributed by:
Endo Pharmaceuticals Inc.
Malvern, PA 19355

OS529E-01-90-01

I01/2019

3003827

NDC 63481-529-10

Rx Only

Cortisporin®-TC

(colistin sulfate–neomycin sulfate–
thonzonium bromide–hydrocortisone
acetate otic suspension)

STERILE **10 mL**
FOR USE IN EARS ONLY

 **endo**
pharmaceuticals
Endo International Company



Each mL contains: Colistin sulfate equivalent to 3 mg of colistin base activity, neomycin sulfate equivalent to 3.3 mg neomycin base activity, hydrocortisone acetate 10 mg (1%), thonzonium bromide 0.5 mg (0.05%), and Polysorbate 80 in an aqueous vehicle buffered with acetic acid and sodium acetate. Thimerosal (mercury derivative) 0.002% added as a preservative.

SHAKE WELL BEFORE USING.

Usual Adult Dosage: See full prescribing information.

Warning: For external use only.

Store between 20° to 25°C (68° to 77°F).
(See USP controlled room temperature.)

Distributed by:
Endo Pharmaceuticals Inc.
Malvern, PA 19355

R10/18

LA529E-52-90-02




(01)00363481 529102

3 0 0 3 8 2 6 A

LOT

EXP



3 63481-529-10 2

NDC 63481-529-10

Cortisporin®-TC

*(colistin sulfate–neomycin sulfate–
thonzonium bromide–hydrocortisone
acetate otic suspension)*

NOTE – The enclosed dropper has been sterilized. If dropper wrapper is broken or open, sterility cannot be guaranteed.

Rx Only

**STERILE
Shake Well Before Using**

10 mL bottle with dropper



FOR USE IN EARS ONLY



NDC 63481-529-10
Cortisporin®-TC
*(colistin sulfate–neomycin sulfate–
thonzonium bromide–hydrocortisone
acetate otic suspension)*

10 mL bottle with dropper

347

Each mL contains: Colistin sulfate equivalent to 3 mg of colistin base activity, Neomycin sulfate equivalent to 3.3 mg neomycin base activity, Hydrocortisone acetate 10 mg (1%), Thonzonium bromide 0.5 mg (0.05%), and Polysorbate 80 in an aqueous vehicle buffered with acetic acid and sodium acetate. Thimerosal (mercury derivative) 0.002% added as a preservative.
Usual Adult Dosage Five (5) drops instilled into affected ear 3 or 4 times a day using calibrated dropper.
For full prescribing information, see package insert.
Warning: For external use only.
Store at 20° to 25°C (68° to 77°F)(See USP controlled room temperature.)
See bottom of carton for lot number and expiration date.
Distributed by:
Endo Pharmaceuticals Inc.
Malvern, PA 19355
UC529E-52-90-02
R10/18

NDC 63481-529-10

Cortisporin®-TC

*(colistin sulfate–neomycin sulfate–
thonzonium bromide–hydrocortisone
acetate otic suspension)*

NOTE – The enclosed dropper has been sterilized. If dropper wrapper is broken or open, sterility cannot be guaranteed.

Rx Only

**STERILE
Shake Well Before Using**


10 mL bottle with dropper



FOR USE IN EARS ONLY



3003828A



NDC 63481-529-05

Rx Only

Cortisporin®-TC

(colistin sulfate—neomycin sulfate—
thonzonium bromide—hydrocortisone
acetate otic suspension)

STERILE **5 mL**
FOR USE IN EARS ONLY

 **endo**
pharmaceuticals
an endo international company

SAMPLE USE ONLY
NOT FOR RESALE



Each mL contains: Colistin sulfate equivalent to 3 mg of colistin base activity, neomycin sulfate equivalent to 3.3 mg neomycin base activity, hydrocortisone acetate 10 mg (1%), thonzonium bromide 0.5 mg (0.05%), and Polysorbate 80 in an aqueous vehicle buffered with a citric acid and sodium acetate. Thimerosal (mercury derivative) 0.002% added as a preservative.

SHAKE WELL BEFORE USING.

Usual Adult Dosage: See full prescribing information.

Warning: For external use only.

Store between 20° to 25°C (68° to 77°F).
(See USP controlled room temperature.)

Distributed by:
Endo Pharmaceuticals Inc.
Malvern, PA 19355

11Q/18

LA529E-52-90-01



(01)0036348 1529058

3 0 0 0 0 0 0 0

LOT

EXP



2 63481-529-05 8

NDC 63481-529-05

SAMPLE USE ONLY - NOT FOR RESALE

Cortisporin®-TC

(colistin sulfate-neomycin sulfate-thonzonium bromide-hydrocortisone acetate otic suspension)

NOTE – The enclosed dropper has been sterilized. If dropper wrapper is broken or open, sterility cannot be guaranteed.

Rx Only

STERILE
Shake Well Before Using

5 mL bottle with dropper

SAMPLE USE ONLY
NOT FOR RESALE



FOR USE IN EARS ONLY



5 mL bottle with dropper

Cortisporin®-TC
(colistin sulfate-neomycin sulfate-thonzonium bromide-hydrocortisone acetate otic suspension)

NDC 63481-529-05

SAMPLE USE ONLY - NOT FOR RESALE

346

Each mL contains: Colistin sulfate equivalent to 3 mg of colistin base activity, Neomycin sulfate equivalent to 3.3 mg neomycin base activity, Hydrocortisone acetate 10 mg (1%), Thonzonium bromide 0.5 mg (0.05%), and Polysorbate 80 in an aqueous vehicle buffered with acetic acid and sodium acetate. Thimerosal (mercury derivative) 0.002% added as a preservative.

Usual Adult Dosage Five (5) drops instilled into affected ear 3 or 4 times a day using calibrated dropper.

For full prescribing information, see package insert.

Warning: For external use only. **Store at 20° to 25°C (68° to 77°F)**(See USP controlled room temperature.)

See bottom of carton for lot number and expiration date. Distributed by: **Endo Pharmaceuticals Inc.** Malvern, PA 19355

UC529E-52-90-01
110/18

NDC 63481-529-05

SAMPLE USE ONLY - NOT FOR RESALE

Cortisporin®-TC

(colistin sulfate-neomycin sulfate-thonzonium bromide-hydrocortisone acetate otic suspension)

NOTE – The enclosed dropper has been sterilized. If dropper wrapper is broken or open, sterility cannot be guaranteed.

Rx Only

STERILE
Shake Well Before Using

5 mL bottle with dropper

FOR USE IN EARS ONLY



000000



This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

DMITRI IARIKOV
01/05/2019 01:32:00 PM