

## HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use DYSFORT safely and effectively. See full prescribing information for DYSFORT.

DYSFORT® (abobotulinumtoxinA) for injection, for intramuscular use

Initial U.S. Approval: 2009

### WARNING: DISTANT SPREAD OF TOXIN EFFECT

*See full prescribing information for complete boxed warning*

The effects of DYSFORT and all botulinum toxin products may spread from the area of injection to produce symptoms consistent with botulinum toxin effects. These symptoms have been reported hours to weeks after injection. Swallowing and breathing difficulties can be life-threatening and there have been reports of death. The risk of symptoms is probably greatest in children treated for spasticity but symptoms can occur in adults, particularly in those patients who have underlying conditions that would predispose them to these symptoms (5.1).

### RECENT MAJOR CHANGES

Indications and Usage (1.4)	9/2019
Dosage and Administration (2.2, 2.6)	9/2019
Warnings and Precautions (5.3, 5.6)	11/2018

### INDICATIONS AND USAGE

DYSFORT is an acetylcholine release inhibitor and a neuromuscular blocking agent indicated for:

- The treatment of cervical dystonia in adults (1.1)
- The temporary improvement in the appearance of moderate to severe glabellar lines associated with procerus and corrugator muscle activity in adults < 65 years of age (1.2)
- The treatment of upper and lower limb spasticity in adults (1.3)
- The treatment of upper limb spasticity in pediatric patients 2 years of age and older, excluding spasticity caused by cerebral palsy (1.4)
- The treatment of lower limb spasticity in pediatric patients 2 years of age and older (1.4)

### DOSAGE AND ADMINISTRATION

Preparation of DYSFORT Solution for Administration (2.2)

- Once reconstituted, store in original container in a refrigerator at 2°C to 8°C (36°F to 46°F) and use within 24 hours
- Do not freeze after reconstitution
- Reconstitution instructions are specific for the 300 Unit and 500 Unit vials
- Reconstituted DYSFORT is intended for intramuscular injection only. After reconstitution, DYSFORT should be used for only one injection session and for only one patient

#### Cervical Dystonia (2.3)

- Initial dose is 500 Units given intramuscularly as a divided dose among the affected muscles
- Re-treatment every 12 to 16 weeks or longer, as necessary, based on return of clinical symptoms with doses administered between 250 Units and 1000 Units to optimize clinical benefit
- Re-treatment should not occur in intervals of less than 12 weeks
- Titrate in 250 Unit steps according to patient's response

#### Glabellar Lines (2.4)

- Administer a total dose of 50 Units, divided in five equal aliquots of 10 Units each, intramuscularly to affected muscles to achieve clinical effect
- Re-treatment should be administered no more frequently than every 3 months

#### Spasticity in Adults (2.5)

- Select dose based on muscles affected, severity of spasticity, and treatment and adverse reaction history with botulinum toxins
- Dosing for upper limb spasticity: between 500 Units and 1000 Units
- Dosing for lower limb spasticity: up to 1500 Units
- The maximum recommended total dose per treatment session (upper and lower limb combined) in adults is 1500 Units
- Re-treatment, based on return of clinical symptoms, should not occur in intervals of less than 12 weeks

#### Spasticity in Pediatric Patients (2.6)

- Select dose based on the affected muscle, severity of spasticity, and treatment and adverse reaction history with all botulinum toxins.
- Recommended dosing for upper limb spasticity, excluding spasticity caused by cerebral palsy: 8 Units/kg to 16 Units/kg per limb. The maximum recommended total dose administered per treatment session must not exceed 16 Units/kg or 640 Units, whichever is lower.
- Recommended dosing for lower limb spasticity: 10 Units/kg to 15 Units/kg per limb. Total dose per treatment session must not exceed 15 Units/kg for unilateral lower limb injections, 30 Units/kg for bilateral injections, or 1000 Units, whichever is lower.
- The maximum recommended total dose per treatment session is 30 Units/kg or 1000 Units, whichever is lower. Re-treatment, based on return of clinical symptoms, should not occur in intervals of less than 3 months.

### DOSAGE FORMS AND STRENGTHS

- For Injection: 300 Units or 500 Units lyophilized powder in a single-dose vial (3)

### CONTRAINDICATIONS

- Hypersensitivity to:
  - any botulinum toxin product or excipients (4, 5.3)
  - cow's milk protein (4, 5.3)
- Infection at the proposed injection site(s) (4)

### WARNINGS AND PRECAUTIONS

- The potency units of DYSFORT are not interchangeable with other preparations of botulinum toxin products (5.2)
- Immediate medical attention may be required in cases of respiratory, speech or swallowing difficulties (5.4)
- Recommended dose and frequency of administration should not be exceeded (5.5)
- Dry eye may occur with glabellar line treatment; if symptoms persist, consider referring patient to an ophthalmologist (5.6)
- Concomitant neuromuscular disorder may exacerbate clinical effects of treatment (5.7)

### ADVERSE REACTIONS

Most commonly observed adverse reactions are (6.1):

#### Cervical Dystonia

(≥5%): muscular weakness, dysphagia, dry mouth, injection site discomfort, fatigue, headache, musculoskeletal pain, dysphonia, injection site pain and eye disorders

#### Glabellar Lines

(≥2%): nasopharyngitis, headache, injection site pain, injection site reaction, upper respiratory tract infection, eyelid edema, eyelid ptosis, sinusitis, nausea, and blood present in urine

#### Spasticity in Adults

- Upper limb spasticity (≥4%): muscular weakness
- Lower limb spasticity (≥5%): falls, muscular weakness, and pain in extremity

#### Spasticity in Pediatric Patients

- Upper limb spasticity (≥10%): upper respiratory tract infection and pharyngitis
- Lower limb spasticity (≥10%): nasopharyngitis, cough, and pyrexia

To report SUSPECTED ADVERSE REACTIONS, contact Ipsen Biopharmaceuticals, Inc. at 855-463-5127 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch)

### DRUG INTERACTIONS

- Concomitant use of DYSFORT and aminoglycosides or other agents interfering with neuromuscular transmission or muscle relaxants, should be observed closely because effect of DYSFORT may be potentiated (7.1, 7.4)
- Anticholinergic drugs may potentiate systemic anticholinergic effects (7.2)
- The effect of administering different botulinum neurotoxins during the course of treatment with DYSFORT is unknown (7.3)

### USE IN SPECIFIC POPULATIONS

- Administer DYSFORT with care in elderly patients, reflecting the greater frequency of concomitant disease and other drug therapy (8.5)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide

Revised: 9/2019

























Other adverse reactions with incidences of less than 5% in the DYSPORT 500 Units group in the double-blind phase of clinical trials included dizziness in 3.5% of DYSPORT-treated patients and 1% of placebo-treated patients, and muscle atrophy in 1% of DYSPORT-treated patients and in none of the placebo-treated patients.

#### Laboratory Findings

Patients treated with DYSPORT exhibited a small increase from baseline (0.23 mol/L) in mean blood glucose relative to placebo-treated patients. This was not clinically significant among patients in the development program but could be a factor in patients whose diabetes is difficult to control.

#### Electrocardiographic Findings

ECG measurements were only recorded in a limited number of patients in an open-label study without a placebo or active control. This study showed a statistically significant reduction in heart rate compared to baseline, averaging about three beats per minute, observed thirty minutes after injection.

#### Glabellar Lines

In placebo-controlled clinical trials of DYSPORT, the most common adverse reactions ( $\geq 2\%$ ) following injection of DYSPORT were nasopharyngitis, headache, injection site pain, injection site reaction, upper respiratory tract infection, eyelid edema, eyelid ptosis, sinusitis, nausea, and blood present in urine.

Table 9 reflects exposure to DYSPORT in 398 patients 19 to 75 years of age who were evaluated in the randomized, placebo-controlled clinical studies that assessed the use of DYSPORT for the temporary improvement in the appearance of glabellar lines [see *Clinical Studies (14.2)*]. Adverse reactions of any cause occurred in 48% of the DYSPORT-treated patients and 33% of the placebo-treated patients.

**Table 9: Most Common Adverse Reactions with > 1% Incidence in Pooled, Placebo-Controlled Trials for Glabellar Lines**

Adverse Reactions by Body System	DYSPORT (N=398) %*	Placebo (N=496) %*
<b>Any Adverse Reaction</b>	48	33
<b>Eye Disorders</b>		
Eyelid Edema	2	0
Eyelid Ptosis	2	<1
<b>Gastrointestinal Disorders</b>		
Nausea	2	1
<b>General Disorders and Administration Site Conditions</b>		
Injection Site Pain	3	2
Injection Site Reaction	3	<1
<b>Infections and Infestations</b>		
Nasopharyngitis	10	4
Upper Respiratory Tract Infection	3	2
Sinusitis	2	1
<b>Investigations</b>		
Blood Present in Urine	2	<1
<b>Nervous System Disorders</b>		
Headache	9	5

\*Patients who received treatment with placebo and DYSPORT are counted in both treatment columns.

In the clinical trials safety database, where some patients received up to twelve treatments with DYSPORT, adverse reactions were reported for 57% (1425/2491) of patients. The most frequently reported of these adverse reactions were headache, nasopharyngitis, injection site pain, sinusitis, URI, injection site bruising, and injection site reaction (numbness, discomfort, erythema, tenderness, tingling, itching, stinging, warmth, irritation, tightness, swelling).

Adverse reactions that occurred after repeated injections in 2–3% of the population included bronchitis, influenza, pharyngolaryngeal pain, cough, contact dermatitis, injection site swelling, and injection site discomfort.

The incidence of eyelid ptosis did not increase in the long-term safety studies with multiple re-treatments at intervals  $\geq$  three months. The majority of the reports of eyelid ptosis were mild to moderate in severity and resolved over several weeks [see *Dosage and Administration (2.4)*].

## Spasticity in Adults

### Injection Site Reactions

Injection site reactions (e.g., pain, bruising, hemorrhage, erythema/hematoma, etc.) have occurred following administration of DYSPORT in adults treated for spasticity.

### Upper Limb Spasticity in Adults

Table 10 lists the adverse reactions that occurred in  $\geq 2\%$  of patients in any DYSPORT dose group and more frequent than placebo in double-blind studies evaluating the treatment of upper limb spasticity in adults. The most common adverse reactions ( $\geq 4\%$ ) in any DYSPORT dose group was muscular weakness.

**Table 10: Most Common Adverse Reactions Observed in at Least 2% of Patients Treated in Pooled, Double-Blind Trials of Adults with Upper Limb Spasticity Reported More Frequently than with Placebo**

Adverse Reactions	DYSPORT		Placebo (N=279) %
	500 Units (N=197) %	1000 Units (N=194) %	
<b>Infections and infestations</b>			
Influenza	1	2	1
Infection	1	2	1
<b>Musculoskeletal and connective tissue disorders</b>			
Muscular weakness	2	4	1
Pain in extremity	0	2	1
Back pain	1	2	1
<b>Nervous system disorders</b>			
Headache	1	2	1
Convulsion	2	2	1
Syncope	1	2	0
Hypoesthesia	0	2	<1
Partial seizures	0	2	0
<b>General disorders and administration site conditions</b>			
Fatigue	2	2	0
Asthenia	2	1	<1
<b>Injury, poisoning and procedural complications</b>			
Fall	2	3	2
Injury	2	2	1
Contusion	1	2	<1
<b>Gastrointestinal disorders</b>			
Diarrhea	1	2	<1
Constipation	0	2	1
<b>Investigation</b>			
Blood triglycerides increased	2	1	0
<b>Respiratory, thoracic and mediastinal disorders</b>			
Cough	1	2	1
<b>Vascular disorders</b>			
Hypertension	1	2	<1
<b>Psychiatric disorders</b>			
Depression	2	3	1

### Less Common Adverse Reactions

In a pooled analysis of clinical studies, adverse reactions with an incidence of less than 2% reported in DYSPORT treatment groups included dysphagia 0.5%, gait disturbance 0.5%, hypertonia 0.5%, and sensation of heaviness 0.3%.

### Lower Limb Spasticity in Adults

The data described below reflect exposure to DYSPORT in 255 adults with lower limb spasticity. Of this population, 89% were Caucasian, 66% male, and the median age was 55 years (range 23–77 years). Table 11 lists the adverse reactions that occurred in  $\geq 2\%$  of patients in any DYSPORT dose group and more frequent than placebo in the double-blind study evaluating the treatment of lower limb spasticity in adults. The most common of these adverse reactions ( $\geq 5\%$ ) in any DYSPORT dose group were falls, muscular weakness, and pain in extremity.

**Table 11: Adverse Reactions Observed in at Least 2% of Patients Treated in the Double-Blind Trial of Adults with Lower Limb Spasticity and Reported More Frequently than with Placebo**

<b>Adverse Reactions</b>	<b>DYSPORT 1000 U (N=127) %</b>	<b>DYSPORT 1500 U (N=128) %</b>	<b>Placebo (N=130) %</b>
<b>Musculoskeletal and connective tissue disorders</b>			
Muscular weakness	2	7	3
Pain in extremity	6	6	2
Arthralgia	4	2	1
<b>Injury, poisoning and procedural complications</b>			
Fall	9	6	3
<b>Nervous system disorders</b>			
Headache	0	3	1
<b>General disorders and administration site conditions</b>			
Fatigue	1	4	0
Influenza-like illness	2	0	0
Edema peripheral	2	0	0
<b>Investigations</b>			
Alanine aminotransferase increased	2	0	1
<b>Gastrointestinal disorders</b>			
Constipation	0	2	1
<b>Psychiatric disorders</b>			
Depression	2	3	0
Insomnia	0	2	0

In the efficacy and safety studies of DYSPORT for the treatment of lower limb spasticity in adults, muscular weakness was reported more frequently in women (10%) treated with 1500 units of DYSPORT compared to men (5%). Falls were reported more frequently in patients 65 years of age and over [see *Use in Specific Populations (8.5)*].

#### Upper Limb Spasticity in Pediatric Patients

Table 12 reflects exposure to DYSPORT in 210 patients, 2 to 17 years of age, who were evaluated in a double blind, active-controlled, multicenter study in patients treated for upper limb spasticity [see *Clinical Studies (14.4)*]. The most commonly observed adverse reactions ( $\geq 10\%$  of patients) were: upper respiratory tract infection and pharyngitis.

**Table 12: Adverse Reactions Observed in  $\geq 3\%$  of Patients Treated in Double-Blind Study of Pediatric Patients with Upper Limb Spasticity and Reported More Frequently than Control Group**

<b>Adverse Reactions</b>	<b>DYSPORT 2 Units/kg<sup>1</sup> (N=70) %</b>	<b>DYSPORT 8 Units/kg (N=70) %</b>	<b>DYSPORT 16 Units/kg (N=70) %</b>
<b>Infections and infestations</b>			
Upper respiratory tract infection	7	9	11
Influenza	1	1	3
Pharyngitis <sup>2</sup>	9	6	10
<b>Gastrointestinal disorders</b>			
Nausea	0	3	1
<b>Musculoskeletal and connective tissue disorders</b>			
Muscular weakness	1	4	6
<b>Nervous system disorders</b>			
Headache	0	6	3
Epilepsy	1	0	4

1 Low dose active comparator arm

2 includes pharyngitis, pharyngitis streptococcal, pharyngotonsillitis

Additional adverse reactions occurring below 3% and considered to be drug related include: myalgia, pain in extremity, fatigue, influenza-like illness, injection site eczema, injection site bruising, injection site rash, injection site pain, and injection site swelling.

#### Lower Limb Spasticity in Pediatric Patients

Table 13 reflects exposure to DYSPORT in 160 patients, 2 to 17 years of age, who were evaluated in the randomized, placebo-controlled clinical study that assessed the use of DYSPORT for the treatment of unilateral or bilateral lower limb spasticity in pediatric cerebral palsy patients [see *Clinical Studies (14.4)*]. The most commonly observed adverse reactions ( $\geq 10\%$  of patients) were: upper respiratory tract infection, nasopharyngitis, influenza, pharyngitis, cough and pyrexia.

**Table 13: Adverse Reactions Observed in  $\geq 4\%$  of Patients Treated in the Double-Blind Trial of Pediatric Patients with Lower Limb Spasticity and Reported More Frequently than with Placebo**

Adverse Reactions	Unilateral		Bilateral		Placebo (N=79) %
	DYSPORT 10 Units/kg (N=43) %	DYSPORT 15 Units/kg (N=50) %	DYSPORT 20 Units/kg (N=37) %	DYSPORT 30 Units/kg (N=30) %	
<b>Infections and infestations</b>					
Nasopharyngitis	9	12	16	10	5
Bronchitis	0	0	8	7	3
<b>Respiratory, thoracic and mediastinal disorders</b>					
Cough	7	6	14	10	6
<b>General disorders and administration site conditions</b>					
Pyrexia	7	12	8	7	5
<b>Musculoskeletal and connective tissue disorders</b>					
Pain in extremity	0	2	5	7	5
<b>Nervous system disorders</b>					
Convulsion/Epilepsy	7	4	0	7	0

## 6.2 Immunogenicity

As with all therapeutic proteins, there is a potential for immunogenicity.

The incidence of antibody formation is highly dependent on the sensitivity and specificity of the assay. In addition, the observed incidence of antibody positivity in an assay may be influenced by several factors including assay methodology, sample handling, timing of sample collection, concomitant medications, and underlying disease. For these reasons, comparison of the incidence of antibodies across products in this class may be misleading.

#### Cervical Dystonia

About 3% of subjects developed antibodies (binding or neutralizing) over time with DYSPORT treatment.

#### Glabellar Lines

Testing for antibodies to DYSPORT was performed for 1554 subjects who had up to nine cycles of treatment. Two subjects (0.13%) tested positive for binding antibodies at baseline. Three additional subjects tested positive for binding antibodies after receiving DYSPORT treatment. None of the subjects tested positive for neutralizing antibodies.

#### Spasticity in Adults

##### Upper Limb Spasticity

From 230 subjects treated with DYSPORT and tested for the presence of binding antibodies, 5 subjects were positive at baseline and 17 developed antibodies after treatment. Among those 17 subjects, 10 subjects developed neutralizing antibodies. An additional 51 subjects from a separate repeat-dose study were tested for the presence of neutralizing antibodies only. None of the subjects tested positive.

In total, from the 281 subjects treated in the long-term studies and tested for the presence of neutralizing antibodies, 3.6% developed neutralizing antibodies after treatment. In the presence of binding and neutralizing antibodies to DYSPORT some patients continued to experience clinical benefit.

### *Lower Limb Spasticity*

From 367 subjects treated with DYSPORT and tested for the presence of binding antibodies, 4 subjects were positive at baseline and 2 developed binding antibodies after treatment. No subjects developed neutralizing antibodies. An additional 85 subjects from two separate studies were tested for the presence of neutralizing antibodies only. One subject tested positive for the presence of neutralizing antibodies.

In total, from the 452 subjects treated with DYSPORT and tested for the presence of neutralizing antibodies, 0.2% developed neutralizing antibodies after treatment.

### Spasticity in Pediatric Patients 2 Years of Age or Older

#### *Upper Limb Spasticity*

From 178 subjects treated with DYSPORT for up to 4 treatment cycles and tested for the presence of binding antibodies at baseline and end of study, 7 subjects previously receiving botulinum toxin injections had binding antibodies after treatment. Among those 7 subjects, 4 subjects (2.3%) developed neutralizing antibodies when tested in the mice bioassay. In the presence of binding and/or neutralizing antibodies to DYSPORT some patients continue to experience clinical benefit.

#### *Lower Limb Spasticity*

From 226 subjects treated with DYSPORT and tested for the presence of binding antibodies, 5 subjects previously receiving botulinum toxins were positive at baseline and 9 patients developed binding antibodies after injections. Among those 9 subjects, 3 subjects developed neutralizing antibodies, while one subject developed neutralizing antibodies from the 5 subjects testing positive for binding antibodies at baseline who previously received botulinum toxin injections.

From a separate repeat-dose study, 203 subjects were tested for the presence of neutralizing antibodies. Two subjects were positive for neutralizing antibodies at baseline and 5 subjects developed neutralizing antibodies after treatments. In total, from the 429 patients tested for the presence of neutralizing antibodies, 2.1% developed neutralizing antibodies after treatment. In the presence of binding and neutralizing antibodies to DYSPORT, some patients continued to experience clinical benefit.

## **6.3 Postmarketing Experience**

Because adverse reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

The following adverse reactions have been identified during post-approval use of DYSPORT: vertigo, photophobia, influenza-like illness, amyotrophy, burning sensation, facial paresis, hypoesthesia, erythema, dry eye, and excessive granulation tissue. Hypersensitivity reactions including anaphylaxis have been reported.

## **7 DRUG INTERACTIONS**

### **7.1 Aminoglycosides and Other Agents Interfering with Neuromuscular Transmission**

Co-administration of DYSPORT and aminoglycosides or other agents interfering with neuromuscular transmission (e.g., curare-like agents) should only be performed with caution because the effect of the botulinum toxin may be potentiated. If co-administered, observe the patient closely.

### **7.2 Anticholinergic Drugs**

Use of anticholinergic drugs after administration of DYSPORT may potentiate systemic anticholinergic effects such as blurred vision.

### **7.3 Other Botulinum Neurotoxin Products**

The effect of administering botulinum neurotoxin products including DYSPORT, at the same time or within several months of each other is unknown. Excessive weakness may be exacerbated by another administration of botulinum toxin prior to the resolution of the effects of a previously administered botulinum toxin.

### **7.4 Muscle Relaxants**

Excessive weakness may also be exaggerated by administration of a muscle relaxant before or after administration of DYSPORT.

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

#### Risk Summary

There are no adequate and well-controlled clinical studies with DYSPORT in pregnant women.

DYSPORT should only be used during pregnancy if the potential benefit justifies the potential risk to the fetus.

DYSPOSPORT produced embryo-fetal toxicity in relation to maternal toxicity when given to pregnant rats and rabbits at doses lower than or similar to the maximum recommended human dose (MRHD) of 1000 Units on a body weight (Units/kg) basis (*see Data*).

In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2-4% and 15-20%, respectively. The background risk of major birth defects and miscarriage for the indicated populations is unknown.

## Data

### *Animal Data*

In a study in which pregnant rats received daily intramuscular injections of DYSPOSPORT (2.2, 6.6, or 22 Units/kg on gestation days 6 through 17 or intermittently 44 Units/kg on gestation days 6 and 12 only) during organogenesis, increased early embryonic death was observed with both schedules at the highest tested doses (22 and 44 Units/kg), which were associated with maternal toxicity. The no-effect dose for embryo-fetal developmental toxicity was 2.2 Units/kg (less than the maximum recommended human dose [MRHD] on a body weight basis).

In a study in which pregnant rabbits received daily intramuscular injections of DYSPOSPORT (0.3, 3.3, or 6.7 Units/kg) on gestation days 6 through 19 or intermittently (13.3 Units/kg on gestation days 6 and 13 only) during organogenesis, no embryofetal data were available at the highest dose administered daily (6.7 Units/kg) because of premature death in all does at that dose. At the lower daily doses or with intermittent dosing, no adverse developmental effects were observed. All doses for which data were available are less than the MRHD on a body weight basis.

In a study in which pregnant rats received 6 weekly intramuscular injections of DYSPOSPORT (4.4, 11.1, 22.2, or 44 Units/kg) beginning on day 6 of gestation and continuing through parturition to weaning, an increase in stillbirths was observed at the highest dose tested, which was maternally toxic. The no-effect dose for pre- and post-natal developmental toxicity was 22.2 Units/kg (similar to the MRHD).

## **8.2 Lactation**

### Risk Summary

There are no data on the presence of DYSPOSPORT in human or animal milk, the effects on the breastfed infant, or the effects on milk production.

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for DYSPOSPORT and any potential adverse effects on the breastfed infant from DYSPOSPORT or from the underlying maternal condition.

## **8.3 Females and Males of Reproductive Potential**

### Infertility

In rats, DYSPOSPORT produced adverse effects on mating behavior and fertility [*see Nonclinical Toxicology (13.1)*].

## **8.4 Pediatric Use**

### Cervical Dystonia

Safety and effectiveness in pediatric patients have not been established [*see Boxed Warning and Warnings and Precautions (5.2)*].

### Glabellar Lines

DYSPOSPORT is not recommended for use in pediatric patients less than 18 years of age.

### Upper Limb Spasticity, Excluding Spasticity Caused by Cerebral Palsy

Safety and effectiveness have been established in pediatric patients 2 years of age and older [*see Warnings and Precautions (5.1), Adverse Reactions (6.1), and Clinical Studies (14.4)*]. The safety and effectiveness of DYSPOSPORT have been established by evidence from adequate and well-controlled studies of DYSPOSPORT in patients 2 years of age and older with upper limb spasticity. A pediatric assessment for DYSPOSPORT demonstrates that DYSPOSPORT is safe and effective in another pediatric population. However, DYSPOSPORT is not approved for such patient population due to marketing exclusivity for another botulinum toxin.

Safety and effectiveness in pediatric patients below the age of 2 years have not been established [*see Boxed Warning and Warnings and Precautions (5.2)*].

### Lower Limb Spasticity in Pediatric Patients

Safety and effectiveness have been established in pediatric patients 2 years of age and older [*see Warnings and Precautions (5.1), Adverse Reactions (6.1), and Clinical Studies (14.4)*]. The safety and effectiveness of DYSPOSPORT have been established by evidence from adequate and well-controlled studies of DYSPOSPORT in patients 2 years of age and older with lower limb spasticity. The safety

and effectiveness of DYSPORT injected into proximal muscles of the lower limb for the treatment of spasticity in pediatric patients has not been established [see *Boxed Warning, Warnings and Precautions (5.2) and Adverse Reactions (6.1)*].

Safety and effectiveness in pediatric patients below the age of 2 years have not been established [see *Boxed Warning, Warnings and Precautions (5.2)*].

#### *Juvenile Animal Data*

In a study in which juvenile rats received a single intramuscular injection of DYSPORT (1, 3, or 10 Units/animal) on postnatal day 21, decreased growth and bone length (injected and contralateral limbs), delayed sexual maturation, and decreased fertility were observed at the highest dose tested, which was associated with excessive toxicity during the first week after dosing.

In a study in which juvenile rats received weekly intramuscular injections of DYSPORT (0.1, 0.3, or 1.0 Units/animal) from postnatal day 21 to 13 weeks of age, decreases in bone mineral content in the injected limb, associated with atrophy of injected and adjacent muscles, were observed at the highest dose tested. No adverse effects were observed on neurobehavioral development. However, dose levels were not adjusted for growth of the pups. On a body weight basis, the doses at the end of the dosing period were approximately 15% of those at initiation of dosing. Therefore, the effects of DYSPORT throughout postnatal development were not adequately evaluated.

## **8.5 Geriatric Use**

### Cervical Dystonia

There were insufficient numbers of patients aged 65 years and over in the clinical studies to determine whether they respond differently than younger patients. In general, elderly patients should be observed to evaluate their tolerability of DYSPORT, due to the greater frequency of concomitant disease and other drug therapy [see *Dosage and Administration (2.3)*].

### Glabellar Lines

Of the total number of subjects in the placebo-controlled clinical studies of DYSPORT, 8 (1%) were 65 years and over. Efficacy was not observed in subjects aged 65 years and over [see *Clinical Studies (14.2)*]. For the entire safety database of geriatric subjects, although there was no increase in the incidence of eyelid ptosis, geriatric subjects did have an increase in the number of ocular adverse reactions compared to younger subjects (11% vs. 5%) [see *Dosage and Administration (2.4)*].

### Adult Spasticity

#### *Upper Limb Spasticity*

Of the total number of subjects in placebo-controlled clinical studies of DYSPORT, 30 percent were aged 65 years and over, while 8 percent were aged 75 years and over. No overall differences in safety or effectiveness were observed between these subjects and younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out.

#### *Lower Limb Spasticity*

Of the total number of subjects in placebo controlled clinical studies of DYSPORT, 18% (n = 115) were 65 and over, while 3% (n = 20) were 75 and over. Subjects aged 65 years and over who were treated with DYSPORT reported a greater percentage of adverse reactions as compared to younger subjects (46% versus 39%). Fall and asthenia were observed with greater frequency in older subjects, as compared to those younger (10% versus 6% and 4% versus 2%, respectively).

## **8.6 Ethnic Groups**

Exploratory analyses in trials for glabellar lines in African-American subjects with Fitzpatrick skin types IV, V, or VI and in Hispanic subjects suggested that response rates at Day 30 were comparable to and no worse than the overall population.

## **10 OVERDOSAGE**

Excessive doses of DYSPORT may be expected to produce neuromuscular weakness with a variety of symptoms. Respiratory support may be required where excessive doses cause paralysis of respiratory muscles. In the event of overdose, the patient should be medically monitored for symptoms of excessive muscle weakness or muscle paralysis [see *Boxed Warning and Warnings and Precautions (5.2)*]. Symptomatic treatment may be necessary.

Symptoms of overdose are likely not to be present immediately following injection. Should accidental injection or oral ingestion occur, the person should be medically supervised for several weeks for signs and symptoms of excessive muscle weakness or paralysis.

There is no significant information regarding overdose from clinical studies.

In the event of overdose, antitoxin raised against botulinum toxin is available from the Centers for Disease Control and Prevention (CDC) in Atlanta, GA. However, the antitoxin will not reverse any botulinum toxin-induced effects already apparent by the time of

antitoxin administration. In the event of suspected or actual cases of botulinum toxin poisoning, please contact your local or state Health Department to process a request for antitoxin through the CDC. If you do not receive a response within 30 minutes, please contact the CDC directly at 770-488-7100. More information can be obtained at <https://www.cdc.gov/laboratory/drugservice/index.html>.

## 11 DESCRIPTION

Botulinum toxin type A, the active ingredient in DYSPORT, is a purified neurotoxin type A complex produced by fermentation of the bacterium *Clostridium botulinum* type A, Hall Strain. It is purified from the culture supernatant by a series of precipitation, dialysis, and chromatography steps. The neurotoxin complex is composed of the neurotoxin, hemagglutinin proteins and non-toxin non-hemagglutinin protein.

DYSPORT® (abobotulinumtoxinA) for injection is a sterile, lyophilized powder supplied in a single-dose vial for reconstitution intended for intramuscular injection. Each vial contains 300 Units or 500 Units of lyophilized abobotulinumtoxinA, human serum albumin (125 mcg) and lactose (2.5 mg). DYSPORT may contain trace amounts of cow's milk proteins [see *Contraindications (4) and Warnings and Precautions (5.3)*].

The primary release procedure for DYSPORT uses a cell-based potency assay to determine the potency relative to a reference standard. The assay and reference material are specific to DYSPORT. One unit of DYSPORT corresponds to the calculated median lethal intraperitoneal dose (LD<sub>50</sub>) in mice. Due to specific details of the assay system, such as vehicle, dilution scheme and laboratory protocols, Units of biological activity of DYSPORT cannot be converted into Units of any other botulinum toxin or any toxin assessed with any other specific assay method.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

DYSPORT inhibits release of the neurotransmitter, acetylcholine, from peripheral cholinergic nerve endings. Toxin activity occurs in the following sequence: Toxin heavy chain mediated binding to specific surface receptors on nerve endings, internalization of the toxin by receptor mediated endocytosis, pH-induced translocation of the toxin light chain to the cell cytosol and cleavage of SNAP25 leading to intracellular blockage of neurotransmitter exocytosis into the neuromuscular junction. This accounts for the therapeutic utility of the toxin in diseases characterized by excessive efferent activity in motor nerves.

Recovery of transmission occurs gradually as the neuromuscular junction recovers from SNAP25 cleavage and as new nerve endings are formed.

### 12.2 Pharmacodynamics

The primary pharmacodynamic effect of DYSPORT is due to chemical denervation of the treated muscle resulting in a measurable decrease of the compound muscle action potential, causing a localized reduction of muscle activity.

### 12.3 Pharmacokinetics

Using currently available analytical technology, it is not possible to detect DYSPORT in the peripheral blood following intramuscular injection at the recommended doses.

## 13 NONCLINICAL TOXICOLOGY

### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

#### Carcinogenesis

Studies to evaluate the carcinogenic potential of DYSPORT have not been conducted.

#### Mutagenesis

Genotoxicity studies have not been conducted for DYSPORT.

#### Impairment of Fertility

In a fertility and early embryonic development study in rats in which either males (2.9, 7.2, 14.5 or 29 Units/kg) or females (7.4, 19.7, 39.4 or 78.8 Units/kg) received weekly intramuscular injections prior to and after mating, dose-related increases in pre-implantation loss and reduced numbers of corpora lutea were noted in treated females. Failure to mate was observed in males that received the high dose. The no-effect dose for effects on fertility was 7.4 Units/kg in females and 14.5 Units/kg in males (approximately one-half and equal to, respectively, the maximum recommended human dose of 1000 Units on a body weight basis).

## 14 CLINICAL STUDIES

### 14.1 Cervical Dystonia

The efficacy of DYSPORT was evaluated in two randomized, double-blind, placebo-controlled, single-dose, parallel-group studies in treatment-naïve cervical dystonia patients. The principal analyses from these trials provide the primary demonstration of efficacy involving 252 patients (121 on DYSPORT, 131 on placebo) with 36% male and 64% female. Ninety-nine percent of the patients were Caucasian.

In both placebo-controlled studies (Study 1 and Study 2), a dose of 500 Units of DYSPORT was given by intramuscular injection divided among two to four affected muscles. These studies were followed by long-term open-label extensions that allowed titration in 250 Unit steps to doses in a range of 250 to 1000 Units, after the initial dose of 500 Units. In the extension studies, re-treatment was determined by clinical need after a minimum of 12 weeks. The median time to re-treatment was 14 weeks and 18 weeks for the 75<sup>th</sup> percentile.

The primary assessment of efficacy was based on the total Toronto Western Spasmodic Torticollis Rating Scale (TWSTRS) change from baseline at Week 4 for both studies. The scale evaluates the severity of dystonia, patient-perceived disability from dystonia, and pain. The adjusted mean change from baseline in the TWSTRS total score was statistically significantly greater for the DYSPORT group than the placebo group at Week 4 in both studies (see Table 14).

**Table 14: TWSTRS Total Score Efficacy Outcome from the Phase 3 Cervical Dystonia Studies Intent to Treat Population**

	Study 1		Study 2	
	DYSPORT 500 Units N=55	Placebo N=61	DYSPORT 500 Units N=37	Placebo N=43
<b>Baseline (Week 0)</b> Mean (SD)	43.8 (8.0)	45.8 (8.9)	45.1 (8.7)	46.2 (9.4)
<b>Week 4</b> Mean (SD)	30.0 (12.7)	40.2 (11.8)	35.2 (13.8)	42.4 (12.2)
Change from Baseline*	-15.6 (2.0)	-6.7 (2.0)	-9.6 (2.0)	-3.7 (1.8)
Treatment difference 95% confidence interval	-8.9 <sup>‡</sup> [-12.9 to -4.7]		-5.9 <sup>‡</sup> [-10.6 to -1.3]	
<b>Week 8</b> Mean (SD)	29.3 (11.0)	39.6 (13.5)		
Change from Baseline*	-14.7 (2.0)	-5.9 (2.0)		
Treatment difference 95% confidence interval	-8.8 <sup>‡</sup> [-12.9 to -4.7]			

\*Change from baseline is expressed as adjusted least squares mean (SE)

<sup>‡</sup>Significant at  $p$ -value < 0.05

Analyses by gender, weight, geographic region, underlying pain, cervical dystonia severity at baseline and history of treatment with botulinum toxin did not show any meaningful differences between groups.

Table 15 indicates the average DYSPORT dose, and percentage of total dose, injected into specific muscles in the pivotal clinical trials.

**Table 15: DYSPORT 500 Units starting dose (units and % of the total dose) by Unilateral Muscle Injected During Double-blind Pivotal Phase 3 studies 2 and 1 Combined**

Number of patients injected per muscle*		DYSPORT Dose Injected		Percentage of the total DYSPORT Dose Injected	
		Median [DYSPORT Units] (min, max)	75th percentile [DYSPORT Units]	Median [%] (min, max)	75th percentile [%]
Sternocleidomastoid	90	125 Units (50, 350)	150 Units	26.5 % (10, 70)	30.0 %
Splenius capitis	85	200 Units (75, 450)	250 Units	40.0 % (15, 90)	50.0 %
Trapezius	50	102.6 Units (50, 300)	150 Units	20.6 % (10, 60)	30.0 %
Levator scapulae	35	105.3 Units (50, 200)	125 Units	21.1 % (10, 40)	25.0 %
Scalenus (medius and anterior)	26	115.5 Units (50, 300)	150 Units	23.1 % (10, 60)	30.0 %
Semispinalis capitis	21	131.6 Units (50, 250)	175 Units	29.4 % (10, 50)	35.0 %
Longissimus	3	150 Units (100, 200)	200 Units	30.0 % (20, 40)	40.0 %

\*Total number of patients in combined studies 2 and 1 who received initial treatment = 121.

## 14.2 Glabellar Lines

Three double-blind, randomized, placebo-controlled, clinical studies evaluated the efficacy of DYSPORT for use in the temporary improvement of the appearance of moderate to severe glabellar lines. These three studies enrolled healthy adults (ages 19-75) with glabellar lines of at least moderate severity at maximum frown. Subjects were excluded if they had marked ptosis, deep dermal scarring, or a substantial inability to lessen glabellar lines, even by physically spreading them apart. The subjects in these studies received either DYSPORT or placebo. The total dose was delivered in equally divided aliquots to specified injection sites (see Figure 1).

Investigators and subjects assessed efficacy at maximum frown by using a 4-point scale (none, mild, moderate, severe).

Overall treatment success was defined as post-treatment glabellar line severity of none or mild with at least 2 grade improvement from baseline for the combined investigator and subject assessments (composite assessment) on Day 30 (see Table 16). Additional endpoints for each of the studies were post-treatment glabellar line severity of none or mild with at least a 1 grade improvement from baseline for the separate investigator and subject assessments on Day 30.

After completion of the randomized studies, subjects were offered participation in a two-year, open-label re-treatment study to assess the safety of multiple treatments.

**Table 16: Treatment Success at Day 30 (None or Mild with at least 2 Grade Improvement from Baseline at Maximum Frown for the combined Investigator and Subject Assessments (Composite))**

Study	2 Grade Improvement	
	DYSPORT n/N (%)	Placebo n/N (%)
GL-1	58/105 (55%)	0/53 (0%)
GL-2	37/71 (52%)	0/71 (0%)
GL-3	120/200 (60%)	0/100 (0%)

Treatment with DYSPORT reduced the severity of glabellar lines for up to four months.

### Study GL-1

Study GL-1 was a single-dose, double-blind, multicenter, randomized, placebo-controlled study in which 158 previously untreated subjects received either placebo or 50 Units of DYSPORT, administered in five aliquots of 10 Units (see Figure 1). Subjects were followed for 180 days. The mean age was 43 years; most of the subjects were women (85%), and predominantly Caucasian (49%) or

Hispanic (47%). At Day 30, 55% of DYSPORT-treated subjects achieved treatment success: a composite 2 grade improvement of glabellar line severity at maximum frown (see Table 16).

In study GL-1, the reduction of glabellar line severity at maximum frown was greater at Day 30 in the DYSPORT group compared to the placebo group as assessed by both Investigators and subjects (see Table 17).

**Table 17: GL-1: Investigators' and Subjects' Assessment of Glabellar Line Severity at Maximum Frown Using a 4-point Scale (% and Number of Subjects with Severity of None or Mild)**

Day	Investigators' Assessment		Subjects' Assessment	
	DYSPORT N=105	Placebo N=53	DYSPORT N=105	Placebo N=53
14	90% 95	17% 9	77% 81	9% 5
30	88% 92	4% 2	74% 78	9% 5
60	64% 67	2% 1	60% 63	6% 3
90	43% 45	6% 3	36% 38	6% 3
120	23% 24	4% 2	19% 20	6% 3
150	9% 9	2% 1	8% 8	4% 2
180	6% 6	0% 0	7% 7	8% 4

#### Study GL-2

Study GL-2 was a repeat-dose, double-blind, multicenter, placebo-controlled, randomized study. The study was initiated with two or three open-label treatment cycles of 50 Units of DYSPORT administered in five aliquots of 10 Units DYSPORT (see Figure 1). After the open-label treatments, subjects were randomized to receive either placebo or 50 Units of DYSPORT. Subjects could have received up to four treatments through the course of the study. Efficacy was assessed in the final randomized treatment cycle. The study enrolled 311 subjects into the first treatment cycle and 142 subjects were randomized into the final treatment cycle. Overall, the mean age was 47 years; most of the subjects were women (86%) and predominantly Caucasian (80%).

At Day 30, 52% of DYSPORT-treated subjects achieved treatment success: a composite 2 grade improvement of glabellar line severity at maximum frown (see Table 16).

The proportion of responders in the final treatment cycle was comparable to the proportion of responders in all prior treatment cycles.

After the final repeat treatment with DYSPORT, the reduction of glabellar line severity at maximum frown was greater at Day 30 in the DYSPORT group compared to the placebo group as assessed by both Investigators and subjects (see Table 18).

**Table 18: GL-2: Investigators' and Subjects' Assessments of Glabellar Line Severity at Maximum Frown Using a 4-point Scale (% and Number of Subjects with Severity of None or Mild)**

Day	Investigators' Assessment		Subjects' Assessment	
	DYSPORT N=71	Placebo N=71	DYSPORT N=71	Placebo N=71
30	85% 60	4% 3	79% 56	1% 1

#### Study GL-3

Study GL-3 was a single-dose, double-blind, multicenter, randomized, placebo-controlled study in which 300 previously untreated subjects received either placebo or 50 Units of DYSPORT, administered in five aliquots of 10 Units (see Figure 1). Subjects were

followed for 150 days. The mean age was 44 years; most of the subjects were women (87%), and predominantly Caucasian (75%) or Hispanic (18%).

At Day 30, 60% of DYSPORT-treated subjects achieved treatment success: a composite 2 grade improvement of glabellar line severity at maximum frown (see Table 16).

In study GL-3, the reduction of glabellar line severity at maximum frown was greater at Day 30 in the DYSPORT group compared to the placebo group as assessed by both Investigators and subjects (see Table 19).

**Table 19. GL-3: Investigators' and Subjects' Assessment of Glabellar Line Severity at Maximum Frown Using a 4-point Scale (% and Number of Subjects with Severity of None or Mild)**

Day	Investigators' Assessment		Subjects' Assessment	
	DYSPORT N=200	Placebo N=100	DYSPORT N=200	Placebo N=100
14	83% 166	5% 5	83% 165	2% 2
30	86% 171	0% 0	82% 163	2% 2
60	75% 150	1% 1	65% 130	4% 4
90	51% 102	1% 1	46% 91	2% 2
120	29% 58	1% 1	31% 61	3% 3
150	16% 32	1% 1	16% 31	3% 3

#### Geriatric Subjects

In GL1, GL2, and GL3, there were 8 subjects aged 65 and older who were randomized to DYSPORT 50 Units in 5 equal aliquots of 10 Units (4) or placebo (4). None of the geriatric DYSPORT subjects were a treatment success at maximum frown at Day 30.

### **14.3 Spasticity in Adults**

#### Upper Limb Spasticity

The efficacy and safety of DYSPORT for the treatment of upper limb spasticity in adults was evaluated in a randomized, multicenter, double-blind, placebo-controlled study that included 238 patients (159 DYSPORT and 79 placebo) with upper limb spasticity (Modified Ashworth Scale (MAS) score  $\geq 2$  in the primary targeted muscle group for toxin-naive patients or MAS score  $\geq 3$  in the primary targeted muscle group for toxin non-naive patients at least 4 months after the last botulinum toxin injection, of any serotype) who were at least 6 months post-stroke or post-traumatic brain injury. The median age of the patients in this study was 55 years (range 18 to 78 years), 64% were male, and 86% were Caucasian.

DYSPORT 500 Units (N=80), DYSPORT 1000 Units (N=79), or placebo (N=79) was injected intramuscularly into the affected upper limb muscles. After injection of the primary targeted muscle groups (PTMG), the remainder of the dose was injected into at least two additional upper limb muscles determined by the patient's individual presentation. Table 20 provides the mean and range of DYSPORT doses injected and the number of injections into specific muscles of the upper limb.

**Table 20: DYSPORT Dose Injected and Number of Injections per Muscle in Adults with Upper Limb Spasticity**

Muscle	DYSPORT Treatment Group	Number of Patients	Mean DYSPORT Units Injected (Min, Max)	Number of Injection Sites Median, [Q1 ; Q3]
Flexor digitorum profundus (FDP)*	500 U	54	93.5 Units (50 to 100)	1, [1 ; 2]
	1000 U	65	195.5 Units (100 to 300)	2, [1 ; 2]
Flexor digitorum superficialis (FDS)*	500 U	63	95.4 Units (50 to 100)	2, [1 ; 2]
	1000 U	73	196.8 Units (100 to 300)	2, [1 ; 2]
Flexor carpi radialis (FCR)*	500 U	57	92.2 Units (25 to 100)	1, [1 ; 2]
	1000 U	57	178.1 Units (80 to 300)	1, [1 ; 2]
Flexor carpi ulnaris (FCU)*	500 U	47	89.9 Units (25 to 180)	1, [1 ; 2]
	1000 U	49	171.2 Units (80 to 200)	1, [1 ; 2]
Brachialis*	500 U	60	148.5 Units (50 to 200)	2, [1 ; 2]
	1000 U	43	321.4 Units (100 to 400)	2, [2 ; 2]
Brachioradialis*	500 U	42	88.3 Units (50 to 200)	1, [1 ; 2]
	1000 U	28	172.1 Units (50 to 200)	1, [1 ; 2]
Biceps Brachii (BB)	500 U	28	106.4 Units (50 to 200)	2, [1 ; 2]
	1000 U	19	207.4 Units (100 to 400)	2, [1 ; 2]
Pronator Teres	500 U	14	81.8 Units (45 to 200)	1, [1 ; 1]
	1000 U	30	157.3 Units (80 to 200)	1, [1 ; 1]

\*Primary Targeted Muscle Group

The co-primary efficacy variables were muscle tone assessed by the MAS at the primary targeted muscle group at Week 4 and the Physician Global Assessment (PGA; ranges from -4 = markedly worse to +4= markedly improved) at Week 4 (see Table 21).

**Table 21: Primary Endpoints (PTMG MAS and PGA) and MAS by Muscle Group at Week 4 in Adults with Upper Limb Spasticity**

	Placebo (N=79)	DYSPORT	
		(500 Units) (N=80)	(1000 Units) (N=79)
LS Mean Change from Baseline in PTMG Muscle Tone on the MAS	-0.3	-1.2*	-1.4*
LS Mean PGA of Response to Treatment	0.7	1.4*	1.8*
LS Mean Change from Baseline in Wrist Flexor Muscle Tone on the MAS	-0.3 (n=54)	-1.4 (n=57)	-1.6 (n=58)
LS Mean Change from Baseline in Finger Flexor Muscle Tone on the MAS	-0.3 (n=70)	-0.9 (n=66)	-1.2 (n=73)
LS Mean Change from Baseline in Elbow Flexor Muscle Tone on the MAS	-0.3 (n=56)	-1.0 (n=61)	-1.2 (n=48)

LS= Least Square;

\*p≤0.05

#### Lower Limb Spasticity

The efficacy of DYSPORT for the treatment of lower limb spasticity was evaluated in a randomized, multicenter, double-blind, placebo-controlled study that included 381 patients (253 DYSPORT and 128 placebo). Patients had lower limb spasticity (Modified Ashworth Scale (MAS) score ≥2 in the affected ankle joint for toxin-naïve patients, or MAS score ≥3 in the affected ankle joint for toxin non-naïve patients) and were at least 6 months post-stroke or post-traumatic brain injury.

Table 22 provides the median DYSPORT doses injected and the number of injections into specific muscles of the lower limb as reported in the double-blind study. In the study, the gastrocnemius and soleus muscles, and at least one additional lower limb muscle were injected, according to the clinical presentation.

**Table 22: DYSPORT Dose Injected and Number of Injections per Muscle in the Lower Limb - Median for the 1000 Unit and 1500 Unit Dose Groups**

Injected Muscle	DYSPORT Units Injected	Number of Injection Sites
Gastrocnemius <i>Lateral</i>	100 Units to 150 Units	1
<i>Medial</i>	100 Units to 150 Units	1
Soleus	333 Units to 500 Units	3
Tibialis posterior	200 Units to 300 Units	2
Flexor digitorum longus	133 Units to 200 Units	1 to 2
Flexor hallucis longus	67 Units to 200 Units	1

The primary efficacy variable was muscle tone assessed by the MAS at the ankle joint at Week 4. The first secondary endpoint was the Physician Global Assessment at Week 4 (see Table 23).

**Table 23: Primary Endpoint Change in MAS and the First Secondary Endpoint PGA at Week 4 in Adults with Lower Limb Spasticity**

LS Mean Change from Baseline on the Modified Ashworth Scale	DYSPORT 1000 Units (N=125)	DYSPORT 1500 Units (N=128)	Placebo (N=128)
Week 4	-0.6	-0.8*	-0.5
LS Mean Physician Global Assessment Score Investigator			
Week 4	0.9	0.9	0.7

\*p<0.05

#### 14.4 Spasticity in Pediatric Patients

##### Upper Limb Spasticity in Pediatric Patients

The efficacy of DYSPORT for the treatment of upper limb spasticity in pediatric patients 2 years of age and older was evaluated in a double-blind, low-dose controlled, multicenter study (NCT02106351). A total of 208 toxin naive or non-naive (66% had prior treatment with a botulinum toxin) patients weighing at least 10 kgs, with a baseline Modified Ashworth Score (MAS) of grade 2 or greater (99% patients) at the primary targeted muscle groups (PTMG), were enrolled in the modified Intention to Treat population (mITT). Patients received DYSPORT 16 Units/kg (n=70), DYSPORT 8 Units/kg (n=69), or DYSPORT 2 Units/kg (n=69) injected into the upper limb. The elbow flexors and wrist flexors respectively were the PTMG in 57% and in 43% of patients. The median age of the patients in this study was 9 years (range 2 to 17 years; 57% were between 2 and 9 years of age); 60% of patients were male, and 75% were White.

The primary efficacy endpoint was the mean change from baseline in MAS in the PTMG at Week 6 (see Table 24). The secondary efficacy endpoint was the mean Physician Global Assessment (PGA) score assessed at Week 6 (Table 25). Although PGA scores numerically favored DYSPORT treatment over the low-dose control, the difference was not statistically significant.

**Table 24: Modified Ashworth Scale Score in the PTMG Change from Baseline at Week 6 in Pediatric Patients with Upper Limb Spasticity (mITT Population)**

	Control Group	Treatment Groups	
	DYSPORT 2 U/kg (N=69)	DYSPORT 8 U/kg (N=69)	DYSPORT 16 U/kg (N=70)
<b>Baseline</b>			
Mean (SD)	3.1 (0.3)	3.1 (0.3)	3.1 (0.5)
<b>Week 6</b>			
LS <sup>a</sup> mean change from baseline in PTMG <sup>b</sup> on MAS	-1.6	-2.0	-2.3
Difference from control in LS <sup>a</sup> means		-0.4	-0.7
p-value <sup>c</sup>		0.0118 <sup>d</sup>	<0.0001
<b>Week 16</b>			
LS <sup>a</sup> mean change from baseline in PTMG <sup>b</sup> on MAS	-0.9	-1.2	-1.5
Difference from control in LS <sup>a</sup> means		-0.3 <sup>d</sup>	-0.6 <sup>d</sup>

<sup>a</sup> LS=Least Square

<sup>b</sup> PTMG=Primary Targeted Muscle Group

<sup>c</sup> p-value is derived from ANCOVA on ranked MAS score change from baseline with treatment, baseline score, age range at baseline, prior botulinum toxin treatment status at baseline, and center as explanatory variables

<sup>d</sup> Nominal p-value <0.05

**Table 25: Physician Global Assessment of Treatment Response at Week 6 in Pediatric Patients with Upper Limb Spasticity (mITT Population)**

	Control Group	Treatment Groups	
	Dysport 2 U/kg (N=68)	Dysport 8 U/kg (N=69)	Dysport 16 U/kg (N=70)
<b>Week 6</b>			
Mean score (SD)	1.7 (0.9)	2.0 (0.9)	2.0 (0.9)
LS <sup>a</sup> mean in PGA	1.8	2.0	2.0
Difference from control in LS <sup>a</sup> means		0.2	0.2
p-value <sup>b</sup>		0.2043	0.1880
<b>Week 16</b>			
Mean score (SD)	1.7 (1.0)	1.6 (1.1)	1.9 (1.2)
LS <sup>a</sup> mean in PGA	1.8	1.7	1.9
Difference from control in LS <sup>a</sup> means		-0.1	0.1
p-value <sup>b</sup>		0.7001	0.4041

<sup>a</sup> LS=Least Square

<sup>b</sup> p-value is derived from ANOVA on ranked PGA score with treatment, age range at baseline, prior botulinum toxin treatment status at baseline, and center as explanatory variables

#### Lower Limb Spasticity in Pediatric Patients

The efficacy of DYSPORT was evaluated in a double-blind, placebo-controlled, multicenter study in patients 2 to 17 years of age treated for lower limb spasticity because of cerebral palsy causing dynamic equinus foot deformity. A total of 235 (158 DYSPORT and 77 placebo) toxin-naïve or non-naïve patients with a Modified Ashworth Score (MAS) of grade 2 or greater at the ankle plantar flexors were enrolled to receive DYSPORT 10 Units/kg/leg (n=79), DYSPORT 15 Units/kg/leg (n=79) or placebo (n=77) injected into the gastrocnemius and soleus muscles (see Table 27). Forty-one percent of patients (n=66) were treated bilaterally and received a total lower limb DYSPORT dose of either 20 Units/kg (n=37) or 30 Units/kg (n=29). The median age of the patients in this study was 5 years (range 2 to 17 years); 60% of patients were male, and 73% were Caucasian.

The primary efficacy endpoint was the mean change from baseline in MAS in ankle plantar flexor at Week 4; a co-primary endpoint was the mean Physician's Global Assessment (PGA) score at Week 4 (see Table 26).

**Table 26: MAS and PGA Change from Baseline at Week 4 in Pediatric Patients with Lower Limb Spasticity (ITT Population)**

		<b>Placebo (N=77)</b>	<b>DYSPO RT 10 Units/kg/leg (N=79)</b>	<b>DYSPO RT 15 Units/kg/leg (N=79)</b>
LS Mean Change from Baseline in Ankle plantar flexor Muscle Tone on the MAS	Week 4	-0.5	-0.9*	-1.0*
	Week 12	-0.5	-0.8*	-1.0*
LS Mean PGA of Response to Treatment	Week 4	0.7	1.5*	1.5*
	Week 12	0.4	0.8*	1.0*

LS=Least Square

\*p<0.05

## 16 HOW SUPPLIED/STORAGE AND HANDLING

DYSPO<sup>RT</sup>® (abobotulinumtoxinA) for injection is a sterile, lyophilized powder supplied in a single-dose, glass vial. Unopened vials of DYSPO<sup>RT</sup> must be stored refrigerated between 2°C to 8°C (36°F to 46°F). Protect from light.

Do not use after the expiration date on the vial. All vials, including expired vials, or equipment used with DYSPO<sup>RT</sup> should be disposed of carefully as is done with all medical waste.

DYSPO<sup>RT</sup> contains a unique hologram on the carton. If you do not see the hologram, do not use the product. Instead contact 855-463-5127.

### Cervical Dystonia, Spasticity in Adults and Pediatric Patients

#### 500 Unit Vial

Each vial contains 500 Units of freeze-dried abobotulinumtoxinA.

Box containing 1 vial—NDC 15054-0500-1

Box containing 2 vials—NDC 15054-0500-2

#### 300 Unit Vial

Each vial contains 300 Units of freeze-dried abobotulinumtoxinA.

Box containing 1 vial—NDC 15054-0530-6

### Glabellar Lines

Each vial contains 300 Units of freeze-dried abobotulinumtoxinA.

Box containing 1 vial— NDC 0299-5962-30

## 17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide).

### Swallowing, Speaking, or Breathing Difficulties, or Other Unusual Symptoms

Advise patients to inform their doctor or pharmacist if they develop any unusual symptoms (including difficulty with swallowing, speaking or breathing), or if any known symptom persists or worsens [see *Warnings and Precautions (5.1, 5.4)*].

### Dry Eye with the Treatment of Glabellar Lines

Inform patients that DYSPO<sup>RT</sup> injection may cause eye dryness [see *Warnings and Precautions (5.6)*]. Advise patients to report symptoms of eye dryness (e.g., eye pain, eye irritation, photosensitivity, or changes in vision) to their doctor.

### Ability to Operate Machinery or Vehicles

Inform patients that if loss of strength, muscle weakness, blurred vision or drooping eyelids occur, they should avoid driving a car or engaging in other potentially hazardous activities.

Manufactured by:  
Ipsen Biopharm Ltd.  
Wrexham, LL13 9UF, UK  
U.S. License No. 1787

Distributed by:  
Ipsen Biopharmaceuticals, Inc.  
Cambridge, MA 02142  
and  
Galderma Laboratories, L.P.  
Fort Worth, TX 76177 USA

DYSPORT is a registered trademark of Ipsen Biopharm Limited.  
© 2019. All rights reserved.

**MEDICATION GUIDE**  
**DYSPO<sup>®</sup>RT (DIS-port)**  
**(abobotulinumtoxinA)**  
**for Injection**

**What is the most important information I should know about DYSPO<sup>®</sup>RT?**

**DYSPO<sup>®</sup>RT may cause serious side effects that can be life threatening including:**

- **Problems breathing or swallowing**
- **Spread of toxin effects**

**These problems can happen within hours, or days to weeks after an injection of DYSPO<sup>®</sup>RT. Call your doctor or get medical help right away if you have any of these problems after treatment with DYSPO<sup>®</sup>RT:**

**1. Problems swallowing, speaking, or breathing.** These problems can happen within hours, or days to weeks after an injection of DYSPO<sup>®</sup>RT usually because the muscles that you use to breathe and swallow can become weak after the injection. Death can happen as a complication if you have severe problems with swallowing or breathing after treatment with DYSPO<sup>®</sup>RT.

- People with certain breathing problems may need to use muscles in their neck to help them breathe. These patients may be at greater risk for serious breathing problems with DYSPO<sup>®</sup>RT.
- Swallowing problems may last for several weeks. People who cannot swallow well may need a feeding tube to receive food and water. If swallowing problems are severe, food or liquids may go into your lungs. People who already have swallowing or breathing problems before receiving DYSPO<sup>®</sup>RT have the highest risk of getting these problems.

**2. Spread of toxin effects.** In some cases, the effect of botulinum toxin may affect areas of the body away from the injection site and cause symptoms of a serious condition called botulism. The symptoms of botulism include:

- loss of strength and muscle weakness all over the body
- blurred vision and drooping eyelids
- trouble saying words clearly (dysarthria)
- trouble breathing
- double vision
- hoarseness or change or loss of voice (dysphonia)
- loss of bladder control
- trouble swallowing

**These symptoms can happen within hours, or days to weeks after you receive an injection of DYSPO<sup>®</sup>RT. These problems could make it unsafe for you to drive a car or do other dangerous activities. See “What should I avoid while receiving DYSPO<sup>®</sup>RT?”**

**What is DYSPO<sup>®</sup>RT?**

DYSPO<sup>®</sup>RT is a prescription medicine that is injected into muscles and used:

- to treat cervical dystonia (CD) in adults
- to improve the look of moderate to severe frown lines between the eyebrows (glabellar lines) in adults younger than 65 years of age for a short period of time (temporary)
- to treat increased muscle stiffness in adults with spasticity
- to treat increased muscle stiffness in children 2 years of age and older with upper limb spasticity, excluding spasticity caused by cerebral palsy
- to treat increased muscle stiffness in children 2 years of age and older with lower limb spasticity

It is not known whether DYSPO<sup>®</sup>RT is safe or effective in people younger than:

- 18 years of age for the treatment of cervical dystonia
- 18 years of age for the treatment of glabellar lines
- 2 years of age for the treatment of spasticity

It is not known whether DYSPO<sup>®</sup>RT is safe or effective for the treatment of other types of muscle spasms.

It is not known whether DYSPO<sup>®</sup>RT is safe or effective for the treatment of other wrinkles.

### **Who should not take DYSPORT?**

#### **Do not take DYSPORT if you:**

- are allergic to DYSPORT or any of the ingredients in DYSPORT. See the end of this Medication Guide for a list of ingredients in DYSPORT
- are allergic to cow's milk protein
- had an allergic reaction to any other botulinum toxin product such as Myobloc® (rimabotulinumtoxinB), Botox® (onabotulinumtoxinA), or Xeomin® (incobotulinumtoxinA)
- have a skin infection at the planned injection site

### **What should I tell my doctor before taking DYSPORT?**

#### **Tell your doctor about all your medical conditions, including if you:**

- have a disease that affects your muscles and nerves (such as amyotrophic lateral sclerosis [ALS or Lou Gehrig's disease], myasthenia gravis or Lambert-Eaton syndrome). See "What is the most important information I should know about DYSPORT?"
- have allergies to any botulinum toxin product
- had any side effect from any botulinum toxin product in the past
- have or have had a breathing problem, such as asthma or emphysema
- have or have had swallowing problems
- have or have had bleeding problems
- have diabetes
- have or have had a slow heart beat or other problem with your heart rate or rhythm
- have plans to have surgery
- had surgery on your face
- have weakness of your forehead muscles (such as trouble raising your eyebrows)
- have drooping eyelids
- experienced dry eye with previous use of botulinum toxin products
- have any other change in the way your face normally looks
- are pregnant or plan to become pregnant. It is not known if DYSPORT can harm your unborn baby
- are breast-feeding or planning to breast-feed. It is not known if DYSPORT passes into breast milk

**Tell your doctor about all the medicines you take**, including prescription and over-the-counter medicines, vitamins and herbal products. Using DYSPORT with certain other medicines may cause serious side effects. **Do not start any new medicines until you have told your doctor that you have received DYSPORT in the past.**

#### **Especially tell your doctor if you:**

- have received any other botulinum toxin product in the last four months
- have received injections of botulinum toxin, such as Myobloc® (rimabotulinumtoxinB), Botox® (onabotulinumtoxinA) or Xeomin® (incobotulinumtoxinA) in the past; be sure your doctor knows exactly which product you received
- have recently received an antibiotic by injection
- take muscle relaxants
- take an allergy or cold medicine
- take a sleep medicine

#### **Ask your doctor if you are not sure if your medicine is one that is listed above.**

Know the medicines you take. Keep a list of your medicines with you to show your doctor and pharmacist each time you get a new medicine.

### **How will I receive DYSPORT?**

- DYSPORT is an injection that your doctor will give you
- DYSPORT is injected into the affected muscles
- If you are an adult, your doctor may give you another dose of DYSPORT after 12 weeks or longer, if it is needed
- If you are an adult being treated for CD or spasticity or you are a child (2 to 17 years of age) being treated for spasticity, your doctor may change your dose of DYSPORT until you and your doctor find the best dose for you. Children should not be retreated sooner than every 12-16 weeks
- The dose of DYSPORT is not the same as the dose of any other botulinum toxin product

### **What should I avoid while receiving DYSPORT?**

DYSPORT may cause loss of strength or general muscle weakness, blurred vision, or drooping eyelids within hours to weeks of taking DYSPORT. **If this happens, do not drive a car, operate machinery, or do other dangerous activities. See "What is the most important information I should know about DYSPORT?"**

**What are the possible side effects of DYSPORT?**

**DYSPORT can cause serious side effects. See "What is the most important information I should know about DYSPORT?"**

**The most common side effects of DYSPORT in people with cervical dystonia include:**

- muscle weakness
- dry mouth
- feeling of tiredness
- muscle pain
- problems speaking
- eye problems
- difficulty swallowing
- injection site pain or discomfort
- headache

**The most common side effects of DYSPORT in people with glabellar lines include:**

- stuffy or runny nose and sore throat
- injection site pain
- upper respiratory infection
- blood in urine
- headache
- injection site reaction
- swelling of eyelids
- drooping eyelids
- sinus infection
- nausea

**The most common side effect of DYSPORT in adults with upper limb spasticity include:**

- muscle weakness

**The most common side effects of DYSPORT in adults with lower limb spasticity include:**

- muscle weakness
- pain in your arms or legs
- fall

**The most common side effects of DYSPORT in children (2 to 17 years of age) with upper limb spasticity include:**

- upper respiratory tract infection
- sore throat

**The most common side effects of DYSPORT in children (2 to 17 years of age) with lower limb spasticity include:**

- stuffy or runny nose and sore throat
- cough
- fever

**Tell your doctor if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of DYSPORT. For more information, ask your doctor or pharmacist.**

Tell your doctor if you have dry eye or changes in vision following use of DYSPORT.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**General information about DYSPORT:**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide.

This Medication Guide summarizes the most important information about DYSPORT. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about DYSPORT that is written for healthcare professionals.

**What are the ingredients in DYSPORT?**

Active ingredient: (botulinum toxin Type A)

Inactive ingredients: human albumin and lactose. DYSPORT may contain cow's milk protein.

Distributed by: Ipsen Biopharmaceuticals, Inc. Cambridge, MA 02142 and Galderma Laboratories, L.P. Fort Worth, TX 76177; Manufactured by: Ipsen Biopharm Ltd., Wrexham, LL13 9UF, UK U.S. License No. 1787

For more information about DYSPORT, call 855-463-5127 or go to [www.dysport.com](http://www.dysport.com) or [www.DysportUSA.com](http://www.DysportUSA.com).

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Revised 9/2019

DYSPORT is a registered trademark of Ipsen Biopharm Limited. Botox, Xeomin and Myobloc are registered trademarks of their respective owners.

© 2019. All rights reserved.