

	
<p>Drug Facts</p> <p>Active ingredient (in each capsule) Purposes</p> <p>Aspirin (NSAID)* 325 mg Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug</p> <p>Uses</p> <ul style="list-style-type: none"> ■ for the temporary relief of minor aches and pains associated with: <ul style="list-style-type: none"> ■ headache ■ backache ■ muscular aches ■ a cold ■ toothache ■ minor pain of arthritis ■ premenstrual and menstrual cramps ■ temporarily reduces fever ■ ask your doctor about other uses for aspirin capsules <p>Warnings</p> <p>Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p>Allergy alert: Aspirin may cause a severe allergic reaction which may include:</p> <ul style="list-style-type: none"> ■ hives ■ facial swelling ■ shock ■ asthma (wheezing) <p>This product contains soy.</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:</p> <ul style="list-style-type: none"> ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed <p>Do not use if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer.</p> <p>Ask doctor before use if</p> <ul style="list-style-type: none"> ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma <p>Ask doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> ■ taking a prescription drug for diabetes, gout or arthritis 	<p>Drug Facts (continued)</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ an allergic reaction occurs. Seek medical help right away, if you are experiencing any of the following signs of stomach bleeding: <ul style="list-style-type: none"> ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that doesn't get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts for more than 3 days ■ ringing in the ears or loss of hearing occurs ■ redness or swelling is present in the painful area ■ any new symptoms occur. These could be signs of a serious condition. <p>If pregnant or breast-feeding, ask a health care professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> ■ drink a full glass of water with each dose ■ adults and children 12 years and over: <ul style="list-style-type: none"> ■ take 1 or 2 capsules every 4 hours or 3 capsules every 6 hours while symptoms persist ■ do not exceed 12 capsules in 24 hours ■ children under 12 years: consult a doctor <p>Other information</p> <ul style="list-style-type: none"> ■ save carton for full directions and warnings ■ store at 15-30°C (59-86°F) ■ avoid excessive heat above 40°C (104°F) ■ do not use if the blue band around the capsule is missing or broken <p>Inactive ingredients</p> <p>anhydrous citric acid, carrageenan, colloidal silicon dioxide, FD&C Blue #1, hypromellose, medium chain triglycerides, pharmaceutical ink, potassium chloride, soy lecithin, titanium dioxide</p> <p>Questions or comments?</p> <p>Call 1-888-842-3678 8:30 AM - 4:00 PM ET, Monday - Friday</p>



Consult your healthcare provider before using this product for your heart.

ASPIRIN THERAPY

VAZALORE™

Aspirin Capsules, 325 mg Pain Reliever/Fever Reducer (NSAID)

30 Liquid-Filled Capsules

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL WITH REPEATED TEXT "SEALED FOR YOUR PROTECTION" UNDER CAP IS MISSING OR BROKEN.

NDC XXXXX-XXX-XX Consult your healthcare provider before using this product for your heart.

New!

ASPIRIN THERAPY

VAZALORE™

Aspirin Capsules, 325 mg Pain Reliever/Fever Reducer (NSAID)

325 mg FIRST LIQUID-FILLED ASPIRIN CAPSULES

30 Liquid-Filled Capsules

112519 XXXXXXXX



LOT
EXP

Consult your healthcare provider before using this product for your heart.

ASPIRIN THERAPY

VAZALORE™

Aspirin Capsules, 325 mg Pain Reliever/Fever Reducer (NSAID)

30 Liquid-Filled Capsules

Distributed by: PLX Pharma Inc.
6285 El Rio, Suite 150
Houston, TX 77054
Visit us at: www.vazalore.com

RETAIN CARTON FOR COMPLETE WARNINGS AND INFORMATION

<p>1A</p> <p>VAZALORE™ Aspirin Capsules, 325mg (NSAID)</p> <p>EXPIRES: MM/DD/YYYY LOT#: 012345 Plx Pharma Inc. 8285 El Rio, Suite 130, Houston, TX 77054</p> <p>TO OPEN: Peel back paper from arrow.</p>	<p>1B</p> <p>VAZALORE™ Aspirin Capsules, 325mg (NSAID)</p> <p>EXPIRES: MM/DD/YYYY LOT#: 012345 Plx Pharma Inc. 8285 El Rio, Suite 130, Houston, TX 77054</p> <p>TO OPEN: Peel back paper from arrow.</p>
<p>1C</p> <p>VAZALORE™ Aspirin Capsules, 325mg (NSAID)</p> <p>EXPIRES: MM/DD/YYYY LOT#: 012345 Plx Pharma Inc. 8285 El Rio, Suite 130, Houston, TX 77054</p> <p>TO OPEN: Peel back paper from arrow.</p>	<p>VAZALORE™ Aspirin Capsules, 325mg (NSAID)</p> <p>XXXXXXX 042919</p>
<p>1D</p> <p>VAZALORE™ Aspirin Capsules, 325mg (NSAID)</p> <p>EXPIRES: MM/DD/YYYY LOT#: 012345 Plx Pharma Inc. 8285 El Rio, Suite 130, Houston, TX 77054</p> <p>TO OPEN: Peel back paper from arrow.</p>	<p>Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed</p>
<p>1E</p> <p>VAZALORE™ Aspirin Capsules, 325mg (NSAID)</p> <p>EXPIRES: MM/DD/YYYY LOT#: 012345 Plx Pharma Inc. 8285 El Rio, Suite 130, Houston, TX 77054</p> <p>TO OPEN: Peel back paper from arrow.</p>	<p>1F</p> <p>VAZALORE™ Aspirin Capsules, 325mg (NSAID)</p> <p>EXPIRES: MM/DD/YYYY LOT#: 012345 Plx Pharma Inc. 8285 El Rio, Suite 130, Houston, TX 77054</p> <p>TO OPEN: Peel back paper from arrow.</p>

Drug Facts

Active ingredient (in each capsule) Purposes
Aspirin (NSAID)* 325 mg Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses
For the temporary relief of minor aches and pains associated with:

- headache ■ backache ■ muscular aches ■ a cold
- toothache ■ minor pain of arthritis
- premenstrual and menstrual cramps
- temporarily reduces fever
- ask your doctor about other uses for aspirin capsules

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives ■ facial swelling ■ shock ■ asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer.

Ask doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout or arthritis

Drug Facts (continued)

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you are experiencing any of the following signs of stomach bleeding:
 - feel faint ■ vomit blood
 - have bloody or black stools
 - have stomach pain that doesn't get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts for more than 3 days
- ringing in the ears or loss of hearing occurs
- redness or swelling is present in the painful area
- any new symptoms occur. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health care professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over:
 - take 1 or 2 capsules every 4 hours or 3 capsules every 6 hours while symptoms persist
 - do not exceed 12 capsules in 24 hours
- children under 12 years: consult a doctor

Other information

- save carton for full directions and warnings
- store at 15-30°C (59-86°F)
- avoid excessive heat above 40°C (104°F)
- do not use if the blue band around the capsule is missing or broken

Inactive ingredients
anhydrous citric acid, carrageenan, colloidal silicon dioxide, FD&C Blue #1, hypromellose, medium chain triglycerides, pharmaceutical ink, potassium chloride, soy lecithin, titanium dioxide

Questions or comments?
Call 1-888-842-3567 8:30 AM - 4:00 PM ET, Monday - Friday

6 Liquid-Filled Capsules

VAZALORE™
Aspirin Capsules, 325 mg
Pain Reliever/Fever Reducer (NSAID)

WXC XXXXX-XXXX-XXX Consult your healthcare provider before using this product for your heart.



325 mg FIRST LIQUID-FILLED ASPIRIN CAPSULES

6 Liquid-Filled Capsules

SAMPLE ONLY - NOT FOR SALE

LOT EXP

112519 XXXXXXXX

PLX PHARMA INC. Distributed by: PLX Pharma Inc.
8785 Ed Rio, Suite 150
Houston, TX 77054
Visit us at: www.vazalore.com

VAZALORE™
Aspirin Capsules, 325 mg
Pain Reliever/Fever Reducer (NSAID)

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPEN OR IF PRINTED SEAL ON BLISTER IS BROKEN OR MISSING.
RETAIN CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts

Active ingredient (in each capsule) Purposes
 Aspirin (NSAID) 325 mg Pain Reliever/Fever Reducer
 Nonsteroidal anti-inflammatory drug

Uses

- for the temporary relief of minor aches and pains associated with:
 - headache
 - backache
 - muscular aches
 - a cold
 - toothache
 - minor pain of arthritis
 - premenstrual and menstrual cramps
- temporarily reduces fever
- ask your doctor about other uses for aspirin capsules

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

This product contains soy.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer.

Ask doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout or arthritis

Drug Facts (continued)

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away, if you are experiencing any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that doesn't get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts for more than 3 days
- ringing in the ears or loss of hearing occurs
- redness or swelling is present in the painful area
- any new symptoms occur. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health care professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over:
 - take 1 or 2 capsules every 4 hours or 3 capsules every 6 hours while symptoms persist
 - do not exceed 12 capsules in 24 hours
- children under 12 years: consult a doctor

Other information

- save carton for full directions and warnings
- store at 15-30°C (59-86°F)
- avoid excessive heat above 40°C (104°F)
- do not use if the blue band around the capsule is missing or broken

Inactive ingredients
 anhydrous citric acid, carrageenan, colloidal silicon dioxide, FD&C Blue #1, hypromellose, medium chain triglycerides, pharmaceutical ink, potassium chloride, soy lecithin, titanium dioxide

Questions or comments?
 Call 1-888-842-3657 8:30 AM - 4:00 PM ET, Monday - Friday

30 Liquid-Filled Capsules

ASPIRIN THERAPY

VAZALORE™

Aspirin Capsules, 325 mg Pain Reliever/Fever Reducer (NSAID)

Consult your healthcare provider before using this product for your heart.




30 Liquid-Filled Capsules

TAMPER EVIDENT:
 DO NOT USE IF CARTON IS OPEN OR IF PRINTED SEAL ON BLISTER IS BROKEN OR MISSING.

Consult your healthcare provider before using this product for your heart.

ASPIRIN THERAPY

VAZALORE™

Aspirin Capsules, 325 mg Pain Reliever/Fever Reducer (NSAID)

325 mg FIRST LIQUID-FILLED ASPIRIN CAPSULES

30 Liquid-Filled Capsules

112519 XXXXXXXX



LOT
EXP

Consult your healthcare provider before using this product for your heart.

ASPIRIN THERAPY

VAZALORE™

Aspirin Capsules, 325 mg Pain Reliever/Fever Reducer (NSAID)

Consult your healthcare provider before using this product for your heart.

PLX PHARMA INC. Distributed by: PLX Pharma Inc.
 8285 El Rio, Suite 130, Houston, TX 77054
 Visit us at: www.vazalore.com

RETAIN CARTON FOR COMPLETE WARNINGS AND INFORMATION

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

KAREN M MAHONEY
12/11/2019 11:45:39 PM