



Marketed by: Pfizer, Madison, NJ 07940, USA. Made in France. © 2019 Pfizer Inc. For most recent product information, visit www.Nexium24HR.com

Do Not Use if foil blister barrier or yellow band on capsule is broken.

Lot: Exp:

PAA114832.FDA01

PAA114832.FDA01


DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8.0 pt
• DRUG FACTS CONTINUED	7.0 pt
• HEADINGS	7.0 pt
• SUBHEADINGS/BODY TEXT	6.0 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5.0 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



**TOP
PANEL 1**



**BASE
PANEL 3**



**RELEASED BACK
PANEL 2**

PAA114834.FDA01


DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.5 pt
• BULLETS	3.5 pt
• HAIRLINES	.5 pt



**TOP
PANEL 1**



**BASE
PANEL 3**



**RELEASED BACK
PANEL 2**

PAA114836.FDA01

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.5 pt
• BULLETS	3.5 pt
• HAIRLINES	.5 pt



KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

Drug Facts
Active ingredient (in each capsule)
 Esomeprazole 20 mg..... Acid reducer
 (Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)

Purpose
 Acid reducer

Uses
 ■ treats frequent heartburn (occurs 2 or more days a week)
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to esomeprazole
Do not use if you have:
 ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
 ■ heartburn with lightheadedness, sweating or dizziness
 ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
 ■ frequent chest pain
 These may be signs of a serious condition. See your doctor.
Ask a doctor before use if you have
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss
 ■ nausea or vomiting ■ stomach pain
Ask a doctor or pharmacist before use if you are
 ■ taking a prescription drug. Acid reducers may interact with certain prescription drugs.
Stop use and ask a doctor if
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea
 ■ you develop a rash or joint pain

Drug Facts (continued)
 If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 ■ adults 18 years of age and older
 ■ this product is to be used once a day (every 24 hours), every day for 14 days
 ■ may take 1 to 4 days for full effect
14-Day Course of Treatment
 ■ swallow 1 capsule with a glass of water before eating in the morning
 ■ take every day for 14 days
 ■ do not take more than 1 capsule a day
 ■ swallow whole. Do not crush or chew capsules.
 ■ do not use for more than 14 days unless directed by your doctor
Repeated 14-Day Courses (if needed)
 ■ you may repeat a 14-day course every 4 months
 ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor
 ■ children under 18 years of age: ask a doctor before use.
 Heartburn in children may sometimes be caused by a serious condition.

Other Information
 ■ read the directions and warnings before use
 ■ keep the carton. It contains important information.
 ■ store at 20-25°C (68-77°F)

Inactive ingredients corn starch, D&C red no. 28, FD&C blue no. 1, FD&C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, pharmaceutical ink, polysorbate 80, sucrose, talc, titanium dioxide, triethyl citrate

Questions or comments? call toll-free weekdays 9 AM to 5 PM EST at 1-866-226-1600



Marketed by:
 Pfizer, Madison, NJ 07940 USA
 © 2019 Pfizer Inc.
 Made in France
 For most recent product information, visit
 www.Nexium24HR.com
 Nexium and the colors purple and gold
 as applied to the capsule are registered
 trademarks of AstraZeneca AB and are
 used under license.

Nexium[®]
 Capsules **24HR**
 Clinically Proven to
 Treat Frequent
 Heartburn

See new warning information NDC 0573-2450-14

Treats Frequent Heartburn

Nexium[®]
 esomeprazole magnesium
 delayed-release capsules **24HR**
 20 mg/acid reducer

14 CAPSULES
 One 14-day course of treatment

May take 1 to 4 days for full effect



Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" or yellow band around the center of each capsule is broken or missing.

PAA114833.FDA01

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9.0 pt
• DRUG FACTS CONTINUED	8.0 pt
• HEADINGS	8.0 pt
• SUBHEADINGS/BODY TEXT	6.0 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5.0 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

NDC 0573-2450-15

See new warning information

Treats Frequent Heartburn

24HR

May take 1 to 4 days for full effect

Capsules



Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

14 CAPSULES

One 14 day course of treatment

Marketed by: Pfizer, Madison, NJ 07940 USA © 2019 Pfizer Inc. Made in France For most recent product information, visit www.Nexium24HR.com

Drug Facts

Active ingredient (in each capsule) Purpose
Esomeprazole 20 mg Acid reducer
(Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)

Uses
■ treats frequent heartburn (occurs 2 or more days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to esomeprazole
Do not use if you have:
■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
■ heartburn with lightheadedness, sweating or dizziness
■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
■ frequent chest pain
These may be signs of a serious condition. See your doctor.
■ Ask a doctor before use if you have
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss ■ nausea or vomiting
■ stomach pain

Ask a doctor or pharmacist before use if you are
■ taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
■ your heartburn continues or worsens
■ you need to take more than 1 course of treatment every 4 months
■ you get diarrhea
■ you develop a rash or joint pain

Questions or comments? call toll-free weekdays 9 AM to 5 PM EST at 1-866-226-1600

Other information
■ read the directions and warnings before use
■ keep the card. It contains important information.
■ store at 20-25°C (68-77°F)

Inactive ingredients corn starch, D.C red no. 28, FD&C blue no. 1, FD&C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, polyorbate 80, sucrose, talc, titanium dioxide, methyl citrate

Directions
■ adults 18 years of age and older
■ this product is to be used once a day (every 24 hours), every day for 14 days
■ may take 1 to 4 days for full effect
14-Day Course of Treatment
■ swallow 1 capsule with a glass of water before eating in the morning
■ take every day for 14 days
■ do not take more than 1 capsule a day
■ do not take more than 14 days unless directed by your doctor
Repeated 14-Day Courses (if needed)
■ you may repeat a 14-day course every 4 months
■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor
■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

KEEP CARD FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

Do Not Use if seal under bottle cap imprinted with "SEALED FOR YOUR PROTECTION" or yellow band around the center of each capsule is broken or missing.

FPO

3 0573 2450 15 5

D000063771.FDA01

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

Drug Facts	
Active ingredient (in each capsule) Esomeprazole 20 mg (Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)	Purpose Acid reducer
Uses ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect	
Warnings Allergy alert: Do not use if you are allergic to esomeprazole Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain These may be signs of a serious condition. See your doctor. Ask a doctor before use if you have ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain Ask a doctor or pharmacist before use if you are ■ taking a prescription drug. Acid reducers may interact with certain prescription drugs. Stop use and ask a doctor if ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain	
Drug Facts (continued) If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ may take 1 to 4 days for full effect 14-Day Course of Treatment ■ swallow 1 capsule with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 capsule a day ■ swallow whole. Do not crush or chew capsules. ■ do not use for more than 14 days unless directed by your doctor Repeated 14-Day Courses (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.	
Other Information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F)	
Inactive ingredients corn starch, D&C red no. 28, FD&C blue no. 1, FD&C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, pharmaceutical ink, polysorbate 80, sucrose, talc, titanium dioxide, triethyl citrate	
Questions or comments? call toll-free weekdays 9 AM to 5 PM EST at 1-866-226-1600	



Marketed by:
Pfizer, Madison, NJ 07940 USA
© 2019 Pfizer Inc.
Made in France
For most recent product information, visit
www.Nexium24HR.com
Nexium and the colors purple and gold as applied to the capsule are registered trademarks of AstraZeneca AB and are used under license.

Easy Open Cap!

Nexium[®] 24HR

Capsules

See new warning information
Treats Frequent Heartburn

NDC 0573-2450-28

28 CAPSULES
Two 14-day courses of treatment

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.

May take 1 to 4 days for full effect

Arthritis Foundation Ease of Use CAP

2 BOTTLES INSIDE

EASY OPEN CAP!


Actual Size

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" or yellow band around the center of each capsule is broken or missing.



PAA114835.FDA01

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9.0 pt
• DRUG FACTS CONTINUED	8.0 pt
• HEADINGS	8.0 pt
• SUBHEADINGS/BODY TEXT	6.0 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5.0 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

Drug Facts
Active ingredient (in each capsule) Purpose
Esomeprazole 20 mg Acid reducer
(Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)

Drug Facts (continued)
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Uses
■ treats frequent heartburn (occurs 2 or more days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Directions
■ adults 18 years of age and older
■ this product is to be used once a day (every 24 hours), every day for 14 days
■ may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to esomeprazole
Do not use if you have:
■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
■ heartburn with lightheadedness, sweating or dizziness
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
■ frequent chest pain
These may be signs of a serious condition. See your doctor.

14-Day Course of Treatment
■ swallow 1 capsule with a glass of water before eating in the morning
■ take every day for 14 days
■ do not take more than 1 capsule a day
■ swallow whole. Do not crush or chew capsules.
■ do not use for more than 14 days unless directed by your doctor
Repeated 14-Day Courses (if needed)
■ you may repeat a 14-day course every 4 months
■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor
■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Ask a doctor before use if you have
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss
■ nausea or vomiting ■ stomach pain

Other Information
■ read the directions and warnings before use
■ keep the carton. It contains important information.
■ store at 20-25°C (68-77°F)

Ask a doctor or pharmacist before use if you are
■ taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Inactive ingredients
corn starch, D&C red no. 28, FD&C blue no. 1, FD&C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, pharmaceutical ink, polysorbate 80, sucrose, talc, titanium dioxide, triethyl citrate

Stop use and ask a doctor if
■ your heartburn continues or worsens
■ you need to take this product for more than 14 days
■ you need to take more than 1 course of treatment every 4 months
■ you get diarrhea
■ you develop a rash or joint pain

Questions or comments?
call toll-free weekdays 9 AM to 5 PM EST at 1-866-226-1600

Tips for Managing Heartburn
• Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
• Eat slowly and do not eat big meals.
• Do not eat late at night or just before bedtime.
• Do not lie flat or bend over soon after eating.
• Raise the head of your bed.
• Wear loose-fitting clothing around your stomach.
• If you are overweight, lose weight.
• If you smoke, quit smoking.



Marketed by:
Pfizer, Madison, NJ 07940 USA
© 2019 Pfizer Inc.
Made in France
For most recent product information, visit www.Nexium24HR.com
Nexium and the colors purple and gold as applied to the capsule are registered trademarks of AstraZeneca AB and are used under license.

Nexium[®] Clinically Proven to Treat Frequent Heartburn
Capsules 24HR

See new warning information NDC 0573-2450-42
Treats Frequent Heartburn




42 CAPSULES
Three 14-day courses of treatment
Capsules
May take 1 to 4 days for full effect


3 BOTTLES INSIDE



ACTUAL SIZE
Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" or yellow band around the center of each capsule is broken or missing.



PAA114837.FDA01
FPO
3 0573 2450 42 1



PAA114837.FDA01

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9.0 pt
• DRUG FACTS CONTINUED	8.0 pt
• HEADINGS	8.0 pt
• SUBHEADINGS/BODY TEXT	6.0 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5.0 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

NDC 0573-2450-43

See new warning information

Treats Frequent Heartburn



24HR

May take 1 to 4 days for full effect

3 Pack

Capsules 

42 CAPSULES

Three 14-day courses of treatment

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" or yellow band around the center of each capsule is broken or missing.



24HR

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

KEEP CARD FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

<p>Drug Facts</p> <p>Active ingredient (in each capsule) Esomeprazole 20 mg (Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)</p> <p>Purpose Acid reducer</p> <p>Uses</p> <ul style="list-style-type: none"> ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect <p>Warnings</p> <p>Allergy alert: Do not use if you are allergic to esomeprazole</p> <p>Do not use if you have:</p> <ul style="list-style-type: none"> ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain <p>These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> ■ taking a prescription drug. Acid reducers may interact with certain prescription drugs. <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain 	<p>Drug Facts (continued)</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ may take 1 to 4 days for full effect <p>14-Day Course of Treatment</p> <ul style="list-style-type: none"> ■ swallow 1 capsule with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 capsule a day ■ swallow whole. Do not crush or chew capsules. ■ do not use for more than 14 days unless directed by your doctor <p>Repeated 14-Day Courses (if needed)</p> <ul style="list-style-type: none"> ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor <p>■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p> <p>Other Information</p> <ul style="list-style-type: none"> ■ read the directions and warnings before use ■ keep the card. It contains important information. ■ store at 20-25°C (68-77°F) <p>Inactive ingredients</p> <p>corn starch, D&C red no. 28, FD&C blue no. 1, FD&C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, pharmaceutical ink, polysorbate 80, sucrose, talc, titanium dioxide, triethyl citrate</p> <p>Questions or comments? call toll-free weekdays 9 AM to 5 PM EST at 1-866-226-1600</p>
---	--

Marketed by: Pfizer, Madison, NJ 07940 USA
© 2019 Pfizer Inc.

Made in France

For most recent product information, visit www.Nexium24HR.com

Nexium and the colors purple and gold as applied to the capsule are registered trademarks of AstraZeneca AB and are used under license.

FPO

3 0573 2450 43 8

D000063792.FDA01

D000063792.FDA01

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	10.0 pt
• DRUG FACTS CONTINUED	9.0 pt
• HEADINGS	8.0 pt
• SUBHEADINGS/BODY TEXT	7.0 pt
• LEADING	7.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	6.0 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



Drug Facts
Active ingredient (in each capsule) Esomeprazole 20 mg Acid reducer
Purpose (Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)
Uses ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn, this drug may take 1 to 4 days for full effect
Warnings
Allergy alert: Do not use if you are allergic to esomeprazole
Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain These may be signs of a serious condition. See your doctor.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Drug Facts (continued)

KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

Tips for Managing Heartburn • Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables. • Eat slowly and do not eat late at night or just before bedtime. • Do not lie flat or bend over soon after eating. • Raise the head of your bed. • Wear loose-fitting clothing around your stomach. • If you are overweight, lose weight. • If you smoke, quit smoking.

PAA114749.FDA01

Do not use if foil blister wrapper or yellow band on capsule is broken.

Made by: Pfizer, Madison, NJ 07740 USA Made in France © 2019 Pfizer Inc. For most recent product information, visit www.Nexium24HR.com

Lot: Exp:

Drug Facts (continued)
Directions ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ may take 1 to 4 days for full effect
14-Day Course of Treatment ■ swallow 1 capsule with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 capsule a day ■ do not take more than 14 days unless directed by your doctor
Repeated 14-Day Courses (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor
Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F)
Inactive ingredients carmine, corn starch, FD&C blue no. 2, FD&C blue no. 2 aluminum lake, FD&C red no. 3, FD&C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, titanium dioxide, triethyl citrate
Questions or comments? call toll-free weekdays 9 AM to 5 PM EST at 1-866-226-1600
Drug Facts (continued)



PAA114749.FDA01


DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



TOP
PANEL 1



BASE
PANEL 3



RELEASED BACK
PANEL 2

PAA115135.FDA01

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.5 pt
• BULLETS	3.5 pt
• SPACE BEFORE BULLET	2 ems
• WARNING BOX LINE, HAIRLINES	.5 pt



KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

Drug Facts	Purpose
Active ingredient (in each capsule) Esomeprazole 20 mg (Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)	Acid reducer
Uses ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect	
Warnings Allergy alert: Do not use if you are allergic to esomeprazole Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain These may be signs of a serious condition. See your doctor. Ask a doctor before use if you have ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain Ask a doctor or pharmacist before use if you are ■ taking a prescription drug. Acid reducers may interact with certain prescription drugs. Stop use and ask a doctor if ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain	
Drug Facts (continued) If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ may take 1 to 4 days for full effect 14-Day Course of Treatment ■ swallow 1 capsule with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 capsule a day ■ swallow whole. Do not crush or chew capsules. ■ do not use for more than 14 days unless directed by your doctor Repeated 14-Day Courses (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.	
Other Information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F)	
Inactive ingredients carmine, corn starch, FD&C blue no. 2, FD&C blue no. 2 aluminum lake, FD&C red no. 3, FD&C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, pharmaceutical ink, polysorbate 80, sucrose, talc, titanium dioxide, triethyl citrate	
Questions or comments? call toll-free weekdays 9 AM to 5 PM EST at 1-866-226-1600	



Marketed by:
Pfizer, Madison, NJ 07940 USA
© 2019 Pfizer Inc.
Made in France
For most recent product information, visit www.Nexium24HR.com
Nexium is a registered trademark of AstraZeneca AB and is used under license.

Nexium[®] 24HR Clear Minis
Clinically Proven to Treat Frequent Heartburn

See new warning information NDC 0573-2452-14

Treats Frequent Heartburn

Nexium[®] 24HR
esomeprazole magnesium delayed-release capsules 20 mg/acid reducer

14 CAPSULES
One 14-day course of treatment

Clear Minis

May take 1 to 4 days for full effect

1 BOTTLE INSIDE

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" or yellow band around the center of each capsule is broken or missing.

PAA115134.FDA01

PAA115134.FDA01

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

Drug Facts
Active ingredient (in each capsule) Purpose
 Esomeprazole 20 mg Acid reducer
 (Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)

Uses
 ■ treats frequent heartburn (occurs 2 or more days a week)
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to esomeprazole
Do not use if you have:
 ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
 ■ heartburn with lightheadedness, sweating or dizziness
 ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
 ■ frequent chest pain
 These may be signs of a serious condition. See your doctor.
Ask a doctor before use if you have
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss
 ■ nausea or vomiting ■ stomach pain
Ask a doctor or pharmacist before use if you are
 ■ taking a prescription drug. Acid reducers may interact with certain prescription drugs.
Stop use and ask a doctor if
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea
 ■ you develop a rash or joint pain

Drug Facts (continued)
If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 ■ adults 18 years of age and older
 ■ this product is to be used once a day (every 24 hours), every day for 14 days
 ■ may take 1 to 4 days for full effect
14-Day Course of Treatment
 ■ swallow 1 capsule with a glass of water before eating in the morning
 ■ take every day for 14 days
 ■ do not take more than 1 capsule a day
 ■ swallow whole. Do not crush or chew capsules.
 ■ do not use for more than 14 days unless directed by your doctor
Repeated 14-Day Courses (if needed)
 ■ you may repeat a 14-day course every 4 months
 ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor
 ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other Information
 ■ read the directions and warnings before use
 ■ keep the carton. It contains important information.
 ■ store at 20-25°C (68-77°F)

Inactive ingredients
 carmine, corn starch, FD&C blue no. 2, FD&C blue no. 2 aluminum lake, FD&C red no. 3, FD&C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, pharmaceutical ink, polysorbate 80, sucrose, talc, titanium dioxide, triethyl citrate

Questions or comments? call toll-free weekdays 9 AM to 5 PM EST at 1-866-226-1600



Marketed by:
 Pfizer, Madison, NJ 07940 USA
 © 2019 Pfizer Inc.
 Made in France
 For most recent product information, visit
 www.Nexium24HR.com
 Nexium is a registered trademark of
 AstraZeneca AB and is used under license.

Nexium 24HR Clinically Proven to **Treat Frequent Heartburn**

ClearMinis™

See new warning information

Treats Frequent Heartburn

Nexium 24HR
 esomeprazole magnesium delayed-release capsules
 20 mg/acid-reducer

42 CAPSULES

Three 14-day courses of treatment

May take 1 to 4 days for full effect

ClearMinis™

NDC 0573-2452-42



Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" or yellow band around the center of each capsule is broken or missing.

PAA115133.FDA01

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

NDC 0573-2452-43

See new warning information

Treats Frequent Heartburn

24HR

3 Pack

Clear Minis™

May take 1 to 4 days for full effect

4 CAPSULES

Three 14-day courses of treatment

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" or yellow band around the center of each capsule is broken or missing.

24HR

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
• Eat slowly and do not eat big meals.
• Do not eat late at night or just before bedtime.
• Do not lie flat or bend over soon after eating.
• Raise the head of your bed.
• Wear loose-fitting clothing around your stomach.
• If you are overweight, lose weight.
• If you smoke, quit smoking.

KEEP CARD FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

Drug Facts

Active ingredient (in each capsule) Purpose
Esomeprazole 20 mg Acid reducer
(Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)

Uses

- treats frequent heartburn (occurs 2 or more days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to esomeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
■ heartburn with lightheadedness, sweating or dizziness
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
■ frequent chest pain
These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss
■ nausea or vomiting
■ stomach pain

Ask a doctor or pharmacist before use if you are

- taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
■ you need to take this product for more than 14 days
■ you need to take more than 1 course of treatment every 4 months
■ you get diarrhea
■ you develop a rash or joint pain

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults 18 years of age and older
■ this product is to be used once a day (every 24 hours), every day for 14 days
■ may take 1 to 4 days for full effect

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
■ take every day for 14 days
■ do not take more than 1 capsule a day
■ swallow whole. Do not crush or chew capsules.
■ do not use for more than 14 days unless directed by your doctor
Repeated 14-Day Courses (if needed)
■ you may repeat a 14-day course every 4 months
■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor
■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other Information

- read the directions and warnings before use
■ keep the card. It contains important information.
■ store at 20-25°C (68-77°F)

Inactive ingredients

carmine, corn starch, FD&C blue no. 2, FD&C blue no. 2 aluminum lake, FD&C red no. 3, FD&C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, pharmaceutical ink, polysorbate 80, sucrose, talc, titanium dioxide, triethyl citrate

Questions or comments?

call toll-free weekdays 9 AM to 5 PM EST at 1-866-226-1600



D000063839.FDA01

Marketed by: Pfizer, Madison, NJ 07940 USA
© 2019 Pfizer Inc.

Made in France

For most recent product information, visit www.Nexium24HR.com

Nexium is a registered trademark of AstraZeneca AB and is used under license.



D000063839.FDA01

Table with 2 columns: DRUG FACTS TEXT DEFINED, TYPE SIZE. Rows include DRUG FACTS TITLE (10 pt), DRUG FACTS CONTINUED (9 pt), HEADINGS (9 pt), SUBHEADINGS/BODY TEXT (7 pt), LEADING (7.5 pt), # OF CHARACTERS PER INCH (<39), BULLETS (6 pt), SPACE BEFORE BULLET (2 ems), BARLINES, HAIRLINES (1.5 pt, .5 pt), SPACE BETWEEN HAIRLINES AND BOX END (2 spaces).

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

VALERIE S PRATT
04/04/2019 04:29:28 PM