

Batch No.

EXP

8U174 00 V5 102218

Location for lot number and expiration date

Drug Facts	Warnings	Do not use	Ask a doctor before use if you have	Other information	Inactive ingredients	Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST	
Drug Facts (continued)	Active ingredient (in each tablet) Lansoprazole 15 mg;Acid Reducer	Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect	Do not use ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. See your doctor.	Ask a doctor before use if you have ■ frequent heartburn over 3 months. This may be a sign of a more serious condition. ■ heartburn with hightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or hightheadedness ■ frequent chest pain ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain	Other information ■ read the directions, warnings and package insert before use ■ keep the carton and package insert. They contain important information. ■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity. ■ protect product from moisture	Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopollose, fumed silica, hypromellose, hydroxyethylcellulose, maize maltoedextrin, maltitol, mannitol, magnesium, microcrystalline cellulose, polyethylene glycol, propylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, talc, titanium dioxide, triethyl citrate	Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST
Directions ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours	14-Day Course of Treatment ■ take 1 tablet before eating in the morning ■ do not crush or chew tablets ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.	14-Day Course (if needed) ■ do not take this medicine with alcohol ■ do not use for more than 14 days unless directed by your doctor	Do not take for 14 days ■ take every day for 14 days	do not take for more than 14 days unless directed by your doctor ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor	children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.		

NOT FOR RESALE

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

Safety Feature – Do not use if printed tablet blister unit is open or torn.

Compare to the active ingredient in Prevacid® 24 HR*



Treats Frequent Heartburn **24 Hour**

Lansoprazole

Delayed Release
Orally Disintegrating Tablets 15 mg
Acid Reducer

Melts in your mouth
Dissolves without water

Strawberry Flavor



14 TABLETS

One 14-day course of treatment
• May take 1 to 4 days for full effect

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

*This product is not manufactured or distributed by
GSK Consumer Healthcare, distributor of Prevacid® 24 HR.

Questions or comments? 1-800-719-9260, weekdays 7:30 AM to 5:00 PM EST

Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopollose, flavin, hypromellose, hypromellose phthalate, maize maltoedextrin, methylcellulose, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium lauryl sulfate, polyvinylpyrrolidone, sucralose, sugars, spheres, talc, titanium dioxide, triethyl citrate

Other information
■ read the directions, warnings and package insert before use
■ keep the carton and package insert. They contain important information.
■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity.
■ protect product from moisture

4 months unless directed by a doctor
■ children under 18 years of age, ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.
■ you may repeat a 14-day course every 4 months
■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

Repeated 14-Day Course (if needed)
■ do not take this medicine with alcohol
■ do not use for more than 14 days unless directed by your doctor
■ do not take more than 1 tablet a day
■ take every day for 14 days
■ The tablets can also be swallowed whole with water.
■ place the tablet on tongue; tablet disintegrates, with or without water.
■ do not crush or chew tablets
■ take 1 tablet before eating in the morning

14-Day Course of Treatment
■ relief of symptoms within 24 hours
■ it may take 1 to 4 days for full effect, although some people get complete relief sooner
■ this product is to be used once a day (every 24 hours), every day for 14 days
■ adults 18 years of age and older

Directions (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

If pregnant or breast-feeding, ask a health care professional before use.
■ you get diarrhea ■ you develop a rash or joint pain
■ you need to take more than 1 course of treatment every 4 months
■ you need to take this product for more than 14 days
■ your heartburn continues or worsens
Stop use and ask a doctor if

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.
■ nausea or vomiting ■ stomach pain
■ unexplained weight loss
■ frequent wheezing, particularly with heartburn
■ frequent chest pain
■ spreading to arms, neck or shoulders, or lightheadedness
■ chest pain or shoulder pain with shortness of breath; sweating; pain
■ heartburn with **lightheadedness, sweating or dizziness**
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ liver disease

Ask a doctor before use if you have
■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools; these may be signs of a serious condition. See your doctor.

Warnings
■ **Allergy alert:** do not use if you are allergic to lansoprazole
■ Do not use

Use
■ treats frequent heartburn (occurs 2 or more days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Purpose
Lansoprazole 15 mg.....Acid Reducer

Active ingredient (in each tablet)

Location for lot number and expiration date

8U174 FA C6 092818



KEEP THE CARTON AND PACKAGE INSERT.
THEY CONTAIN IMPORTANT INFORMATION.

Safety Feature –
Do not use if printed tablet blister unit is open or torn.

NDC 64239-181-01

Treats Frequent Heartburn



Compare to the active ingredient
in Prevacid® 24 HR*

24 Hour

Lansoprazole

Delayed Release

Orally Disintegrating Tablets 15 mg



Strawberry Flavor

Acid Reducer

■ May take 1 to 4 days for full effect

14 TABLETS

ONE 14-DAY COURSE
OF TREATMENT

Melts in your mouth
Dissolves without water



Actual size

STANDARD FORMAT
TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box



Made in Israel
 Manufactured by:
 Dexcel Pharma Technologies Ltd
 10 Hakdima Street, Yokneam 2069200, Israel
 *This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Prevacid® 24 HR.

Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg

Drug Facts
 Active ingredient (in each tablet) Lansoprazole 15 mg.....Acid Reducer
Use
 ■ treats frequent heartburn (occurs 2 or more days a week)
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect
Warnings
 ■ **Do not use** if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
 ■ **Allergy alert:** do not use if you are allergic to lansoprazole
Ask a doctor before use if you have
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ heartburn with lightheadedness, sweating or dizziness
 ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
 ■ frequent wheezing, particularly with heartburn
 ■ frequent chest pain
 ■ unexplained weight loss
 ■ nausea or vomiting ■ stomach pain
Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.
Stop use and ask a doctor if
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea
 ■ you develop a rash or joint pain
If pregnant or breast-feeding, ask a health care professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).
Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

Drug Facts (continued)
Directions
 ■ adults 18 years of age and older
 ■ this product is to be used once a day (every 24 hours), every day for 14 days
 ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours
14-Day Course of Treatment
 ■ take 1 tablet before eating in the morning
 ■ do not crush or chew tablets
 ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
 ■ take every day for 14 days
 ■ do not take more than 1 tablet a day
 ■ do not use for more than 14 days unless directed by your doctor
 ■ do not take this medicine with alcohol
Repeated 14-Day Course (if needed)
 ■ you may repeat a 14-day course every 4 months
 ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor
 ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.
Other information
 ■ read the directions, warnings and package insert before use
 ■ keep the carton and package insert. They contain important information.
 ■ store at 20-25°C (68-77°F) ■ Keep product out of high heat and humidity
 ■ protect product from moisture
Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, crospovidone, flavor, hypromellose, hypromellose phthalate, maize maltoedextrin, maltitol, mannitol, microcrystalline cellulose, polyethylene glycol, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

8U130 FA C6 092818

Location for lot number and expiration date

Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg



Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

NDC 64239-181-02

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

24 Hour
Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg



Strawberry Flavor

Acid Reducer
 ■ May take 1 to 4 days for full effect

28 TABLETS TWO 14-DAY COURSES OF TREATMENT

Melts in your mouth
 Dissolves without water



STANDARD FORMAT
TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box

Location for lot number and expiration date

8U155 FA C6 092818

Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg

Drug Facts

Active ingredient (in each tablet) Acid Reducer
Lansoprazole 15 mg

Use
■ treats frequent heartburn (occurs 2 or more days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Do not use
■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
Ask a doctor before use if you have
■ liver disease
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ heartburn with lightheadedness, sweating or dizziness
■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
Frequent chest pain
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss
■ nausea or vomiting
■ stomach pain
Ask a doctor or pharmacist before use if you are taking a prescription drug.
Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
■ your heartburn continues or worsens
■ you need to take this product for more than 14 days
■ you need to take more than 1 course of treatment every 4 months
■ you get diarrhea
■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Other information
■ read the directions, warnings and package insert before use
■ keep the carton and package insert. They contain important information.
■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity
■ protect product from moisture

Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopolone, flavor, hypromellose, hypromellose phthalate, maize starch, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

Drug Facts (continued)

Directions
■ adults 18 years of age and older
■ this product is to be used once a day (every 24 hours), every day for 14 days
■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
■ take 1 tablet before eating in the morning
■ do not crush or chew tablets
■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
■ take every day for 14 days
■ do not take more than 1 tablet a day
■ do not use for more than 14 days unless directed by your doctor.
■ do not take this medicine with alcohol

Repeated 14-Day Course (if needed)
■ you may repeat a 14-day course every 4 months
■ do not take for more than 14 days or more often than every 4 months
unless directed by a doctor
■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg



Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

NDC 64239-181-03

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

24 Hour
Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg



Strawberry Flavor

Acid Reducer

- May take 1 to 4 days for full effect

42 TABLETS

THREE 14-DAY COURSES OF TREATMENT

Melts in your mouth
Dissolves without water



Actual size

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

Made in Israel
 Manufactured by:
Dexel Pharma Technologies Ltd
 10 Hakidma Street, Yokneam 2069200, Israel

*This product is not manufactured or distributed by
 GSK Consumer Healthcare, distributor of Prevacid® 24 HR.

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Lansoprazole 15 mg; Acid Reducer</p> <p>Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Warnings ■ Allergy alert: do not use if you are allergic to lansoprazole</p> <p>Do not use ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have ■ liver disease ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness</p> <p>Chest pain ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain</p> <p>Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain</p> <p>If pregnant or breast-feeding, ask a health care professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p>	<p>Directions ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours</p> <p>14-Day Course of Treatment ■ take 1 tablet before eating in the morning ■ do not crush or chew tablets ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water. ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ do not take this medicine with alcohol</p> <p>Repeated 14-Day Course (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor</p> <p>Other information ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition. ■ read the directions, warnings and package insert before use ■ keep the carton and package insert. They contain important information. ■ store at 20-25°C (68-77°F) ■ Keep product out of high heat and humidity. ■ protect product from moisture</p> <p>Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopollose, HAVON, hypromellose, hypromellose pthalate, maize maibedextrin, malitol, mannitol, meglumine, microcrystalline cellulose, polyvinylpyrrolidone, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate</p> <p>Questions or comments? 1-800-719-9266; weekdays 7:30 AM to 5:00 PM EST</p>
---	---

Location for lot number and expiration date

8U174 FA C6-V 100418



KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

Safety Feature –
 Do not use if printed tablet blister unit is open or torn.

NDC 64239-181-01

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

24 Hour
Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg



Strawberry Flavor

Acid Reducer

■ May take 1 to 4 days for full effect

Orally Disintegrating Tablets



Actual size

14 TABLETS

ONE 14-DAY COURSE OF TREATMENT

Melts in your mouth
 Dissolves without water

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box



Manufactured by:
 Dexcel Pharma Technologies Ltd
 10 Hakdima Street, Yokneam 2069200, Israel
 Made in Israel

*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Prevacid® 24 HR.

Drug Facts
 Active ingredient (in each tablet) Lansoprazole 15 mg
 Purpose Acid Reducer

Use
 ■ treats frequent heartburn (occurs 2 or more days a week)
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
 Allergy alert: do not use if you are allergic to lansoprazole
 Do not use
 ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
 Ask a doctor before use if you have
 ■ liver disease
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ heartburn with lightheadedness, sweating or dizziness
 ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness
 ■ frequent wheezing, particularly with heartburn
 ■ frequent chest pain
 ■ unexplained weight loss
 ■ nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug.
 Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea
 ■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.
 ■ If pregnant or breast-feeding, ask a health care professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

5:00 PM EST
 Questions or comments? 1-800-719-9260: weekdays 7:30 AM to

Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg

Location for lot number and expiration date

8U130 FA C6-V 100418

Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg



Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

NDC 64239-181-02

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

24 Hour
Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg



Acid Reducer
 ■ May take 1 to 4 days for full effect

Orally Disintegrating Tablets

Melts in your mouth
 Dissolves without water



28 TABLETS TWO 14-DAY COURSES OF TREATMENT

STANDARD FORMAT

TYPE SIZES:
 TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued) 8 pt Helvetica Neue LT Std 57 Condensed
 HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
 TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
 LEADING: 6.5 pt
 BULLETS: 5 pt Square with 2 M'S spacing between statements
 TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
 HEAVY LINES: 1.5 pt
 HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box

Location for lot number and expiration date

8U155 FA C6-V 100418

Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg

Drug Facts

Active ingredient (in each tablet) Lansoprazole 15 mg, Acid Reducer

Purpose

Use

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: do not use if you are allergic to lansoprazole

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Questions or comments? 1-800-719-9260, weekdays 7:30 AM to 5:00 PM EST

Drug Facts (continued)

Directions

- adults 18 years of age and older: this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours
- take 1 tablet before eating in the morning

14-Day Course of Treatment

- take 1 tablet before eating in the morning
- do not crush or chew tablets
- place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- do not take this medicine with alcohol

Repeated 14-Day Course (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor

Other information

- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.
- read the directions, warnings and package insert before use
- keep the carton and package insert. They contain important information.
- store at 20-25°C (68-77°F). ■ keep product out of high heat and humidity
- protect product from moisture

Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, crospovidone, flavor, hypromellose, hypromellose phthalate, maize malodextrin, methyl, mannitol, magnesium, microcrystalline cellulose, polyorbate 80, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hekidma Street, Yokneam 209200, Israel
*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Prevacid® 24 HR.



Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg

Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

NDC 64239-181-03

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

24 Hour
Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg



Acid Reducer

- May take 1 to 4 days for full effect

Orally Disintegrating Tablets

42 TABLETS THREE 14-DAY COURSES OF TREATMENT

Melts in your mouth
Dissolves without water



STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box

Made in Israel
 Manufactured by:
Dexcel Pharma Technologies Ltd
 10 Hakidma Street, Yokneam 2069200, Israel

*This product is not manufactured or distributed by
 GSK Consumer Healthcare, distributor of Prevacid® 24 HR.

<p>Drug Facts (continued)</p> <p>Directions ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours</p> <p>14-Day Course of Treatment ■ take 1 tablet before eating in the morning ■ do not crush or chew tablets ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water. ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ do not take this medicine with alcohol</p> <p>Repeated 14-Day Course (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor</p> <p>■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p> <p>Other Information ■ read the directions, warnings and package insert before use ■ keep the carton and package insert. They contain important information. ■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity. ■ protect product from moisture</p> <p>Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopolone, hawthorn, hypromellose, hypromellose phthalate, maize maltoedextrin, mannitol, magnesium, microcrystalline cellulose, poly sorbate 80, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate</p> <p>Questions or comments? 1-800-719-9260, weekdays 7:30 AM to 6:00 PM EST</p>	<p>Drug Facts</p> <p>Active ingredient (in each tablet) Lansoprazole 15 mg;Acid Reducer</p> <p>Purpose ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Warnings ■ Allergy alert: do not use if you are allergic to lansoprazole ■ Do not use ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools; these may be signs of a serious condition. See your doctor. ■ Ask a doctor before use if you have ■ liver disease ■ had heartburn over 3 months; this may be a sign of a more serious condition. ■ heartburn with difficulty breathing, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating, pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain</p> <p>Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if ■ your heartburn continues for more than 14 days ■ you need to take this product for more than 14 months ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain</p> <p>If pregnant or breast-feeding, ask a health care professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p>
--	---

Location for lot number and expiration date

8U174 FA C6-J 100418



KEEP THE CARTON AND PACKAGE INSERT.
 THEY CONTAIN IMPORTANT INFORMATION.

Safety Feature –
 Do not use if printed tablet blister unit is open or torn.

NDC 64239-181-01

Treats Frequent Heartburn

24 Hour

Lansoprazole

Delayed Release Orally Disintegrating Tablets 15 mg

Acid Reducer

- May take 1 to 4 days for full effect

14 TABLETS ONE 14-DAY COURSE OF TREATMENT

MELTech™
 Melts In Your Mouth

Compare to the active ingredient in Prevacid® 24 HR*

Child-resistant



Strawberry Flavor



Actual size

Melts in your mouth
 Dissolves without water

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued); 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box



*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Prevacid® 24 HR.

Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakdima Street, Yokneam 2069200, Israel

Lansoprazole

Delayed Release
Orally Disintegrating Tablets 15 mg

Drug Facts (continued)

Directions
adults 18 years of age and older
this product is to be used once a day (every 24 hours), every day for 14 days
it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
take 1 tablet before eating in the morning
do not crush or chew tablets
place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
take every day for 14 days
do not take more than 1 tablet a day
do not use for more than 14 days unless directed by your doctor
do not take this medicine with alcohol

Repeated 14-Day Course (if needed)
you may repeat a 14-day course every 4 months
do not take for more than 14 days or more often than every 4 months
unless directed by a doctor

children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information
read the directions, warnings and package insert before use
keep the carton and package insert. They contain important information.
store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity
protect product from moisture

Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopolone, flavor, hypromellose, hypromellose phthalate, maize macrodextrin, maltitol, mannitol, meglumine, microcrystalline cellulose, polyborate 80, propylene glycol, silicon dioxide, sodium stearoyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-719-9260, weekdays 7:30 AM to 5:00 PM EST

Active ingredient (in each tablet)
Lansoprazole 15 mg.....Acid Reducer

Purpose
Acid Reducer

Use
treats frequent heartburn (occurs 2 or more days a week)
not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: do not use if you are allergic to lansoprazole
Do not use
if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
Ask a doctor before use if you have
liver disease
head heartburn over 3 months. This may be a sign of a more serious condition.
heartburn with lightheadedness, sweating or dizziness
chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
frequent chest pain
frequent wheezing, particularly with heartburn
unexplained weight loss
nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug.
Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
your heartburn continues or worsens
you need to take this product for more than 14 days
you need to take more than 1 course of treatment every 4 months
you get diarrhea
you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

8U130 FA C6-J 100418

Location for lot number and expiration date




Lansoprazole

Delayed Release
Orally Disintegrating Tablets 15 mg

Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.



NDC 64239-181-02

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

Child-resistant

24 Hour Lansoprazole

Delayed Release
Orally Disintegrating Tablets 15 mg





Strawberry Flavor

Acid Reducer
■ May take 1 to 4 days for full effect

28 TABLETS TWO 14-DAY COURSES OF TREATMENT

Melts in your mouth
Dissolves without water



STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box



Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel
Made in Israel

*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Prevacid® 24 HR.

Drug Facts

Active ingredient (in each tablet)
Lansoprazole 15 mg.....Acid Reducer

Use
■ treats frequent heartburn (occurs 2 or more days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: do not use if you are allergic to lansoprazole

Do not use
■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have
■ liver disease
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ heartburn with lightheadedness, sweating or dizziness
■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness

Frequent chest pain
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss
■ nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug.
Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
■ your heartburn continues or worsens
■ you need to take this product for more than 14 days
■ you need to take more than 1 course of treatment every 4 months
■ you get diarrhea
■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Questions or comments? 1-800-719-9260. weekdays 7:30 AM to 5:00 PM EST

Other information
■ read the directions, warnings and package insert before use
■ keep the carton and package insert. They contain important information.
■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity
■ protect product from moisture

Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopolone, flavor, hypromellose, hypromellose phthalate, maize starch, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucrose, sugar spheres, talc, titanium dioxide, triethyl citrate

Directions
■ adults 18 years of age and older
■ this product is to be used once a day (every 24 hours), every day for 14 days
■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
take 1 tablet before eating in the morning

do not crush or chew tablets
place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.

do not use for more than 14 days unless directed by your doctor.
do not take this medicine with alcohol

Repeated 14-Day Course (if needed)
you may repeat a 14-day course every 4 months
do not take for more than 14 days or more often than every 4 months unless directed by a doctor


■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15mg

8U130 FA C6-A 100418

Location for lot number and expiration date

Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg



Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

NDC 64239-181-02


Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

Value 2 Pack

24 Hour
Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg



Strawberry Flavor

Acid Reducer
■ May take 1 to 4 days for full effect

28 TABLETS TWO 14-DAY COURSES OF TREATMENT

Melts in your mouth
Dissolves without water



Actual size

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

Location for lot number and expiration date

8U155 FA C6-B 100418

Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg

Drug Facts

Active ingredient (in each tablet) Lansoprazole 15 mg.....Acid Reducer

Use
■ treats frequent heartburn (occurs 2 or more days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
■ Ask a doctor before use if you have liver diseases
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ heartburn with lightheadedness, sweating or dizziness
■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
■ frequent chest pain
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss
■ nausea or vomiting
■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug.
Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
■ your heartburn continues or worsens
■ you need to take this product for more than 14 days
■ you need to take more than 1 course of treatment every 4 months
■ you get diarrhea
■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Questions or comments? 1-800-719-9260. weekdays 7:30 AM to 5:00 PM EST

Drug Facts (continued)

Directions
■ this product is to be used once a day (every 24 hours), every day for 14 days
■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours
■ take 1 tablet before eating in the morning

14-Day Course of Treatment
■ take 1 tablet before eating in the morning
■ do not crush or chew tablets
■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
■ take every day for 14 days
■ do not take more than 1 tablet a day
■ do not use for more than 14 days unless directed by your doctor
■ do not take this medicine with alcohol

Repeated 14-Day Course (if needed)
■ you may repeat a 14-day course every 4 months
■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

Other information
■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.
■ read the directions, warnings and package insert before use
■ keep the carton and package insert. They contain important information.
■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity
■ protect product from moisture

Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, crospovidone, flavor, hypromellose, hypromellose phthalate, maize polyorbital, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate, poly sorbate 80, propylene glycol, silicon dioxide, sodium stearyl fumarate, naldexidine, methylol, mannitol, magnesium, microcrystalline cellulose, crospovidone, crospovidone, flavor, hypromellose, hypromellose phthalate, maize polyorbital, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Manufactured by:
Decel Pharma Technologies Ltd
10 Hakdima Street, Yokneam 2069200, Israel
Made in Israel

*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Prevacid® 24 HR.



Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg

Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

NDC 64239-181-03

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

Value 3 Pack

24 Hour
Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg



Strawberry Flavor

Acid Reducer

- May take 1 to 4 days for full effect

42 TABLETS

THREE 14-DAY COURSES OF TREATMENT

Melts in your mouth
Dissolves without water



Actual size

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box



*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Prevacid® 24 HR.

Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakdima Street, Yokneam 2069200, Israel

Lansoprazole

Delayed Release
Orally Disintegrating Tablets 15 mg

Drug Facts

Active Ingredient (in each tablet)
Lansoprazole 15 mg
Acid Reducer

Purpose
Acid Reducer

Use
■ Treats frequent heartburn (occurs 2 or more days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: do not use if you are allergic to lansoprazole

Do not use
■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have
■ liver disease
■ had heartburn over 3 months. This may be a sign of a more serious condition.

Chest pain
■ frequent wheezing, particularly with heartburn
■ frequent weight loss
■ unexplained vomiting or nausea or stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug.
Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
■ your heartburn continues or worsens
■ you need to take this product for more than 14 days
■ you need to take more than 1 course of treatment every 4 months
■ you get diarrhea
■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

Drug Facts (continued)

Directions
■ adults 18 years of age and older
■ this product is to be used once a day (every 24 hours), every day for 14 days
■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
■ take 1 tablet before eating in the morning

Warnings
■ do not crush or chew tablets
■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.

Do not use
■ take every day for 14 days
■ do not take more than 1 tablet a day
■ do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Course (if needed)
■ do not take this medicine with alcohol

Other information
■ you may repeat a 14-day course every 4 months
■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor
■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information
■ read the directions, warnings and package insert before use
■ keep the carton and package insert. They contain important information.
■ store at 20-25°C (68-77°F) ■ Keep product out of high heat and humidity
■ protect product from moisture

Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopolone, flavor, hypromellose, hypromellose phthalate, maize polyborate 80, propylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

8U130 FA C6-K 100418

Location for lot number and expiration date



Lansoprazole

Delayed Release
Orally Disintegrating Tablets 15 mg

Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

NDC 64239-181-02

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

Buy One 14 Count, Get One 14 Count Free!

24 Hour Lansoprazole

Delayed Release
Orally Disintegrating Tablets 15 mg



Strawberry Flavor

Acid Reducer
■ May take 1 to 4 days for full effect

28 TABLETS TWO 14-DAY COURSES OF TREATMENT

Melts in your mouth
Dissolves without water



STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box



Manufactured by:
 Dexcel Pharma Technologies Ltd
 10 Hakdama Street, Yokneam 2069200, Israel
 *This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Prevacid® 24 HR.

Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg


Drug Facts
 Active Ingredient (in each tablet) Lansoprazole 15 mg.....Acid Reducer
Use
 ■ treats frequent heartburn (occurs 2 or more days a week)
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect
Warnings
 Allergy alert: do not use if you are allergic to lansoprazole
 Do not use
 ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
 ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a more serious condition.
Ask a doctor before use if you have
 ■ liver disease
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.
chest pain
 ■ frequent wheezing, particularly with heartburn
 ■ frequent weight loss
 ■ nausea or vomiting
Ask a doctor or pharmacist before use if you are taking a prescription drug.
 Acid reducers may interact with certain prescription drugs.
Stop use and ask a doctor if
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea
 ■ you develop a rash or joint pain
If pregnant or breast-feeding, ask a health care professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).
Questions or comments? 1-800-719-9260, weekdays 7:30 AM to 5:00 PM EST

Drug Facts (continued)
Directions
 ■ take 1 tablet before eating in the morning
 ■ do not crush or chew tablets
 ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
 ■ take every day for 14 days
 ■ do not take more than 1 tablet a day
 ■ do not use for more than 14 days unless directed by your doctor
 ■ do not take this medicine with alcohol
Repeated 14-Day Course (if needed)
 ■ you may repeat a 14-day course every 4 months
 ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor
 ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.
Other information
 ■ read the directions, warnings and package insert before use
 ■ keep the carton and package insert. They contain important information.
 ■ store at 20-25°C (68-77°F). ■ keep product out of high heat and humidity
 ■ protect product from moisture
Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopollose, flavor, hypromellose, hypromellose phthalate, maize starch, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

8U130 FA C6-L 100418

Location for lot number and expiration date

Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg



Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

NDC 64239-181-02

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

100% More Free!
 14 Free Tablets!

24 Hour
Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg



Strawberry Flavor

Acid Reducer
 ■ May take 1 to 4 days for full effect


28 TABLETS TWO 14-DAY COURSES OF TREATMENT

Melts in your mouth
 Dissolves without water



Actual size

STANDARD FORMAT
TYPE SIZES:
 TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued) 8 pt Helvetica Neue LT Std 57 Condensed
 HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
 TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
 LEADING: 6.5 pt
 BULLETS: 5 pt Square with 2 M'S spacing between statements
 TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
 HEAVY LINES: 1.5 pt
 HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box



Manufactured by:
 Dexcel Pharma Technologies Ltd
 10 Hakdima Street, Yokneam 2069200, Israel
 Made in Israel

*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Prevacid® 24 HR.

Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg

Drug Facts
Active Ingredient (in each tablet)
 Lansoprazole 15 mg.....Acid Reducer

Use
 ■ Treats frequent heartburn (occurs 2 or more days a week)
 ■ Not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
 ■ Allergy alert: do not use if you are allergic to lansoprazole
 ■ Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have
 ■ Liver disease
 ■ Had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ Heartburn with lightheadedness, sweating or dizziness
 ■ Chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness

Chest pain
 ■ Frequent wheezing, particularly with heartburn
 ■ Unexplained weight loss
 ■ Nausea or vomiting

Ask a doctor or pharmacist before use if you are taking a prescription drug.
 Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
 ■ Your heartburn continues or worsens
 ■ You need to take this product for more than 14 days
 ■ You need to take more than 1 course of treatment every 4 months
 ■ You get diarrhea
 ■ You develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.
 ■ Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Questions or comments? 1-800-719-9260, weekdays 7:30 AM to 5:00 PM EST

Drug Facts (continued)
Directions
 ■ This product is to be used once a day (every 24 hours), every day for 14 days
 ■ Adults 18 years of age and older
 ■ It may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
 ■ Take 1 tablet before eating in the morning
 ■ Do not crush or chew tablets
 ■ Place the tablet on tongue, tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
 ■ Take every day for 14 days
 ■ Do not take for more than 14 days unless directed by your doctor.
 ■ Do not take this medicine with alcohol

Repeated 14-Day Course (if needed)
 ■ You may repeat a 14-day course every 4 months
 ■ Do not take for more than 14 days or more often than every 4 months unless directed by a doctor

Other information
 ■ Children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Inactive ingredients
 ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopolone, flavor, hypromellose, hypromellose phthalate, maize starch, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Keep the carton and package insert. They contain important information.
 ■ Store at 20-25°C (68-77°F) ■ Keep product out of high heat and humidity
 ■ Protect product from moisture

8U130 FA C6-P 100418

Location for lot number and expiration date

Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg



Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

NDC 64239-181-02

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

Bonus Size! 14 Free

24 Hour
Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg



Strawberry Flavor

Acid Reducer
 ■ May take 1 to 4 days for full effect

28 TABLETS TWO 14-DAY COURSES OF TREATMENT

Melts in your mouth
 Dissolves without water



Actual size

STANDARD FORMAT
TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box



This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Prevacid® 24 HR.
 Dexcel Pharma Technologies Ltd
 10 Hakidma Street, Yokneam 2069200, Israel
 Manufactured by:
 Made in Israel

Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg

<p>Drug Facts Active ingredient (in each tablet) Lansoprazole 15 mg Acid Reducer</p>	<p>Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p>
<p>Warnings ■ Allergy alert: do not use if you are allergic to lansoprazole ■ Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.</p>	<p>Ask a doctor before use if you have ■ liver disease ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain</p>
<p>Other information ■ read the directions, warnings and package insert before use ■ keep the carton and package insert. They contain important information. ■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity ■ protect product from moisture</p>	<p>Stop use and ask a doctor if ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain</p>
<p>Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopollose, flavor, hypromellose, hypromellose phthalate, maize starch, mannitol, magnesium stearate, microcrystalline cellulose, polyborate 80, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate</p>	<p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p>
<p>Directions ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours</p>	<p>Questions or comments? 1-800-719-9260, weekdays 7:30 AM to 5:00 PM EST</p>

8U130 FA C6-R 100418

Location for lot number and expiration date

Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg



Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

NDC 64239-181-02

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

Bonus! + 14 Free

24 Hour
Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg



Strawberry Flavor

Acid Reducer
 ■ May take 1 to 4 days for full effect

28 TABLETS TWO 14-DAY COURSES OF TREATMENT

Melts in your mouth
 Dissolves without water



STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box



Manufactured by:
 Dexcel Pharma Technologies Ltd
 10 Hakdima Street, Yokneam 2069200, Israel
 Made in Israel
 *This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Prevacid® 24 HR.

Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg

Drug Facts
Active ingredient (in each tablet)
 Lansoprazole 15 mg.....Acid Reducer

Use
 ■ Treats frequent heartburn (occurs 2 or more days a week)
 ■ Not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
 ■ Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
 ■ If you have heartburn over 3 months. This may be a sign of a more serious condition.
 ■ Heartburn with lightheadedness, sweating or dizziness
 ■ Chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
 ■ Frequent wheezing, particularly with heartburn
 ■ Unexplained weight loss
 ■ Nausea or vomiting
 ■ Stomach pain

Ask a doctor before use if you have
 ■ Liver disease
 ■ Had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ If you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Do not use
 ■ If you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Allergy alert: do not use if you are allergic to lansoprazole

Other information
 ■ Children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.
 ■ Unless directed by a doctor
 ■ Do not take for more than 14 days or more often than every 4 months
 ■ You may repeat a 14-day course every 4 months
 ■ Repeated 14-Day Course (if needed)
 ■ Do not take this medicine with alcohol
 ■ Do not use for more than 14 days unless directed by your doctor
 ■ Do not take more than 1 tablet a day
 ■ Take every day for 14 days
 ■ The tablets can also be swallowed whole with water.
 ■ Place the tablet on tongue, tablet disintegrates, with or without water.
 ■ Do not crush or chew tablets
 ■ Take 1 tablet before eating in the morning
 ■ 14-Day Course of Treatment
 ■ Relief of symptoms within 24 hours
 ■ It may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours
 ■ Adults 18 years of age and older
 ■ This product is to be used once a day (every 24 hours), every day for 14 days

Directions
 ■ Adults 18 years of age and older
 ■ This product is to be used once a day (every 24 hours), every day for 14 days

Purpose
 ■ Adults 18 years of age and older
 ■ This product is to be used once a day (every 24 hours), every day for 14 days

Drug Facts (continued)

Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

Inactive ingredients
 ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopolone, flavor, hypromellose, hypromellose phthalate, maize maltoedextrin, mannitol, meglumine, microcrystalline cellulose, polyvinylpyrrolidone, polyvinylpyrrolidone, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Stop use and ask a doctor if
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea
 ■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

8U130 FA C6-S 100418

Location for lot number and expiration date

Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg



Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

NDC 64239-181-02

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

Bonus! 14 Tablets FREE

24 Hour
Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg



Strawberry Flavor

Acid Reducer
 ■ May take 1 to 4 days for full effect

28 TABLETS TWO 14-DAY COURSES OF TREATMENT

Melts in your mouth
 Dissolves without water



Actual size

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box

Location for lot number and expiration date

8U155 FA C6-M 100418

Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg

Drug Facts

Active ingredient (in each tablet) Lansoprazole 15 mg.....Acid Reducer

Use
 ■ Treats frequent heartburn (occurs 2 or more days a week)
 ■ Not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
 ■ Do not use if you are allergic to lansoprazole
 ■ Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
 ■ Ask a doctor before use if you have:
 ■ liver disease
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness
 ■ frequent chest pain
 ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss
 ■ nausea or vomiting
 ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug.
 Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea
 ■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Questions or comments? 1-800-719-9260. weekdays 7:30 AM to 5:00 PM EST

Other information
 ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.
 ■ unless directed by a doctor
 ■ do not take for more than 14 days or more often than every 4 months
 ■ you may repeat a 14-day course every 4 months
Repeated 14-Day Course (if needed)
 ■ do not take this medicine with alcohol
 ■ do not use for more than 14 days unless directed by your doctor
 ■ take every day for 14 days
 ■ do not take more than 1 tablet a day
 ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
 ■ do not crush or chew tablets
 ■ take 1 tablet before eating in the morning

14-Day Course of Treatment
 ■ take 1 tablet before eating in the morning
 ■ relief of symptoms within 24 hours
 ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

Directions
 ■ this product is to be used once a day (every 24 hours), every day for 14 days
 ■ adults 18 years of age and older

Purpose
 Acid Reducer

Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, crospovidone, flavor, hypromellose, hydroxypropylcellulose, mize, polydextrose, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Manufactured by:
 Dexcel Pharma Technologies Ltd
 10 Hakdima Street, Yokneam 2069200, Israel

*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Prevacid® 24 HR.



Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg

Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

NDC 64239-181-03

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

Bonus Pack: 28 + 14 Free

24 Hour
Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg



Strawberry Flavor

Acid Reducer

- May take 1 to 4 days for full effect

42 TABLETS

THREE 14-DAY COURSES OF TREATMENT

Melts in your mouth
Dissolves without water



Actual size

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box

Location for lot number and expiration date


8U155 FA C6-N 100418

Made in Israel
 Manufactured by:
 Cexel Pharma Technologies Ltd
 10 Hakdama Street, Yokneam 2069200, Israel
 *This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Prevacid® 24 HR.

Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg

Drug Facts
Active ingredient (in each tablet) Lansoprazole 15 mg, Acid Reducer
Purpose
Use
 ■ treats frequent heartburn (occurs 2 or more days a week)
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect
Warnings
 ■ Allergy alert: do not use if you are allergic to lansoprazole
 ■ Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
 ■ Ask a doctor before use if you have liver disease
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ heartburn with **lightheadedness, sweating or dizziness**
 ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
 ■ frequent chest pain
 ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss
 ■ nausea or vomiting
 ■ stomach pain
Ask a doctor or pharmacist before use if you are taking a prescription drug.
 Acid reducers may interact with certain prescription drugs.
Stop use and ask a doctor if
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea
 ■ you develop a rash or joint pain
If pregnant or breast-feeding, ask a health care professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).
 5:00 PM EST
Questions or comments? 1-800-719-9260, weekdays 7:30 AM to



Drug Facts (continued)
Directions
 ■ adults 18 years of age and older
 ■ this product is to be used once a day (every 24 hours), every day for 14 days
 ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours
14-Day Treatment
 ■ take 1 tablet before eating in the morning
Do not crush or chew tablets
 ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
 ■ take every day for 14 days
 ■ do not take more than 1 tablet a day
 ■ do not use for more than 14 days unless directed by your doctor
 ■ do not take this medicine with alcohol
Repeated 14-Day Course (if needed)
 ■ you may repeat a 14-day course every 4 months
 ■ do not take for more than 14 days or more often than every 4 months
unless directed by a doctor
 ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.
Other information
 ■ read the directions, warnings and package insert before use
 ■ keep the carton and package insert. They contain important information.
 ■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity
 ■ protect product from moisture
Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, crospovidone, flavor, hypromellose, hypromellose phthalate, maize starch, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate



Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg

Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.



NDC 64239-181-03

Treats Frequent Heartburn

MELTech™
 Melts In Your Mouth

Compare to the active ingredient in Prevacid® 24 HR*

Bonus Size! 14 Tablets Free

24 Hour
Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg


Acid Reducer
 ■ May take 1 to 4 days for full effect

42 TABLETS THREE 14-DAY COURSES OF TREATMENT

Melts in your mouth
 Dissolves without water

Strawberry Flavor

Actual size



STANDARD FORMAT

TYPE SIZES:
 TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued) 8 pt Helvetica Neue LT Std 57 Condensed
 HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
 TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
 LEADING: 6.5 pt
 BULLETS: 5 pt Square with 2 M'S spacing between statements
 TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
 HEAVY LINES: 1.5 pt
 HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box

Location for lot number and expiration date

8U155 FA C6-R 100418

Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg

Drug Facts

Active ingredient (in each tablet) Lansoprazole 15 mg.....Acid Reducer

Use
■ treats frequent heartburn (occurs 2 or more days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
■ Ask a doctor before use if you have liver diseases
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ heartburn with lightheadedness, sweating or dizziness
■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
■ frequent chest pain
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss
■ nausea or vomiting
■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug.
Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
■ your heartburn continues or worsens
■ you need to take this product for more than 14 days
■ you need to take more than 1 course of treatment every 4 months
■ you get diarrhea
■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Questions or comments? 1-800-719-9260. weekdays 7:30 AM to 5:00 PM EST

Drug Facts (continued)

Directions
■ this product is to be used once a day (every 24 hours), every day for 14 days
■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours
■ take 1 tablet before eating in the morning

14-Day Course of Treatment
■ take 1 tablet before eating in the morning
■ do not crush or chew tablets
■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
■ take every day for 14 days
■ do not take more than 1 tablet a day
■ do not use for more than 14 days unless directed by your doctor
■ do not take this medicine with alcohol

Repeated 14-Day Course (if needed)
■ you may repeat a 14-day course every 4 months
■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

Other information
■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.
■ read the directions, warnings and package insert before use
■ keep the carton and package insert. They contain important information.
■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity
■ protect product from moisture

Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, crospovidone, flavor, hypromellose, hypromellose phthalate, maize polyborate 80, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Made in Israel
Manufactured by:
Decel Pharma Technologies Ltd
10 Hakdama Street, Yokneam 2069200, Israel
*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Prevacid® 24 HR.



Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg

Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

NDC 64239-181-03

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

Bonus! + 14 Free

24 Hour
Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg



Strawberry Flavor

Acid Reducer

- May take 1 to 4 days for full effect

42 TABLETS THREE 14-DAY COURSES OF TREATMENT

Melts in your mouth
Dissolves without water



STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

Location for lot number and expiration date

8U155 FA C6-S 100418

Made in Israel
 Manufactured by:
 Dexcel Pharma Technologies Ltd
 10 Hakidma Street, Yokneam 2069200, Israel

*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Prevacid® 24 HR.



Drug Facts
 Active ingredient (in each tablet)
 Lansoprazole 15 mg.....Acid Reducer

Use
 ■ treats frequent heartburn (occurs 2 or more days a week)
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
 ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
 ■ Ask a doctor before use if you have:
 liver disease
 had heartburn over 3 months. This may be a sign of a more serious condition.
 heartburn with lightheadedness, sweating or dizziness
 chest pain or shoulder pain with shortness of breath; swelling; pain spreading to arms, neck or shoulders; or rightheadedness
 frequent chest pain
 frequent wheezing, particularly with heartburn
 unexplained weight loss
 nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug.
 Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea
 ■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

Drug Facts (continued)
Directions
 ■ this product is to be used once a day (every 24 hours), every day for 14 days
 ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
 ■ take 1 tablet before eating in the morning

Do not crush or chew tablets
 ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
 ■ take every day for 14 days
 ■ do not take more than 1 tablet a day
 ■ do not use for more than 14 days unless directed by your doctor
 ■ do not take this medicine with alcohol
 ■ unless directed by a doctor

Repeated 14-Day Course (if needed)
 ■ you may repeat a 14-day course every 4 months
 ■ do not take for more than 14 days or more often than every 4 months

Other information
 ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Inactive ingredients
 ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopolone, flavoc, hypromellose, hypromellose phthalate, maize polyorbate 80, propylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Keep the carton and package insert. They contain important information.
 ■ keep the carton and package insert. They contain important information.
 ■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity
 ■ protect product from moisture

Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15mg



Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15mg

Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.



NDC 64239-181-03

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

Bonus! 14 Tablets FREE

24 Hour
Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15mg



Strawberry Flavor

Acid Reducer

- May take 1 to 4 days for full effect

42 TABLETS

THREE 14-DAY COURSES OF TREATMENT

Melts in your mouth
 Dissolves without water



Actual size

STANDARD FORMAT

TYPE SIZES:

TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued); 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box

Treats Frequent Heartburn

Lansoprazole

Delayed Release

Orally Disintegrating Tablets 15 mg

acid reducer

- May take 1 to 4 days for full effect

Please read the entire package insert before taking
Lansoprazole Delayed Release Orally Disintegrating Tablets.
Save for future reference.

How Lansoprazole delayed release orally disintegrating tablets Treats Your Frequent Heartburn

Lansoprazole delayed release orally disintegrating tablets stops acid production at the source – the **pumps** that release acid into the stomach. Lansoprazole delayed release orally disintegrating tablet is taken once a day (every 24 hours), every day for 14 days.

What You Can Expect When Taking Lansoprazole delayed release orally disintegrating tablets

Frequent heartburn can occur anytime during the 24-hour period (day or night). Take Lansoprazole delayed release orally disintegrating tablets in the morning before eating. Lansoprazole is clinically proven to treat frequent heartburn. Although some people get complete relief of symptoms within 24 hours, it may take 1 to 4 days for full effect. Make sure you take Lansoprazole delayed release orally disintegrating tablets every day for 14 days to treat your frequent heartburn.

Who Should Take Lansoprazole delayed release orally disintegrating tablets

Adults (18 years and older) with **frequent heartburn** – when you have heartburn 2 or more days a week.

Who Should NOT Take Lansoprazole delayed release orally disintegrating tablets

People who have one episode of heartburn a week or less, or who want immediate relief of heartburn.

How to Take Lansoprazole delayed release orally disintegrating tablets

14-DAY Course of Treatment

- Take 1 tablet before eating in the morning.
- **Do not crush or chew tablets.**
- Place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
- Take every day for 14 days.
- Do not take more than 1 tablet a day.
- Do not use for more than 14 days unless directed by your doctor.
- Do not take this medicine with alcohol.

When to Take Lansoprazole delayed release orally disintegrating tablets Again

You may repeat a 14-day course of therapy every 4 months.

When to Talk to Your Doctor

Do not take for more than 14 days or more often than every 4 months unless directed by a doctor.



Warnings and When to Ask Your Doctor

Allergy alert: Do not use if you are allergic to lansoprazole

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

Clinical studies prove lansoprazole effectively treats frequent heartburn

In three clinical studies, lansoprazole was shown to be significantly better than placebo in treating frequent heartburn.

How Lansoprazole delayed release orally disintegrating tablets is Sold

This medicine is available in 14 tablet, 28 tablet and 42 tablet sizes. These sizes contain one, two and three 14-day courses of treatment, respectively. Do not use for more than 14 days in a row unless directed by your doctor. For the 28 count (two 14-day courses) and the 42 count (three 14-day courses), you may repeat a 14-day course every 4 months.

For Questions or Comments About Lansoprazole delayed release orally disintegrating tablets

Call 1-800-719-9260


weekdays 7:30 AM to 5:00 PM EST

Made in Israel

Manufactured by:

Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

: 80100 00 J3 100418



This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

VALERIE S PRATT
04/09/2019 02:42:39 PM