

Reference ID: 4381841

- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight ■ Quit smoking

Drug Facts (continued)

Strength	Active ingredient (in each tablet)	Purpose
----------	------------------------------------	---------

*This product is not manufactured or distributed by Procter & Gamble, distributor of Pilosec OTC®.

1Y974 FA C4 092618

→ **lot number and expiration date**



MELTech™
Melts In Your Mouth

Compare to Prilosec OTC®*

Treats *FREQUENT* Heartburn!



Omeprazole

Delayed Release
Orally Disintegrating Tablets 20 mg
Acid Reducer

14 TABLETS

One 14-day course of treatment
May take 1 to 4 days for full effect

Melts in your mouth
Dissolves without water



Strawberry Flavor



Actual

STANDARD FORMAT

TYPE SIZES:

TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

(continued) **8 pt Helvetica Neue LT Std 57 Condensed**

HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

SIIBHEADINGS: 6 pt Helvetica Neue IT Std 77 Bold Condensed

TEXT: 6 pt Helvetica Neue LT Std 57 Condensed

TEXT: 6 pt Helve

LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 MS spacing between statements

BULLETS: 5 pt Square with 2 M'S spacing between statement

TELEPHONE #: 6 pt

HEAVY LINES: 1.5

HAIRLINES: 0.5 pt

Hairlines extend to within two spaces of the “Drug Facts” box

Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

Omeprazole

Delayed Release
Orally Disintegrating Tablets 20 mg

Drug Facts (continued)

Directions

- For adults 18 years of age and older
- This product is to be used once a day (every 24 hours), every day for 14 days
- You should take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- Take 1 tablet before eating in the morning
- Do not crush or chew tablets
- Place the tablet on tongue; tablet disintegrates, with or without water.
- The tablet can also be swallowed whole with water.

Take every day for 14 days

- Do not take more than 1 tablet a day
- Do not use for more than 14 days unless directed by your doctor
- Do not take this medicine with alcohol

Repeated 14-Day Course (if needed)

- You may repeat a 14-day course every 4 months
- Do not take for more than 14 days or more often than every 4 months unless directed by a doctor

Children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

- Read the directions and warnings before use
- Keep the carton. It contains important information.
- Store at 20°-25°C (68°-77°F); keep product out of high heat and moisture

Inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscopollose, ferric oxide, hypromellose, hypromellose phthalate, inulin, croscopollose, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

<p>Drug Facts</p> <p>Active ingredient (in each tablet)</p> <p>Acid reducer</p> <p>omeprazole 20 mg</p>	<p>Use</p> <ul style="list-style-type: none"> ■ Treats frequent heartburn (occurs 2 or more days a week) ■ Not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect 	<p>Warnings</p> <p>Alert: do not use if you are allergic to omeprazole</p>	<p>Do not use if you have:</p> <ul style="list-style-type: none"> ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with pain, light-headedness, sweating or dizziness ■ chest or shoulder pain with shortness of breath, sweating, or pain spreading to arms, neck or shoulders; or lightheadedness 	<p>These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have:</p> <ul style="list-style-type: none"> ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain 	<p>Ask a doctor or pharmacist before use if you are taking a prescription drug.</p> <p>Acid reducers may interact with certain prescription drugs.</p>	<p>Stop use and ask a doctor if:</p> <ul style="list-style-type: none"> ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain 	<p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-2222).</p>
---	---	--	--	---	---	---	--

1Y930 FA C4 082718

**Location for
lot number and
expiration date**



- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Safety Feature –
Do not use if printed tablet
blister unit is open or torn.



NDC 64239-XXX-02

Treats *FREQUENT* Heartburn!



MELTech™
Melts In Your Mouth

Compare to Prilosec OTC®*

Omeprazole

Delayed Release
Orally Disintegrating Tablets 20 mg
Acid Reducer

28 TABLETS

Two 14-day courses of treatment
May take 1 to 4 days for full effect

Melts in your mouth
Dissolves without water



Strawberry Flavor



Actual
size

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued) **8 pt Helvetica Neue LT Std 57 Condensed**
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box



STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued) **8 pt Helvetica Neue LT Std 57 Condensed**
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the “Drug Facts” box

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Questions or comments? 1-800-719-9260: weekdays 7:30 AM to 5:00 PM EST

Inactive ingredients: amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscopollose, ferrous fumarate, hypromellose, hypromellose phthalate, maize malboxdextrin, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Other information: ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture

4 months unless directed by a doctor
■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.
■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor
■ you may repeat a 14-day course every 4 months

Repeated 14-Day Course (if needed)
■ do not take this medicine with alcohol
■ do not use for more than 14 days unless directed by your doctor
■ do not take more than 1 tablet a day
■ take every day for 14 days
The tablets can also be swallowed whole with water.

■ place the tablet on tongue; tablet disintegrates, with or without water.
■ do not crush or chew tablets
■ take 1 tablet before eating in the morning

14-Day Course of Treatment
■ take 1 tablet before eating in the morning
■ symptoms within 24 hours
■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours
■ this product is to be used once a day (every 24 hours), every day for 14 days

Directions ■ for adults 18 years of age and older

contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).
■ If pregnant or breast-feeding, ask a health professional before use.
■ you develop a rash or joint pain
■ you get diarrhea
■ you need to take more than 1 course of treatment every 4 months
■ you need to take this product for more than 14 days
■ your heartburn continues or worsens
■ Stop use and ask a doctor if:

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.
■ nausea or vomiting
■ unexplained weight loss
■ frequent wheezing, particularly with heartburn
■ had heartburn over 3 months. This may be a sign of a more serious condition.

Ask a doctor before use if you have:
These may be signs of a serious condition. See your doctor.
■ frequent chest pain
to arms, neck or shoulders, or lightheadedness
■ chest or shoulder pain with shortness of breath, sweating, pain spreading
■ heartburn with lightheadedness, sweating or dizziness
■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools

Do not use if you have:
Allergy alert: do not use if you are allergic to omeprazole
Warnings

■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect
Use ■ treats frequent heartburn (occurs 2 or more days a week)

Omeprazole 20 mg.....Acid reducer
Purpose

Active ingredient (in each tablet)



NDC 64239-XXX-01

Safety Feature –
Do not use if printed tablet blister unit is open or torn.



Compare to Prilosec OTC®*

Treats **FREQUENT** Heartburn!



Omeprazole

Delayed Release
Orally Disintegrating Tablets 20 mg
Acid Reducer

14 TABLETS

One 14-day course of treatment
May take 1 to 4 days for full effect



Strawberry
Flavor

Orally
Disintegrating
Tablets

Melts in your mouth
Dissolves without water



STANDARD FORMAT

TYPE SIZES:

TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

1Y955 FA C4-K 092618

Directions
For adults 18 years of age and older
● this product is to be used once a day (every 24 hours), every day for 14 days
● may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-day Treatment
● take 1 tablet before eating in the morning
● do not crush or chew tablets
● place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
● take every day for 14 days
● do not take more than 1 tablet a day
● do not eat for more than 14 days unless directed by your doctor
● do not take the medicine with alcohol

Repeated 14-Day Course (if needed)
● you may repeat a 14-day course every 6 months
● do not take for more than 14 days or more often than every 6 months unless directed by a doctor

Other information
● children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.
● read the directions and warnings before use
● keep the carton, if container information.
● store at 20-25°C (68-77°F); keep product out of high heat and moisture

Inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscopollose, ferric oxide, flavor, hypromellose, polyethylene glycol, silicon dioxide, sodium stearate, sodium steryl fumarate, sorbitol, sucrose, sugar spheres, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-719-9260: weekdays 7:30 AM to 5:00 PM EST

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking



NDC 64239-XXX-03



Compare to Prilosec OTC®*



Strawberry
Flavour

Orally Disintegrating Tablets

Melts in your mouth
Dissolves without water

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued) **8 pt Helvetica Neue LT Std 57 Condensed**
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box

Directions ■ For adults 18 years of age and older this product is to be used once a day (every 24 hours), every day for 14 days ■ I may take 1 to 4 days for full effect, some people get complete relief of symptoms within 2-4 hours

14-Day Course of Treatment

- Take 1 tablet before eating in the morning
- Do not crush or chew tablets
- Place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
- Take every day for 14 days
- Do not take more than 1 tablet a day
- Do not use for more than 14 days unless directed by your doctor
- Do not take this medicine with alcohol

Repeated 14-Day Course (if needed)

- You may repeat a 14-day course every 4 months
- Do not take for more than 14 days or more often than every 4 months unless directed by a doctor

■ Children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information ■ Read the directions and warnings before use ■ Keep the carton, it contains important information. ■ Store at 20°-25°C (68°-77°F); keep product out of high heat and moisture

Inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscarmellose, ferric oxide, flavor, hypromellose, hypromellose phthalate, maize malodextrin, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

Warnings

Allergy alert: do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Purpose Acid reducer

Active ingredient (in each tablet) Omeprazole 20 mg

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Purpose Acid reducer

Active ingredient (in each tablet) Omeprazole 20 mg

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Purpose Acid reducer

Active ingredient (in each tablet) Omeprazole 20 mg

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Purpose Acid reducer

Active ingredient (in each tablet) Omeprazole 20 mg

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Purpose Acid reducer

Active ingredient (in each tablet) Omeprazole 20 mg

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Purpose Acid reducer

Active ingredient (in each tablet) Omeprazole 20 mg

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Purpose Acid reducer

Active ingredient (in each tablet) Omeprazole 20 mg

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Purpose Acid reducer

Active ingredient (in each tablet) Omeprazole 20 mg

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Purpose Acid reducer

Active ingredient (in each tablet) Omeprazole 20 mg

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Purpose Acid reducer

Active ingredient (in each tablet) Omeprazole 20 mg

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Purpose Acid reducer

Active ingredient (in each tablet) Omeprazole 20 mg

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Purpose Acid reducer

Active ingredient (in each tablet) Omeprazole 20 mg

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■



0 00000 00000 0

Safety Feature –
Do not use if printed tablet
blister unit is open or torn.

NDC 64239-XXX-01

MELTech™
Melts In Your Mouth

Treats **FREQUENT** Heartburn!

 **24 HR**

Compare to Prilosec OTC®*

Child-resistant

Omeprazole

Delayed Release
Orally Disintegrating Tablets 20 mg
Acid Reducer

14 TABLETS

One 14-day course of treatment
May take 1 to 4 days for full effect

Melts in your mouth
Dissolves without water



Strawberry
Flavor



Actual
size



000000000000

STANDARD FORMAT

TYPE SIZES:

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed

HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed

TEXT: 6 pt Helvetica Neue LT Std 57 Condensed

LEADING: 6.5 pt

BULLETS: 5 pt Square with 2 M'S spacing between statements

TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed

HEAVY LINES: 1.5 pt

HAIRLINES: 0.5 pt

Hairlines extend to


Handlines extend to within the spaces of the Drag Race box



<p>Drug Facts</p> <p>Active ingredient (in each tablet) Omeprazole 20 mg.....Acid reducer</p> <p>Purpose Acid reducer</p>	<p>Use</p> <ul style="list-style-type: none"> not intended for immediate relief of heartburn; this drug may take 1 to 4 days trials frequent heartburn (occurs 2 or more days a week) <p>for full effect</p>	<p>Warnings</p> <p>Allergy alert: do not use if you are allergic to omeprazole</p> <p>Do not use if you have:</p> <ul style="list-style-type: none"> trouble or pain swallowing food, vomiting with blood, or bloody or black stools heartburn with lightheadedness, sweating or dizziness chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulder; or lightheadedness <p>Frequent chest pain</p> <ul style="list-style-type: none"> These may be signs of a serious condition. See your doctor. 	<p>Ask a doctor before use if you have:</p> <ul style="list-style-type: none"> heartburn over 5 months. This may be a sign of a more serious condition. unexplained weight loss nausea or vomiting <p>Ask a doctor or pharmacist before use if you are taking a prescription drug.</p> <p>Acid reducers may interact with certain prescription drugs.</p>	<p>Stop use and ask a doctor if:</p> <ul style="list-style-type: none"> your heartburn continues or worsens you need to take this product for more than 14 days you need to take more than 1 course of treatment every 4 months you get diarrhea you develop a rash or joint pain <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact ►</p>	<p>Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST</p>	<p>Directions</p> <ul style="list-style-type: none"> for adults: 18 years of age and older this product is to be used once a day (every 24 hours), every day for 14 days it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours <p>14-Day Course of Treatment</p> <ul style="list-style-type: none"> take 1 tablet before eating in the morning do not crush or chew tablets place the tablet on tongue; tablet disintegrates, with or without water. <p>The tablet can also be swallowed whole with water. </p>	<p>Repeat 14-Day Course (if needed)</p> <ul style="list-style-type: none"> do not take this medicine with alcohol do not use for more than 14 days unless directed by your doctor do not take more than 1 tablet a day take every day for 14 days <p>do not take more than 14 days or more often than every 4 months unless directed by a doctor</p>	<p>Other information</p> <ul style="list-style-type: none"> children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition. you may repeat a 14-day course every 4 months 	<p>Inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscopollose, ferric oxide, flavor, hypromellose, microcrystalline cellulose, polyethylene glycol, silicon stearate, sodium fumarate, sorbitol, croscopollose, sugar spheres, talc, titanium dioxide, triethyl citrate</p> <p>■ store at 20-25°C (68-77°F); keep product out of high heat and moisture</p> <p>■ keep the carton. It contains important information.</p> <p>■ read the directions and warnings before use</p>
--	--	---	--	--	--	--	--	---	---

Omeprazole

Delayed Release
Orally Disintegrating Tablets 20 mg



**Safety Feature –
Do not use if printed tablet
blister unit is open or torn.**

NDC 64239-XXX-03

MELTech™
Melts In Your Mouth

Treats **FREQUENT** Heartburn!

 **24 HR**

Compare to Prilosec OTC®**
Child-resistant

Omeprazole

Delayed Release
Orally Disintegrating Tablets 20 mg
Acid Reducer

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect



Strawberry
Flavor



Actual
size

Melts in your mouth
Dissolves without water

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box

- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight ■ Quit smoking



STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box

1Y955 FA C4-L 092618

<p>Drug Facts</p> <p>Active ingredient (in each tablet)</p> <p>Oneprazole 20 mg. _____ Acid reducer</p> <p>Purpose</p> <p>_____ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Use</p> <p>■ treats frequent heartburn (occurs 2 or more days a week)</p> <p>■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p>	<p>Warnings</p> <p>Allergy alert: do not use if you are allergic to oneprazole</p> <p>Do not use if you have:</p> <ul style="list-style-type: none"> ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain <p>These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have:</p> <ul style="list-style-type: none"> ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain <p>Ask a doctor or pharmacist before use if you are taking a prescription drug.</p> <p>Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if:</p> <ul style="list-style-type: none"> ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact ►</p>	<p>Questions or comments? 1-800-771-9260, weekdays 7:30 AM to 5:00 PM EST</p>	<p>Other information</p> <ul style="list-style-type: none"> ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture <p>Inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscopolone, ferric oxide, flavor, hypromellose, hypromellose phthalate, maize methylacrylate, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium lauryl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate</p>	<p>Directions</p> <ul style="list-style-type: none"> ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours <p>14-Day Course of Treatment</p> <ul style="list-style-type: none"> ■ take 1 tablet before eating in the morning ■ do not crush or chew tablets ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water. ■ take every day for 14 tablets a day ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ do not take this medicine with alcohol <p>Repeated 14-Day Course (if needed)</p> <ul style="list-style-type: none"> ■ you may repeat a 14-day course every 4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition. 	<p>Keep out of reach of children. In case of overdose, get medical help or contact ►</p>	<p>Questions or comments? 1-800-771-9260, weekdays 7:30 AM to 5:00 PM EST</p>	<p>Drug Facts (continued)</p>	<p>Directions</p> <ul style="list-style-type: none"> ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours <p>14-Day Course of Treatment</p> <ul style="list-style-type: none"> ■ take 1 tablet before eating in the morning ■ do not crush or chew tablets ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water. ■ take every day for 14 tablets a day ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ do not take this medicine with alcohol <p>Repeated 14-Day Course (if needed)</p> <ul style="list-style-type: none"> ■ you may repeat a 14-day course every 4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition. 	<p>Other information</p> <ul style="list-style-type: none"> ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture <p>Inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscopolone, ferric oxide, flavor, hypromellose, hypromellose phthalate, maize methylacrylate, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium lauryl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate</p>	
--	--	--	--	---	---	--	--------------------------------------	---	--	--

A barcode consisting of vertical black bars of varying widths on a white background. Below the bars are several groups of numbers: "0", "00000", "00000", and "0".

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

**Safety Feature –
Do not use if printed tablet
blister unit is open or torn.**



NDC 64239-XXX-03



Value 3 Pack



Strawberry
Flavor

Melts in your mouth
Dissolves without water

Actual
size

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the “Drug Facts” box

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Omeprazole

Delayed Release
Orally Disintegrating Tablets 20 mg

Active ingredient (in each tablet) _____ Acid reducer

Purpose

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- chest or shoulder pain; or shortness of breath, sweating, pain spreading to arms, neck, or shoulders; or lightheadedness
- frequent chest pain

These may be signs of a serious condition. See your doctor.

Ask a doctor or pharmacist before use if you are taking a prescription drug.

Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you get diarrhea
- you develop a rash or joint pain

Keep out of reach of children. In case of overdose, get medical help or contact **Poison Control Center** right away (1-800-222-1222).


1Y930 FA C4-D 092718

**Location for
lot number and
expiration date**



- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Safety Feature –
Do not use if printed tablet
blister unit is open or torn.



NDC 64239-XXX-02

Treats FREQUENT Heartburn!



MELTech™
Melts In Your Mouth

Compare to Prilosec OTC®*

100% More Free!
14 Free Tablets!

Omeprazole

**Delayed Release
Orally Disintegrating Tablets 20 mg
Acid Reducer**

28 TABLETS

Two 14-day courses of treatment
May take 1 to 4 days for full effect

Melts in your mouth
Dissolves without water




Strawberry Flavor



Actual
size

Hairlines extend to within two spaces of the “Drug Facts” box



Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

Omeprazole

Delayed Release
Orally Disintegrating Tablets 20 mg

inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscopollose, ferulic oxide, hydroxypropylmethylcellulose, hypromellose, hypromellose phthalate, maize maltodextrinate, mannitol, polypropylene glycol, silicon dioxide, polyvinyl stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F); keep product out of high heat and moisture

- **For adults:** this product is to be used once a day (every 24 hours), every day for 14 days
- **It may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours**
- 14-Day Course of Treatment**
- **Take 1 tablet before eating in the morning**
- **Do not crush or chew tablets**
- **Place the tablet on tongue, tablet disintegrates, with or without water.**
- **The tablet can also be swallowed whole with water.**
- **Take every day for 14 days**
- **Do not take more than 1 tablet a day**
- **Do not use for more than 14 days unless directed by your doctor**
- **Do not take this medicine with alcohol**
- Repeated 14-Day Course (if needed)**
- **You may repeat a 14-day course every 4 months**
- **Do not take for more than 14 days or more often than every 4 months unless directed by your doctor**
- **Children under 18 years of age:** a doctor before use. **Heartburn in children may sometimes be caused by a serious condition.**

Drug Facts (continued)

► **If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Stop use and ask a doctor if:

- your headache continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

Ask a doctor or pharmacist before use if you are taking a prescription drug.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

These may be signs of a serious condition. See your doctor

- frequent chest pain
- arms, neck or shoulders; or lightheadedness
- chest or shoulder pain with shortness of breath; sweating; pain spreading to
- head, arm with light-headedness, sweating or dizziness
- trouble or pain swallowing food, vomiting with blood, or bloody or black stools

Warnings
Allergy alert: do not use if you are allergic to omeprazole

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect


Active ingredient (in each tablet) Omeprazole 20 mg
Acid reducer

Drug Facts

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Safety Feature –
Do not use if printed tablet
blister unit is open or torn.



NDC 64239-XXX-02

Treats *FREQUENT* Heartburn!



MELTech™
Melts In Your Mouth

*Compare to Prilosec OTC®**

Bonus Size!
14 Free

Omeprazole

**Delayed Release
Orally Disintegrating Tablets 20 mg
Acid Reducer**

28 TABLETS

Two 14-day courses of treatment
May take 1 to 4 days for full effect

Melts in your mouth
Dissolves without water



Strawberry Flavor



Actual
size

STANDARD FORMAT

TYPE SIZES:

TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed

HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed

TEXT: 6 pt Helvetica Neue LT Std 57 Condensed

LEADING: 6.5 pt

BULLETS: 5 pt S

TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed

HEAVY LINES: 1.5 pt

HAIRLINES: 0.5 pt

Hairlines extend to within two spaces of the “Drug Facts” box

TYPE SIZES:

HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed

LEADING: 6.5 pt

HEAVY LINES: 1.5 pt

Hairlines extend to within two spaces of the “Drug Facts” box

1Y955 FA C4-E 092618

Omeprazole

Delayed Release
Orally Disintegrating Tablets 20 mg

<p>Drug Facts (continued)</p> <p>Directions</p> <ul style="list-style-type: none"> For adults: 18 years of age and older It may take 1 to 4 days (every 24 hours), every day for 14 days This product is to be used once a day (every 24 hours), every day for 14 days It may take 1 to 4 full effect, some people get complete relief or symptoms within 24 hours Take 1 tablet before eating in the morning Do not crush or chew tablets Place the tablet on tongue; tablet disintegrates, with or without water. The tablet can also be swallowed whole with water. Take every day for 14 days Do not take more than 1 tablet a day Do not use for more than 14 days unless directed by your doctor Do not take this medicine with alcohol <p>Repeated 14-Day Course (if needed)</p> <ul style="list-style-type: none"> You may repeat a 14-day course every 4 months Do not take for more than 14 days or more often than every 4 months unless directed by a doctor <ul style="list-style-type: none"> Children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition. <p>Other information</p> <ul style="list-style-type: none"> Read the directions and warnings before use Keep the cap on. It contains important information. Store at 20-25°C (68-77°F). Keep product out of high heat and moisture <p>Inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscopolone, ferric oxide, flavor, hypromellose, hypromellose phthalate, maize methylcellulose, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, stearate, sodium stearoyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate</p>	<p>Questions or comments? 1-800-719-9260: weekdays 7:30 AM to 5:00 PM EST</p>	<p>Drug Facts</p> <p>Active ingredient (in each tablet)</p> <p>Omeprazole 20 mg _____ Acid reducer</p> <p>Purpose</p> <p>For full effect</p> <ul style="list-style-type: none"> Not intended for immediate relief of heartburn; this drug may take 1 to 4 days Treats frequent heartburn (occurs 2 or more days a week) <p>Use</p> <ul style="list-style-type: none"> For full effect Not intended for immediate relief of heartburn; this drug may take 1 to 4 days <p>Warnings</p> <p>Warning alert: do not use if you are allergic to omeprazole</p> <p>Do not use if you have:</p> <ul style="list-style-type: none"> Trouble or pain swallowing food, vomiting with blood, or bloody or black stools Heartburn with lightheadedness, sweating or dizziness Chest or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness Frequent chest pain <p>Ask a doctor before use if you have:</p> <ul style="list-style-type: none"> Had heartburn over 3 months. This may be a sign of a more serious condition. Frequent wheezing, particularly with heartburn Unexplained weight loss Nausea or vomiting Stomach pain <p>Ask a doctor or pharmacist before use if you are taking a prescription drug.</p> <p>Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if:</p> <ul style="list-style-type: none"> Your heartburn continues or worsens You need to take more than 1 course of treatment every 4 months You get diarrhea You develop a rash or joint pain <p>If pregnant or breast-feeding: ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact Poison Control Center right away (toll free 1-800-222-1222).</p>
---	--	---

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Safety Feature –
Do not use if printed tablet
blister unit is open or torn.

NDC 64239-XXX-03

Treats *FREQUENT* Heartburn!



Compare to Prilosec OTC®*

Bonus Pack: 28 + 14 Free

Omeprazole

Delayed Release
Orally Disintegrating Tablets 20 mg
Acid Reducer

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect

Melts in your mouth
Dissolves without water



Strawberry
Flavour



Actual
size

STANDARD FORMAT

TYPE SIZES:

TITLE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed

HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed

TEXT: 6 pt Helvetica Neue LT Std 57 Condensed

LEADING: 6.5 pt

BULLETS: 5 pt S

TELEPHONE #: 6 pt He

HEAVY LINES: 1.5 pt

HAIRLINES: 0.5 pt

Airlines extend t

Headlines extend to within two spaces of the Drug Facts box.

1Y955 FA C4-F 092618

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Omeprazole 20 mg.....Acid reducer</p> <p>Purpose Acid reducer</p>	<p>Use</p> <p>• treats frequent heartburn (occurs 2 or more days a week)</p> <p>• not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p>	<p>Warnings</p> <p>Allergy alert: do not use if you are allergic to omeprazole</p> <p>Do not use if you have:</p> <ul style="list-style-type: none"> • trouble or pain swallowing food, vomiting with blood, or bloody or black stools • heartburn with lightheadedness, sweating or dizziness • chest or shoulder pain with shortness of breath, sweating, or pain spreading to arms, neck or shoulder; or lightheadedness <p>Frequent chest pain</p> <p>These may be signs of a serious condition. See your doctor.</p>	<p>Ask a doctor before use if you have:</p> <p>• heartburn over 3 months. This may be a sign of a more serious condition.</p> <p>• frequent wheezing, particularly with heartburn</p> <p>• unexplained weight loss</p> <p>• nausea or vomiting</p>	<p>Ask a doctor or pharmacist before use if you are taking a prescription drug.</p> <p>Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if:</p> <ul style="list-style-type: none"> • your heartburn continues or worsens • you need to take this product for more than 14 days • you need to take more than 1 course of treatment every 4 months • you get diarrhea • you get dizziness • you develop a rash or joint pain 	<p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact ► 500-PM EST</p>	<p>Drug Facts (continued)</p> <p>Directions</p> <ul style="list-style-type: none"> • for adults 18 years of age and older • this product is to be used once a day (every 24 hours), every day for 14 days • it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours <p>14-Day Course of Treatment</p> <ul style="list-style-type: none"> • take 1 tablet before eating in the morning • do not crush or chew tablets • place the tablet on tongue; tablet disintegrates, with or without water. • take every day for 14 days • do not take more than 1 tablet a day • do not use for more than 14 days unless directed by your doctor • do not take this medicine with alcohol <p>Repeated 14-Day Course (if needed)</p> <ul style="list-style-type: none"> • you may repeat a 14-day course every 4 months • do not take for more than 14 days or more often than every 4 months unless directed by a doctor <p>• children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p>	<p>Other information</p> <ul style="list-style-type: none"> • keep the carton. It contains important information. • store at 20-25°C (68-77°F); keep product out of high heat and moisture <p>Inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscopolone, ferric oxide, flavoc, hypromellose, hypromellose phthalate, magnesium stearate, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate</p>	<p>Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST</p>	<p>Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST</p>	<p>Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST</p>	<p>Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST</p>	<p>Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST</p>	<p>Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST</p>	<p>Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST</p>	<p>Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST</p>
--	--	---	---	--	---	--	--	--	--	--	--	--	--	--	--

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

**Safety Feature –
Do not use if printed tablet
blister unit is open or torn.**

NDC 64239-XXX-03



Compare to Prilosec OTC®*

Bonus Size!
14 Tablets Free

**Delayed Release
Orally Disintegrating Tablets 20 mg
Acid Reducer**



Strawberry Flavor

Three 14-day courses of treatment
May take 1 to 4 days for full effect

Melts in your mouth
Dissolves without water

Actual
size

TYPE SIZES:

TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) **8 pt Helvetica Neue LT Std 57 Condensed**

HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed

TEXT: 6 pt Helvetica Neue LT Std 57 Condensed

LEADING: 6.5 pt

BULLETS: 5 pt Square

TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed

HEAVY LINES: 1.5 pt

HAIRLINES: 0.5 pt

Hairlines extend to within two spaces of the “Drug Facts” box


010760 B-40 A1 CCE 11

[illegible]

A barcode consisting of vertical black bars of varying widths on a white background. Below the bars are the numbers "0000000000000000".

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

**Safety Feature –
Do not use if printed tablet
blister unit is open or torn.**



MELTech™
Melts In Your Mouth



Bonus Size!
14 Free



Strawberry
Flavor

Melts in your mouth
Dissolves without water

Three 14-day courses of treatment
May take 1 to 4 days for full effect



Actual size

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the “Drug Facts” box

1Y955 FA C4-J 092618

Omeprazole

Delayed Release
Orally Disintegrating Tablets 20 mg

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

**Safety Feature –
Do not use if printed tablet
blister unit is open or torn.**

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued) **8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique**
 (continued) **8 pt Helvetica Neue LT Std 57 Condensed**
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 1.5 pt Square
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box

This label may not be the latest approved by FDA.
current labeling information, please visit <https://www.fda.gov/drugs>

FRONT PANEL

peel off
coupon

Instant Redeemable Coupon - Expires XX-XX-XX

SAVE \$X NOW!

On This Purchase

MELTech™
Melts In Your Mouth

Compare to Prilosec OTC®*

NDC 64239-XXX-01

Treats **FREQUENT** Heartburn!

 **24 HR**

Omeprazole

Delayed Release

Orally Disintegrating Tablets 20 mg

Acid Reducer

14 TABLETS

One 14-day course of treatment
May take 1 to 4 days for full effect



Strawberry
Flavor

Melts in your mouth
Dissolves without water



1Y974 FA CP3 082318

BACK PANEL

SAVE \$X NOW!

MANUFACTURER'S COUPON EXPIRATION DATE XX/XX/XXXX

SAVE \$X^{XX}

on your purchase of
14 COUNT SIZE
"Store Brand" Omeprazole
Delayed Release Orally Disintegrating Tablets 20 mg

CONSUMER: Only one coupon is redeemable per purchase. **RETAILER:** This coupon will be redeemable for face value on specified products (plus a handling fee) if submitted in compliance with the terms of this offer. Cash value 1/100th of one cent. Limit one coupon per customer. **Do Not Double.** Coupon is non-reproducible. Coupons cannot be combined. Mail properly handled coupons to: L. Perrigo Company (#1565), NCH Marketing, PO Box 880001, El Paso, TX 88588-0001

COUPON NOT SUBJECT TO DOUBLING

Customer pays any sales tax.



Omeprazole

Delayed Release
Orally Disintegrating Tablets 20 mg

14 TABLETS

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Reference ID: 4381841

This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drugsat>

FRONT PANEL

peel off
coupon

Instant Redeemable Coupon - Expires XX-XX-XX

SAVE \$X NOW!

On This Purchase

MELTech™
Melts In Your Mouth

Compare to Prilosec OTC®*

NDC 64239-XXX-02

Treats **FREQUENT** Heartburn!

 **24 HR**

Omeprazole

Delayed Release

Orally Disintegrating Tablets 20 mg

Acid Reducer

28 TABLETS

Two 14-day courses of treatment
May take 1 to 4 days for full effect

Melts in your mouth
Dissolves without water



Strawberry
Flavor



Actual
size

1Y930 FA CP3 082318

BACK PANEL

SAVE \$X NOW!

MANUFACTURER'S COUPON EXPIRATION DATE XX/XX/XXXX

SAVE \$X^{XX}

on your purchase of
28 COUNT SIZE

"Store Brand" Omeprazole

Delayed Release Orally Disintegrating Tablets 20 mg

CONSUMER: Only one coupon is redeemable per purchase. **RETAILER:** This coupon will be redeemable for face value on specified products (plus a handling fee) if submitted in compliance with the terms of this offer. Cash value 1/100th of one cent. Limit one coupon per customer. **Do Not Double.** Coupon is non-reproducible. Coupons cannot be combined. Mail properly handled coupons to: L. Perrigo Company (#1565), NCH Marketing, PO Box 880001, El Paso, TX 88588-0001

COUPON NOT SUBJECT TO DOUBLING

Customer pays any sales tax.



Omeprazole

Delayed Release
Orally Disintegrating Tablets 20 mg

28 TABLETS

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Reference ID: 4381841

This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drugsatfda>

FRONT PANEL

peel off
coupon

Instant Redeemable Coupon - Expires XX-XX-XX

SAVE \$X NOW!

On This Purchase

MELTech™
Melts In Your Mouth

Compare to Prilosec OTC®*

NDC 64239-XXX-03

Treats **FREQUENT** Heartburn!

 **24 HR**

Omeprazole

Delayed Release

Orally Disintegrating Tablets 20 mg

Acid Reducer

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect



Strawberry
Flavor

Melts in your mouth
Dissolves without water



Actual
size

1Y955 FA CP3 082318

BACK PANEL

SAVE \$X NOW!

MANUFACTURER'S COUPON EXPIRATION DATE XX/XX/XXXX

SAVE \$X^{XX}

on your purchase of
42 COUNT SIZE
"Store Brand" Omeprazole
Delayed Release Orally Disintegrating Tablets 20 mg

**COUPON NOT SUBJECT
TO DOUBLING**

Customer pays any sales tax.

CONSUMER: Only one coupon is redeemable per purchase. RETAILER: This coupon will be redeemable for face value on specified products (plus a handling fee) if submitted in compliance with the terms of this offer. Cash value 1/100th of one cent. Limit one coupon per customer. Do Not Double. Coupon is non-reproducible. Coupons cannot be combined. Mail properly handled coupons to: L. Perrigo Company (#1565), NCH Marketing, PO Box 880001, El Paso, TX 88588-0001



Omeprazole

Delayed Release
Orally Disintegrating Tablets 20 mg

42 TABLETS

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Reference ID: 4381841

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

KAREN M MAHONEY
01/28/2019 11:50:03 AM